




Dental Science, Materials and Technology

DENTURES

*Types,
Benefits and Potential
Complications*




N
o
v
a

B
i
o
m
e
d
i
c
a
l



TIBOR KAROLY FÁBIÁN
PAL FEJERDY
PETER HERMANN
EDITORS

NOVA



VISIT...

LANZAROTE
Caliente.COM

DENTAL SCIENCE, MATERIALS AND TECHNOLOGY

DENTURES

TYPES, BENEFITS AND

POTENTIAL COMPLICATIONS

No part of this digital document may be reproduced, stored in a retrieval system or transmitted in any form or by any means. The publisher has taken reasonable care in the preparation of this digital document, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained herein. This digital document is sold with the clear understanding that the publisher is not engaged in rendering legal, medical or any other professional services.

DENTAL SCIENCE, MATERIALS AND TECHNOLOGY

Additional books in this series can be found on Nova's website
under the Series tab.

Additional E-books in this series can be found on Nova's website
under the E-book tab.

DENTAL SCIENCE, MATERIALS AND TECHNOLOGY

DENTURES

**TYPES, BENEFITS AND
POTENTIAL COMPLICATIONS**

TIBOR KAROLY FÁBIÁN
PAL FEJERDY
AND
PETER HERMANN
EDITORS



Nova Science Publishers, Inc.
New York

Copyright © 2012 by Nova Science Publishers, Inc.

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic, tape, mechanical photocopying, recording or otherwise without the written permission of the Publisher.

For permission to use material from this book please contact us:

Telephone 631-231-7269; Fax 631-231-8175

Web Site: <http://www.novapublishers.com>

NOTICE TO THE READER

The Publisher has taken reasonable care in the preparation of this book, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained in this book. The Publisher shall not be liable for any special, consequential, or exemplary damages resulting, in whole or in part, from the readers' use of, or reliance upon, this material. Any parts of this book based on government reports are so indicated and copyright is claimed for those parts to the extent applicable to compilations of such works.

Independent verification should be sought for any data, advice or recommendations contained in this book. In addition, no responsibility is assumed by the publisher for any injury and/or damage to persons or property arising from any methods, products, instructions, ideas or otherwise contained in this publication.

This publication is designed to provide accurate and authoritative information with regard to the subject matter covered herein. It is sold with the clear understanding that the Publisher is not engaged in rendering legal or any other professional services. If legal or any other expert assistance is required, the services of a competent person should be sought. FROM A DECLARATION OF PARTICIPANTS JOINTLY ADOPTED BY A COMMITTEE OF THE AMERICAN BAR ASSOCIATION AND A COMMITTEE OF PUBLISHERS.

Additional color graphics may be available in the e-book version of this book.

Library of Congress Cataloging-in-Publication Data

ISBN: 978-1-61942-288-9 (eBook)

Published by Nova Science Publishers, Inc. † New York

Contents

Preface		vii
Chapter I	The Prosthetic Classification of Partially Edentulous Dental Arches and Its Use for Treatment Planning <i>Tibor Fábián, Péter Hermann and Pál Fejérdy</i>	1
Chapter II	Orthodontics for Patient of Prosthetic Dentistry. Possibilities and Complications of Psychosomatic Origin <i>Tibor Károly Fábián, Anita Beck, Boglárka Rencz, Pál Fejérdy, Péter Hermann and Gábor Fábián</i>	15
Chapter III	Effects of Wearing Partial Denture Prosthesis on Jaw Functions and Associated Prefrontal Cortex Activity <i>Noriyuki Narita</i>	27
Chapter IV	Inlay Retained Fixed Partial Dentures <i>Sherif Adel Mohsen</i>	45
Chapter V	Removable Partial Dentures with Double Crowns - Biological and Prophylactic Value and Potential Complications <i>Ivica Stančić and Aleksandra Jelenković Popovac</i>	61
Chapter VI	Systemic Changes of Interest for Rehabilitation with Complete Dentures: Manufacture and Maintenance <i>Leandro Dorigan de Macedo, Soraia Marangoni, Maria do Carmo Favarin de Macedo, Hilton Marcos Alves Ricz and Luciana Assirati Casemiro</i>	85
Chapter VII	Laboratory Improvement of the Clinical Performance of Complete Dentures – A Review <i>Luciana Assirati Casemiro, Soraia Marangoni, André Marcelo Peruchi Minto and Leandro de Dorigan de Macedo</i>	115

Chapter VIII	Alternatives to Removable Complete Dentures: Viability for Public Health Systems in Developing Countries <i>Mara Rodrigues Martins, Wellington Márcio dos Santos Rocha, Débora Emrich Magalhães, Fernanda Piana Santos Lima de Oliveira, Virgínia Sanábio Vilela, Marcos Azeredo Furquim Werneck and Efigênia Ferreira e Ferreira</i>	129
Chapter IX	Denture Adhesives – A Boon or a Bane <i>Puja Chowdhry Malhotra</i>	143
Chapter X	Changes in the Mycological Ecology of Dentures <i>Carlos Madrid, Jacques Bille and Marcelo Abarca</i>	151
Chapter XI	Methods for the Treatment of Denture Induced Psychogenic Symptoms <i>Anita Beck, Gábor Varga, Péter Hermann, Gábor Fábíán, Pál Fejérdy and Tibor Károly Fábíán</i>	165
Chapter XII	Psychogenic Complications of Making Dentures. Theoretical Background, Prevention and Treatment Possibilities <i>Tibor Károly Fábíán, Anita Beck, Laura Gótai, Péter Hermann and Pál Fejérdy</i>	199
Chapter XIII	Pulsed Electromagnetic Field (PEMF) Therapy for Peri-Implant Healing and Bone Remodeling <i>Tibor Károly Fábíán, Anita Beck, Gábor Fábíán, Pál Fejérdy and Péter Hermann</i>	243
Index		255

Preface

This work is an edited collection of miscellaneous chapters from the huge field of prosthetic dentistry, about dentures, and their types, benefits and potential complications. Besides most frequently discussed related topics of the field, this edited collection also includes some chapters dedicated to rarely reviewed research areas like preprosthetic orthodontics, brain correlates of wearing denture, psychosomatic aspects of prosthetic dentistry, denture related microbiology and the use of mind-body therapies, physiotherapies as well as complementary and alternative approaches for prosthetic dentistry, which may supply a great want. This book gathers and presents data on the leading edge of prosthetic dentistry.

Chapter I – Major goal of the “prosthetic classification” is to enhance treatment planning and denture designing procedures. With regard to this aspect partially edentulous arches can be rendered to classes as follows: *Class 1A*: one or more primary fulcrum lines (straight line drawn between two teeth adjacent to the edentulous space) may exist; but none of the primary fulcrum lines is going to become a factual rotation axis when a denture is inserted. No force of rotation will be generated and the denture may not settle. In this class of arches fixed dentures are recommended. *Class 1B*: one or more fulcrum line(s) may exist, which may become a factual rotation axis when a denture is inserted. The rotational forces, however, will not be very powerful, so that the settling of the denture can be equalized (i.e. compensated due to multiple abutments). In this class of arches fixed dentures may be recommended. *Class 2A*: only one primary fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this factual rotation axis the denture may settle in one direction. In this class of arches RPD is recommended and the baseplate of RPD can be moderately reduced. *Class 2A/1*: only one fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. This class is a special case (subclass) of class 2A characterized by very few (maximum one or two) remaining teeth. In this class of arches RPD is recommended and the baseplate of RPD can *not* be reduced. *Class 2B*: Several primary fulcrum lines may exist, one of which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. In this class of arches RPD (combined with or without fixed dentures) are recommended and the baseplate of RPD can be moderately reduced. *Class 3*: one or more primary fulcrum line(s) may exist; from which one or more may become a factual rotation axis when a denture is inserted. Rotating around

this/these rotation axis the denture will tip upward on one side and downward on the other side (i.e. two-directional rotary movements occur). In this class of arches RPD is recommended, the baseplate of RPD can not be reduced and, *resilient* type telescopes (i.e. delayed type dental support) should be used.

Chapter II – Various dental specialties have made such enormous advances that it has become impossible to absorb all that is new in all dental areas. Therefore a collaborative effort between orthodontist and prosthodontist is often necessary to achieve an optimal treatment result. Prosthodontists and restorative dentists rely on orthodontics as a means of providing more comprehensive care to their patients. Such collaboration is not only desirable, but mandatory if dentists are to provide highest level care for their patients. Preprosthetic orthodontics gains significance in both esthetics and functionality of the denture being under preparation. Therefore, an interdisciplinary approach may also be cost-effective for both patient and dentist from the standpoint of producing more stable and durable restorations of high esthetics. Orthodontics may contribute to the success of cases undergoing restorative dental procedures where preliminary minor tooth movement is necessary, to correct local malpositions and/or to move abutment teeth with simple appliances. Success of implant placement may also be improved via orthodontic forced extrusion of periodontally compromised teeth. Collaboration of orthodontist and prosthodontist may be even more demanding when correction of primary orthodontic anomalies would be needed. No wonder that, the integration of orthodontics and prosthetic dentistry into a multidisciplinary rehabilitation is often indispensable for a proper dental care of the patients. A proper plaque control is also essential for a successful treatment in all cases. Along with mechanical and biological principles, unique psychological and social aspects of the combined orthodontic and prosthetic treatment should also be considered for a successful therapy. In this chapter the important role of orthodontics and its contribution to the success of prosthetic dentistry for adult patients is discussed.

Chapter III – *Purpose*: Denture wearing is considered to improve chewing ability, chewing comfort, and quality of life in elderly individuals. In an attempt to clarify the effects of prosthodontic treatment on activation of the prefrontal cortex involved in higher cognitive functions, the authors investigated the efficacy of wearing partial denture from the aspects of jaw functions such as muscle and movement activities and associated prefrontal cortex activity. *Methods*: Twelve partially edentulous patients were enlisted as subjects. Functional near-infrared spectroscopy was used to detect changes of hemoglobin concentration in the prefrontal cortex area, while jaw muscle EMG and jaw motion activities were simultaneously examined. Masticatory scores were also determined to evaluate the subjective awareness of chewing ability and occlusal contact features were examined using a Dental Prescale Occluzer System. Statistical comparisons were performed to compare between data obtained while wearing and not wearing a partial denture prosthesis. *Results*: Significant increases were seen in chewing-related jaw functions such as jaw muscle EMG and jaw motion activities, as well as prefrontal cortex activity associated with wearing a partial denture prosthesis as compared with the not wearing condition. Furthermore, occlusal contact and masticatory scores were also significantly increased by wearing a partial denture prosthesis. *Conclusions*: Occlusal reconstruction obtained by wearing a partial denture prosthesis improved prefrontal cortex activity and the awareness of chewing ability, as well as jaw functions, as shown by increased jaw muscle EMG and jaw motion activities. From these findings, the authors concluded that appropriate denture wearing might have an impact on not

only jaw functions, but also cognitive functions closely related to masticatory performance. The relationship between masticatory function and cognitive ability is discussed from the standpoint of prosthodontic treatment in this chapter.

Chapter IV – Patients often refuse these options since the reasons of high cost and/or surgical intervention. The full coverage retainers for tooth supported fixed partial denture is a conventional approach to replace one or several missing teeth. Two types of full coverage retainers FPDs are used in dental clinics: porcelain fused to metal and all ceramic. Porcelain fused to metal provides great fracture strength and relatively good long-term clinical success. One of significant disadvantages of the porcelain fused to metal, as well as modern all ceramic FPDs is that large amount of sound tooth substance of the abutments needs be removed in FPD fabrication. An inlay-retained FPD (IRFPD) is, however, a less-invasive treatment modality and a more conservative option for restoration of damaged teeth, because it requires minimal tooth reduction, preserves healthy tooth structure, and maintains the periodontal tissue's integrity. Inlay-retained FPDs are, therefore, alternatives to both anterior and posterior complete coverage conventional restorations. Inlay retained FPDs can be constructed by using dental alloys, ceramic materials, and fiber-reinforced composite. Several types of preparations for inlay retained FPDs are used: Occluso – proximal inlay, Tub-shaped , Proximal box, Rest seat on the occlusal surface, Lingual tooth reduction, and Retentive-slot preparations. The size of these preparation features depends on the size of the tooth. Several researches were carried out about the effect of various designs, materials, connectors size on the fracture resistance, marginal integrity and retention of inlay retained FPDs.

Chapter V – Correctly indicated and accurately made, partial denture with double crowns is a dental restoration of the highest biological, prophylactic and aesthetic values. However, insufficient knowledge of double crowns systems, as well as errors and omissions in the clinical and laboratory work invalidate this type of restoration and can have negative effects on the supporting tissues. Double crowns may come in form of telescope crowns and conical crowns. Double crowns as connecting elements should provide: retention, stabilization, physiological transmission of occlusal loads through the abutment teeth as well as the prophylactic value, comfort for the patient, durability and aesthetics. In order to achieve these qualities, it is necessary, among other things: to make proper indication, to execute adequate tooth preparation (grinding), to make the choice of materials for the construction, to establish proper connection of the outer crown and the metal framework and to choose a suitable aesthetic material for veneering. It is necessary to understand the mechanism of double crowns retention so that the optimal value of the retention force which should be 5-9 N per each patrix can be established. Insufficient retention force will reduce the comfort for the patient because would make the separation too easy, whereas an excessive force may negatively affect the periodontium of the abutment teeth. Retention force and durability of restorations with telescope and conical crowns depend also on the characteristics of the material they are made of. The most frequently used are gold and palladium-silver alloys. In recent years, titanium and the combination of ceramic and galvanized gold are used. Prophylactic effect of double crowns is also provided by a proper planning that will enable the physiological transfer of the forces on the abutment teeth and the health of free gingiva. A special challenge is to plan the connection of outer crown with the metal framework. The best results showed optimized approximal extensions. Double crown production demands experience of a clinician and dental technician in the performance of clinical and laboratory phases. A removal of sufficient tooth tissue during grinding in order to avoid getting

oversized artificial teeth and a formation of the appropriate demarcation line which will provide sufficient stability of the system in the gingival area represent a particular challenge. Errors in laboratory work can be avoided with a good knowledge of the milling proceedings, a selection of a suitable material for the external crowns modeling and acquisition of precision castings.

Chapter VI – The aging population has profoundly transformed the profile of patients seeking oral rehabilitation. Care delivery to these patients, who present with multiple comorbidities and are usually being treated with various drugs, is becoming increasingly more frequent. Systemic changes prompted by the use of many medications may impose limitations to clinical practice with regard to indication, planning, materials, preparation techniques, and maintenance of prosthetic rehabilitation. Unfortunately, the dental literature is lacking in scientific papers reporting oral manifestations of systemic diseases, as well as in works describing the type of assistance that is necessary to treat such patients. The reported findings are limited to the description of techniques, which are sometimes adapted, for complete denture treatment in patients with neurological diseases, such as Alzheimer's and Parkinson's syndrome, and burning mouth in cases of allergy to dental materials. Several systemic conditions and different treatments should be taken into account when it comes to the correct time for prosthetic rehabilitation completion, the materials and technique to be used, and the maintenance procedures to be employed. It is known that in the case of patients with severe immunosuppression, whose origin can be constitutional (senile), pathological (hematological, immune, or endocrine), or prompted by medication (use of immunosuppressants, corticosteroids, chemotherapy, among others), there is higher incidence of oral infections by fungi, especially opportunistic infections such as aspergillosis and mucormycosis, whose incidence may increase due to badly fitted and poorly cleaned dentures. Another group of autoimmune diseases, the so-called pemphigus vulgaris, may have repercussions in the oral mucosa. This disease predisposes the oral cavity to serious injury by lack of adherence of epithelial tissue and by infection due to the constant utilization of immunosuppressants, making the use of dentures very difficult. In addition, patients with renal failure and severe liver disease may present changes in both oral mucosa and salivary glands, which demands manufacture and use of dentures. Treatments such as radiotherapy and the chronic use of bisphosphonates imply in severe changes in both the bone and soft tissues. The common denominator of all these diseases or therapeutic protocols is that they have temporary or permanent contraindications for rehabilitation with conventional dentures or implant overdentures. Thus, on the basis of literature and a 10-year clinical group experience in a dental hospital service, this book chapter aims to describe major diseases and drug therapies that directly or indirectly reflect on the oral cavity, including the necessary adjustments in materials, manufacturing techniques, and maintenance of complete dentures.

Chapter VII – Laboratory processing techniques have been refined in an attempt to achieve complete dentures with better clinical performance. Regardless of the type of denture base material (microwave-, light-, or heat-polymerized polymethyl methacrylate, for example), materials as well as handling and polymerization methods have been continuously modified with a view to obtaining more resistant dentures. In an attempt to improve the mechanical properties of base materials and to try to overcome the low impact and flexural strength of currently available dentures, which often results in clinical failure, various laboratory treatments have been proposed. The incorporation of reinforcing structures such as glass fiber, carbon fiber, Kevlar, aramid, and fiber-reinforced composite, which can be found

in different sizes and shapes and are sometimes impregnated with substances such as silanes, has been successfully tested in some cases. However, incorporation of such compounds can culminate in undesirable effects such as larger amount of residual monomer, difficult polishing, and unsatisfactory aesthetics. Metal wires have also been used over time for this purpose, as well as substances such as methyl acetate and methyl formate. The increased bond strength between artificial teeth and the denture base material has also been relentlessly pursued. New positive results are being obtained by means of techniques that promote interpenetration of the polymer networks of the two materials and make use of substances such as dichloromethane. To improve the biological properties of dentures, simple procedures such as immersion in water for defined time periods has been evaluated, in order to reduce their toxicity. Studies have shown that dibenzoyl peroxide, used as initiator in polymethyl methacrylate resin and associated with allergic processes, is released from conventional dentures after polymerization, but this event can be substantially reduced with completion of an additional polymerization cycle. Still, in order to reduce dentures contamination, several antimicrobial agents such as silver have been incorporated into the base material, which has led to positive effects with regard to their antimicrobial action. However, decrease in flexural and impact strength has sometimes been observed. In this context, this book chapter intends to update the reader on materials and laboratory techniques that can optimize the physical, mechanical, and biological properties of dentures that are considered to be the most relevant to the clinical performance of complete dentures.

Chapter VIII – The global population is ageing at an unprecedented rate. The result of this process has produced an important change in the healthcare system, with the cure of acute diseases giving way to the treatment of chronic illnesses. The impact of this demographic change is of considerable concern due to the increase in the cost of healthcare systems, although the current correlation between health costs and the elderly population is weak. In the last ten years, Brazil has made considerable progress in the prevention and control of caries among children. However, the situation among adults and the elderly is among the poorest in the world. However, the rehabilitation of these individuals is hindered by a number of factors. The difficulty of retaining and adapting lower dentures is common and constitutes an obstacle to the complete success of prosthesis use, as reported in a number of studies. Moreover, the lack of information on the part of patients regarding the limitations of conventional complete dentures contributes toward dissatisfaction and the abandonment of complete lower denture use. The aim of this chapter is to carry out a systematic review of methods for the rehabilitation of individuals with an edentulous lower jaw in the search for an alternative solution that is biological, functional and financially viable. This review will consist of the treatment and options for edentulismo including the bone loss risk, the patient satisfaction, the masticatory efficiency, the number of implants and the economic cost/analysis. At the final presents a discussion on clinical decisions in the edentulous rehabilitation.

Chapter IX – Denture Adhesives have always been surrounded by an infinite number of controversies. Prosthodontists and dentists in general have never really accepted the role of denture adhesives in improving denture retention, stability and function. Despite substantial and significant records encouraging the use of adhesives, the dental fraternity by large still considers adhesive use as a poor echo of their treatments and prosthetic proficiencies. However, in view of a longer-living population with an increasing rate of chronic residual ridge resorption and a consequent increase in dentures that are unsteady because of

parameters beyond the control of the patient or dentist, denture adhesives deserve a place in removable prosthodontics as an effective and active aide in denture treatment and denture aftercare. Complete dentures constitute one of the most important treatment options in prosthodontics, more so with an increase in average life expectancy of the individual. However, retention of complete dentures has always posed to be a problem for the dentist and the dental industry. Solutions to the problem, over the years have included overdentures, implants, and denture adhesives. From time immemorial adhesives have been used by denture wearers and advertised commercially, however these dental materials have not been given their due place in prosthetic dentistry. These materials are specially promoted in certain special conditions such as complicated prostheses—obturators, dry mouth, difficult and exacting patients (e.g. Parkinson's disease, Alzheimer's disease), compromised ridges, single complete denture or in public figures. Besides, adhesives are also normally used by denture wearers to psychologically support the patient so as to make the complete dentures more acceptable. Conflicting views still persist. The ancient views regard these materials as a poor replacement for a proper fit and adjustment of dentures. However, existing literature supports the use of denture adhesives i.e. if properly used. Adhesives can be a valuable aid in management of challenging denture patients. Therefore, the following article aims to solve this dilemma regarding the use denture adhesives, oust some of the myths related to adhesives in order to allow good use and avoid misuse of the same.

Chapter X – A wide range of medical devices used in elderly patients shown to support colonization and biofilm formation by *Candida spp.* Among them, more attention must be paid to denture plaque because similar to dental plaque it also serves as a reservoir of potentially infectious microorganisms. Continuous swallowing or aspiration of microorganisms from denture plaque exposes elderly patients to the risk of unexpected infections. The level of oral hygiene in elderly people, especially those living in long-term care institutions, has been reported to be poor, both for dentures and remaining teeth. The upper denture has been suggested to be the major source of infection in long term hospital care patients. It has also been reported that larger quantities of *Candida spp.* reside in the denture base and not in the palate. The most frequent isolated yeast species in dentures is *C. albicans*. Although, recent reports show high proportions of *non-albicans Candida spp.*, thus showing a shift from *C. albicans* towards *non-albicans spp.* This shift has already been reported over the last years with *C. glabrata* emerging as the second most prevalent species found in dentures. Infections related to *C. glabrata* can lead to systemic infections with high mortality rates. The increased inadequate use of topical and systemic antifungals has been proposed to contribute to this fungal profile change. Several studies show a correlation between poor denture hygiene and prevalence of *Candida* species and also a statistically significant relationship between denture related stomatitis (DRS), presence of *Candida* and denture cleanliness. High frequency and quantity of *Candida spp.* detection is sometimes associated with few clinically detectable cases of DRS. Hyphae detection by microscopy is now the gold standard (with the exception of *C. glabrata*) for the diagnosis of oropharyngeal fungal infections (OFI). Presence of oral lesions associated with OFI has been reported only in one third of microscopically affected patients. Frequent use of toothpaste to clean dentures has an influence on dentures surface roughness due to the presence of abrasives. Surface roughness being positively correlated with the rate of fungal colonization of biomaterials, a rougher surface may be a risk factor for microorganism adhesion and biofilm formation. Conclusion: High levels of *Candida spp.* are reported on dentures of elderly patients. A high

frequency of *non-albicans Candida spp.* (particularly *C. glabrata*) has been detected in recent studies. This seems to confirm a change in the mycological ecology involved in yeast denture colonization and DRS. Keywords: *Candida spp.*, dentures, elderly.

Chapter XI – Psychogenic denture intolerance (PDI) patients suffering from several denture induced psychogenic symptoms refuse to accept psychological background of their symptoms frequently. Instead of psychiatrists or psychotherapists, first they visit dentist and insist on the somatic origin of their symptoms. Since most of these patients refuse a referral to psychiatrist and/or psychotherapist, an initial psychosomatic therapy is needed, which is a scope of dental profession's duty. Initiation of a palliative dental care and gradual escalation of any kind of mind-body therapy (as "basic therapeutics" for psychosomatic disorders) are "cornerstones" of such dental psychosomatic interventions. Introduction of any method of complementary/alternative medicine may also be useful especially for the prevention of relapse. To carry out efficient palliative dental care useless repetition of previous dental treatment (or any further invasive dental treatment) should be avoided, certain psychotherapeutic approaches should be introduced and several other treatment modalities like physiotherapy, medication, medicinal herb therapy or acupuncture should also be administered. For mind-body therapies, relaxation, hypnosis, self-hypnosis, meditation, photo-acoustic stimulation or biofeedback may be used. For complementary and alternative medicine prayer, breathing exercises, Tai-Chi and Qigong, Yoga or several other therapies (i.e. sleep deprivation, vigil, fasting therapy etc.) may be considered.

Chapter XII – Psychogenic complications of making dentures as well as consequently appearing psychogenic denture intolerance is a complex and rising problem of dentistry and presents many intricate problems, which are being tackled by various disciplines of both basic and clinical research. Estimations based on the available data and clinical experience indicate that, at least 3-4% of denture wearers suffer from psychogenic symptoms caused by the treatment procedure, insertion or wearing of fixed or removable dentures. No wonder that, there is a high amount of scientific information gathered so far, however data are rather divergent, sometimes even contradictory and there are numerous questions without any available data to answer. Present chapter is primarily dedicated to the clinical aspects of these phenomena, including clinical manifestations, diagnosis, prevention and treatment possibilities. Other relevant subject areas of this chapter include theoretical background and peculiarities of denture-related psychological and psycho-physiological phenomena, background and pathomechanisms of denture induced psychosomatic manifestations, basic principles of communication and patient-nurse-dentist interrelationships. This collection of information helps the reader to be at home in scientific field of denture related psychogenic manifestations.

Chapter XIII – Data in the literature revealed that bone formation around dental implants can be promoted due to PEMF treatment significantly. Since PEMF decreases the rate of residual ridge resorption following tooth extraction, PEMF is likely to improve healing of implants inserted into extraction sites as well. PEMF may also be used advantageously for accelerating bone healing following advanced implant surgery. Although PEMF stimulation during the first two weeks after surgery seems to be the most efficient; late phase bone healing (even four weeks after surgery) may also be accelerated with PEMF administration. Moreover, PEMF stimulation speeds up bone remodeling processes which effect is likely to be a crucial factor of the phenomenon that PEMF accelerates tooth movement (and coupled periodontal processes). PEMF also influence the function of periodontal ligament fibroblasts,

which also may play a significant role in the acceleration of bone remodeling around orthodontically moved teeth as well as around teeth subjected to high mechanical load during mastication (i.e. abutment teeth) or because of parafunction. Although PEMF was found to be beneficial to a wide variety of therapeutic processes also in dentistry; the exact mechanisms by which PEMF affects alveolar bone and other periodontal tissues is far not fully understood yet. However, certain mechanisms are already recognized and the increased amount of knowledge may be utilized also for the clinical practice. In this chapter scientific data of clinical relevance are collected and discussed.

Chapter I

The Prosthetic Classification of Partially Edentulous Dental Arches and Its Use for Treatment Planning

Tibor Fábián*, Péter Hermann and Pál Fejérdy

Semmelweis University Budapest, Faculty of Dentistry,
Clinic of Prosthetic Dentistry, Budapest, Hungary, EU

Abstract

Major goal of the “prosthetic classification” (Fábián and Fejérdy 1979) is to enhance treatment planning and denture designing procedures. With regard to this aspect partially edentulous arches can be rendered to classes as follows:

Class 1A: one or more primary fulcrum lines (straight line drawn between two teeth adjacent to the edentulous space) may exist; but none of the primary fulcrum lines is going to become a factual rotation axis when a denture is inserted. No force of rotation will be generated and the denture may not settle. In this class of arches fixed dentures are recommended.

Class 1B: one or more fulcrum line(s) may exist, which may become a factual rotation axis when a denture is inserted. The rotational forces, however, will not be very powerful, so that the settling of the denture can be equalized (i.e. compensated due to multiple abutments). In this class of arches fixed dentures may be recommended.

Class 2A: only one primary fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this factual rotation axis the denture may settle in one direction. In this class of arches RPD is recommended and the baseplate of RPD can be moderately reduced.

Class 2A/1: only one fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. This class is a special case (subclass) of class 2A characterized by very few

* E-mail address: fab@fok.usn.hu.

(maximum one or two) remaining teeth. In this class of arches RPD is recommended and the baseplate of RPD can *not* be reduced.

Class 2B: Several primary fulcrum lines may exist, one of which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. In this class of arches RPD (combined with or without fixed dentures) are recommended and the baseplate of RPD can be moderately reduced.

Class 3: one or more primary fulcrum line(s) may exist; from which one or more may become a factual rotation axis when a denture is inserted. Rotating around this/these rotation axis the denture will tip upward on one side and downward on the other side (i.e. two-directional rotary movements occur). In this class of arches RPD is recommended, the baseplate of RPD can not be reduced and, *resilient* type telescopes (i.e. delayed type dental support) should be used.

Introduction

A survey and an account of the diverse factors influencing treatment planning are only possible if they are systemized. This is the primary aim of the prosthetic classification of partially edentulous arches (Fábíán and Fejérdy 1979) described in this chapter. This classification is referred to as "prosthetic classification"; because its major goal is to enhance treatment planning and denture designing procedures for those partially edentulous patients being suitable for complete arch reconstruction ("full-arch" reconstruction) but not suitable for implant dentistry (Fábíán and Fejérdy 1979, 1981). (Please consider that a patient may be not suitable for implant dentistry because of dental, medical but also because of social, psychological or numerous other reasons). However, all systemization may lead to simplification which is a huge mistake of treatment planning in dentistry, and lack of attention to all facets of treatment planning may invite treatment failure (Laney 1983, Bensing 2000, Green and Laskin 2000, Fábíán and Fejérdy 2010). Therefore, besides giving an introduction to prosthodontic classification (see second and third part of this chapter) several other aspects recommended for consideration during treatment planning will be mentioned first as an introduction as follows.

It should be emphasized that patient's subjective need as well as the normatively determined need for replacement of missing teeth should be evaluated carefully before any prosthodontic treatment aiming at complete arch ("full-arch") reconstruction (Elias and Sheiham 1998, Heft et al. 2003, Mazurat and Mazurat 2003, Graham et al. 2006, Fábíán and Fejérdy 2010). Accordingly, the success of a prosthetic treatment strongly depends on the patients' perception of the value of such replacement (Mazurat and Mazurat 2003). It should be considered that, patients are rather focused on psychosocial meaning of the mouth than to physiological function of the teeth when defining subjective need for replacement of missing teeth (Graham et al. 2006). Although occlusion of a complete dental arch is preferable normatively in most cases, many people are really satisfied with less than 28 natural ($\pm$ wisdom) teeth (Rosenoer and Sheiham 1995, Elias and Sheiham 1998, Jones et al. 2003, Ekanayake and Perera 2005, Cunha-Cruz et al. 2007). In fact, there is a discrepancy between the normative need and perceived need. Therefore, prompt replacement of absent teeth without a subjective need of patients may lead to an overtreatment and discomfort (Witter et al. 1994, 1999, Mazurat and Mazurat 2003, Fejérdy 2007) especially when removable partial

dentures (RPDs) are used because of financial (or other) reasons (Witter et al. 1994, 1999, Mazurat and Mazurat 2003, Bae et al. 2006).

It should be also considered that aesthetics seems to be more important than other functions for a great majority of individuals (Elias and Sheiham 1998, Shor et al. 2005, Zlatarić and Celebić 2008); therefore there can be a lack of subjective need for replacement of missing posterior (particularly molar) teeth (Witter et al. 1994, Elias and Sheiham 1998, Mazurat and Mazurat 2003) especially if only a few teeth are absent (Zitzmann et al. 2007). Although aesthetics seems to be more important than other functions for a great majority of individuals (Elias and Sheiham 1998, Shor et al. 2005, Zlatarić and Celebić 2008), expectations and wishes of patient related to chewing (Mazurat and Mazurat 2003, Szentpétery et al. 2005, Zlatarić and Celebić 2008, Roumanas 2009) must also not be underestimated (Mazurat and Mazurat 2003, Fábián et al. 2007, Zlatarić and Celebić 2008, Fábián and Fejérdy 2010). Although chewing ability can be acceptable for patients functioning from second premolar to second premolar (i.e. when all molars missing) (Mazurat and Mazurat 2003); more advanced loss of teeth may induce strong difficulties of chewing. In such cases improvement of mastication may be a highly important wish of patients certainly (Mazurat and Mazurat 2003).

Patients' expectations related to speech may also influence satisfaction with dentures strongly (Zlatarić and Celebić 2008, Roumanas 2009, Fábián and Fejérdy 2010). Similarly, numerous other expectations and wishes of patients related to the denture and/or dental treatment may also appear (Levin and Landesman 1976, Fábián and Fábián 2000, Fábián et al. 2007, Roumanas 2009). Importantly, executable and acceptable wishes and expectations of patient should be fulfilled (Fábián and Fábián 2000, Mazurat and Mazurat 2003, Fábián et al. 2007). If inexecutable and/or unacceptable wishes and expectations also appear, the clinician should carefully weigh the option of nontreatment (Levin and Landesman 1976, Stein 1983, Fábián and Fábián 2000, Fábián et al. 2007).

Besides premised subjective need and conscious wishes (Fábián and Fábián 2000, Smith and McCord 2004, Fábián et al. 2007) related to denture function, unconscious wishes of patients related to both dental treatment and denture (as well as to dentist) should also be considered (Fábián and Fábián 2000, Fábián et al. 2007, Fejérdy and Orosz 2007). In contrast to conscious wishes, understanding unconscious wishes (e.g. to look younger, to stop the appearance of aging, to be loved by the dentist etc.) may be more challenging but highly important for preventing psychogenic manifestation (Fábián and Fábián 2000, Fábián et al. 2007, Fejérdy and Orosz 2007, Fábián and Fejérdy 2010). It may occur that, the patient's conscious and/or unconscious wishes related to the dental treatment and/or dentures and/or dentist may not harmonize with the reality and possibilities of a prosthetic treatment. It is a matter of considerable significance that the clinician carefully weighs the option of nontreatment in such cases (Stein 1983, Fábián and Fábián 2000, Fábián et al. 2007, Fejérdy and Orosz 2007).

It should be also considered that, fixed dentures are usually preferable to a removable one for most patient (Szentpétery et al. 2005). Therefore properly planned fixed dentures supported either by natural abutment teeth or by properly inserted (Bartling et al. 1999) dental implants may be preferable for most patients. However, financial means (Ringland et al. 2004, Roumanas 2009) and compliance of the patient as well as the amount of stress and possible complications induced by preparation of teeth (Cronström et al. 1998, Goodacre et al. 2003/a) and/or implant-surgical interventions (Kaptein et al. 1998, Bartling et al. 1999,

Walton 2000, Goodacre et al. 2003/b) should also be considered very carefully. Because of the high success rate of endosseous dental implants (Nedir et al. 2004, Comfort et al. 2005, Goené et al. 2005, Schwartz-Arad et al. 2007), strategic extraction of compromised tooth/teeth and their replacement with implant supported fixed dentures should also be considered (Barone et al. 2006, Schwartz-Arad et al. 2007, Kao 2008), especially because compromised teeth may significantly worsen the perceived oral health (Cunha-Cruz et al. 2007).

Fixed dentures are especially preferable in instances in which a space created by the loss of a single tooth or perhaps two adjacent teeth is bounded by clinically adequate abutments (Laney 1983, Fábíán and Fejérdy 2010). Besides bridges, use of implant supported single crown(s) may also be a proper solution in such cases (Nedir et al. 2004, Schwartz-Arad et al. 2007, Kao 2008). If the edentulous areas are longer, multiple abutments are usually necessary to support fixed restorations (Fábíán and Fejérdy 1979, 1981, Fábíán et al. 1983, Laney 1983, Fejérdy et al. 2007); however, the increased length of the span may invite symptoms in sensitive patients. Implant supported fixed prostheses may also be a proper solution also in such cases (Goené et al. 2005, Schwartz-Arad et al. 2007, Kao 2008).

It should be also considered that, the farther the pontics deviate from the straight line through the abutments, the longer is the operative lever arm that may also invite problems due to tipping forces (Fábíán and Fejérdy 1979, 1981, Fábíán et al. 1983, Laney 1983). In cases of increased estimated tipping forces, either insertion of implants or orthodontic movement of the abutment tooth/teeth or use of an RPD may be preferable. In the absence of sufficient number (and/or quality) of abutment teeth (and/or implants) RPDs should be used (Fábíán and Fejérdy 1979, 1981). Combinations of RPDs and fixed dentures may also be used advantageously in such cases (Fábíán and Fejérdy 1979, 1981). In cases with only a few remaining teeth (and/or implants), RPDs may be used. In such cases, especially rotary movements (especially two-directional rotary movements around intermediary abutment tooth interrupting long edentulous areas) may be difficult to control (Fábíán and Fejérdy 1979, 1981, Fábíán 1980, Laney 1983).

Besides above, there are some other aspects of prosthodontic treatment that should also be considered for proper treatment planning. Possibilities to correct patients' existing malocclusion (if any) should be considered before prosthodontic treatment. Occlusal early contacts (Biondi and Picardi 1993, Learreta et al. 2007), alterations of horizontal maxillomandibular relationship (Biondi and Picardi 1993) and improper vertical dimension (Piquero et al. 1999) should be corrected with the new denture(s) (Fábíán and Fejérdy 2010). Possibilities to decrease the risk of an inflammatory response in the pulp following tooth preparation should also be considered (Fábíán and Fejérdy 2010); and time need of careful preparation technique, making well fitting provisional crowns/bridges, applying chemical defense of the stump, and proper methodology for the cementation of both provisorical and permanent restorations (Kern et al. 1996, Brännström 1996) should also be taken into account. Time need of proper shaping of occlusal contacts for both provisional and permanent dentures (avoiding the "prophylactic" removal of occlusal contacts as a "pain preventive measure" (Creech et al. 1984)) as well as time need of accurate precise shaping of marginal closure and gingival site for both provisorical and permanent dentures should also be considered for planning (Hermann et al. 2009, Fábíán and Fejérdy 2010).

Importantly, planning of a time-schedule ensuring enough time to carry out a high quality, precise, "lege artis" dental treatment is one of the most important points to achieve a

successful prosthodontic treatment (Hermann et al. 2009, Fábián and Fejérdy 2010). Timing of the start of prosthetic treatment and the insertion of dentures may also be of particular importance (Müller-Fahlbusch and Sone 1982, Fábián and Fejérdy 2010), because heavy psychosocial and/or psychoemotional stress condition as well as the active phases of psychiatric disorders significantly increase the risk of the appearance of psychogenic manifestations in relation with the prosthodontic treatment (Müller-Fahlbusch and Sone 1982, Fábián and Fejérdy 2010). Therefore, starting of prosthetic treatment or insertion of dentures under any heavy psychological stress conditions and/or active phases of psychiatric disorders should be avoided (Müller-Fahlbusch and Sone 1982, Fábián and Fejérdy 2010); and the prosthetic treatment or insertion of denture should be delayed (Müller-Fahlbusch and Sone 1982, Fábián and Fejérdy 2010).

Basic Principles of the Prosthetic Classification

As indicated above, this classification is referred to as "prosthetic classification" (Fábián and Fejérdy 1979, 1981); because its major goal is to enhance treatment planning and denture designing procedures for those partially edentulous patients being suitable for complete arch reconstruction ("full-arch" reconstruction) but not suitable for implant dentistry. With regard to these aspects partially edentulous arches can be rendered to five main classes such as 1A, 1B, 2A, 2B and 3, and to one subclass (2A/1) as described below (see also Figure 1).

Class 1A: one or more primary fulcrum lines (straight line drawn between two teeth adjacent to the edentulous space) may exist; but none of the primary fulcrum lines is going to become a factual rotation axis when a denture is inserted. No force of rotation will be generated and the denture may not settle. In case of partially edentulous arches of class 1A fixed dentures are recommended (Fábián and Fejérdy 1979, 1981, Fábián et al. 1981). If still a removable partial denture (RPD) were made, baseplate of RPD can be extremely reduced to a framework. Class 1B: one or more fulcrum line(s) may exist, which may become a factual rotation axis when a denture is inserted. The rotational forces, however, will not be very powerful, so that the settling of the denture can be equalized (i.e. compensated due to multiple abutments). In case of partially edentulous arches of class 1B fixed dentures may be recommended (Fábián and Fejérdy 1979, 1981, Fábián et al. 1983). If still a removable partial denture (RPD) were made, baseplate of RPD can be strongly reduced.

Class 2A: only one primary fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. In case of a partially edentulous arch of class 2A removable partial denture (RPD) is recommended and the baseplate of RPD can be moderately reduced only (Fábián and Fejérdy 1979, 1981, Fábián 1980/b). Class 2A/1: only one fulcrum line exists, which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the denture may settle in one direction. This class is a special case (subclass) of class 2A characterized by very few (maximum one or two) remaining teeth. In case of partially edentulous arches of class 2A/1 removable partial denture (RPD) is recommended and the baseplate of RPD can not be reduced (Fábián and Fejérdy 1979, 1981, 1991, Fábián 1980/a,b). Class 2B: Several primary fulcrum lines may exist, and one of which may become a factual rotation axis when a denture is inserted. Rotating around this rotation axis the

denture may settle in one direction. In case of partially edentulous arches of class 2B removable partial denture (combined with or without fixed dentures) are recommended and the baseplate of RPD can be moderately reduced only (Fábíán and Fejérdy 1979, 1981, 1991, Fábíán 1980/b).

Class 3: one or more primary fulcrum line(s) may exist; from which one or more may become a factual rotation axis when a denture is inserted. Rotating around this/these rotation axis the denture will tip upward on one side and downward on the other side (i.e. two-directional rotary movements occur). In case of partially edentulous arches of class 3 removable partial denture (RPD) is recommended, the baseplate of RPD can not be reduced and, importantly, resilient type telescopes (i.e. delayed type dental support) should be used (Fábíán and Fejérdy 1979, 1981, Fábíán 1980/a,b).




Figure 1. Typical examples for the prosthodontic classification of edentulous dental arches (Fábíán and Fejérdy 1979).

Planning Partial Dentures for Various Types of Edentulous Arches

Treatment Planning for Cases of Class 1A and Class 1B

According to the basic principles of prosthodontic classification introduced above, in classes 1A and 1B the conditions for making a bridge are given. The type of denture to be made should be chosen after weighing the advantages and disadvantages of a bridge as opposed to a removable partial denture. (Please remember that, prosthodontic classification is for patients not suitable for implant dentistry, therefore the advantages/disadvantages of implant based restorations will not be discussed.)

For the restoration of partially edentulous arches of class 1A bridge(s) should be made in the first place, because the full dental support the foremost condition of making a bridge is given. In case of dentures made to restore arches of class 1A, pontics do not deviate from the straight line through the abutments. Consequently there is no primary operative lever arm and there are no significant tipping forces. Therefore involvement of auxiliary abutment teeth for dental support of a bridge is not necessary if the periodontal tissues of abutment teeth are healthy. (Fábián and Fejérdy 1979, 1981, Fábián et al. 1981, Fábián 1983, 2001, Fejérdy 2007, Fejérdy et al. 2007).

If still a removable partial denture (RPD) were made, an RPD with metal framework should be used, the baseplate of RPD can be extremely reduced to framework, and dental support (support on remaining teeth only) should be given primarily to equalize masticatory forces. However, RPDs are expedient only if one of their major advantages (i.e. they can be made without preparing a tooth; they are usually cheaper than fixed dentures etc.) predominates because of medical, social, psychological or other reasons (Fábián and Fejérdy 1979, 1981, Fábián et al. 1981, Fábián 1983, 2001, Fejérdy 2007).

In the case of dentures made to restore edentulous arches of class 1B one or more pontic(s) deviate(s) from the straight line through the abutments. Consequently, primary operative lever arm(s) and significant tipping force(s) appear(s). Therefore, involvement of one or more auxiliary tooth/teeth is needed, to prevent the overload of the abutment teeth. If still a removable partial denture (RPD) were made, an RPD with metal framework should be used, the baseplate of RPD can be reduced significantly but saddle(s) should be used. A dento-muscular support (support primarily on remaining teeth but also on mucosal tissues) should be given to equalize the masticatory forces (Fábián and Fejérdy 1979, 1981, Fábián et al. 1981, Fábián 1983, 2001, Fejérdy 2007, Fejérdy et al. 2008).

Treatment Planning for Cases of Class 2A

In class 2A a large edentulous area is located either on distal end(s) of the arch (in one or both side) or interdentally in the premolar or frontal region, whereas the remaining teeth are located in one group (Fábián and Fejérdy 1979, 1981, Fábián 1980/b, Fábián 1983, 2001, Fejérdy 2007). As a result of such a location of a rather large edentulous area, a relatively long operative lever arm and significant tipping force appears which can not be equalized by the remaining teeth. Therefore, in class 2A an RPD with metal framework should be used, the

baseplate of RPD can be reduced significantly but saddle(s) should be used and a dento-musocal support should be given to equalize masticatory forces (Fábíán and Fejérdy 1979, 1981, Fábíán 1980/b, Fábíán 1983, 2001, Fejérdy 2007, Fejérdy et al. 2008).

The conditions for a dento-mucosal support are relatively advantageous in this class, because there is only one primary fulcrum line (straight line drawn between two teeth adjacent to the edentulous space) which is going to become a factual rotation axis when a denture is inserted; and the artificial teeth of RPD lie only on one side of the primary fulcrum line functioning as the axis of rotation (rotating around this rotation axis the denture may settle or tip in one direction only). Therefore, the masticatory force will cause a small-scale settling of the saddle(s) which will stay stable in this position (Fábíán and Fejérdy 1979, 1981, Fábíán 1980/b, Fábíán 1983, 2001, Fejérdy 2007).

In view of retention class 2A is inadventagous, because tensile forces will tip the saddle(s) of denture (either mesially or distally), therefore RPDs should be completed with an indirect retainer in class 2A. In case of clasp retention, we may put the occlusal support far from the saddle, because it removes the rotation axis of the RPD away from the point of greatest settlement of the saddle(s).

Consequently, an advantageous decrease of the angle of rotation occurs, while the occlusal rest positioned far from the saddle advantageously acts as an effective indirect retainer as well. If precision attachment is used, involvement of auxiliary abutment tooth may be needed for proper support and anchorage of the attachment (Fábíán and Fejérdy 1979, 1981, Fábíán 1980/b, Fábíán 1983, 2001, Fejérdy 2007).

Treatment Planning for Cases of Class 2A/1

This class is a special case (subclass) of class 2A characterized by very few (maximum one or two) remaining teeth. In case of partially edentulous arches of class 2A/1 removable partial denture (RPD) is recommended (Fábíán and Fejérdy 1979, 1981, 1991, Fábíán 1980/a,b). Because of the very few remaining teeth muco-dental support (support primarily on mucosal tissues but also on remaining teeth) should be used to equalize masticatory forces. Therefore the baseplate of RPD can not be reduced, but even may be extended to an optimal dimension using functional impression to improve anchorage (Fábíán and Fejérdy 1979, 1981, 1991, Fábíán 1980/a,b).

There are three basic forms may be considered for the prosthetic restoration of cases of class 2A/1 as follows: (1).: An RPD retained by telescope may be advocated in the first place. In case of two remaining teeth (located near by each other) the primary telescopes should preferably be splinted together. Since the baseplate of RPD can not be reduced, a baseplate of acrylic resin without any metal framework will met the demands in this case usually. (2).: RPD may be retained by clasps also in cases of class 2A/1. In such cases however RPD with metal framework should be used, even if the baseplate is not reduced. Abutment crowns may also be used according to indication, and the remaining teeth may be splinted in this case. (3).: If the remaining teeth are two lower cuspids the denture can be advantageously retained using a bar (preferably of precision attachment) (Fábíán and Fejérdy 1979, 1981, 1991, Fábíán 1980/a,b, Fábíán 1983, 2001, Fejérdy 2007).

Treatment Planning for Cases of Class 2B

Cases of class 2B differ from those of class 2A in so far as the remaining teeth are not situated in one group, but there are edentulous areas between them (Fábián and Fejérdy 1979, 1981, Fábián 1980/b, Fábián 1983, 2001, Fejérdy 2007). The basic similarity between cases of class 2B and those of class 2A is seen in the considerable, not compensable extension of the edentulous ridge on one side of the primary fulcrum line functioning as the axis of rotation (rotating around this rotation axis the denture may settle in one direction only). (Fábián and Fejérdy 1979, 1981, Fábián 1980/b, Fábián 1983, 2001, Fejérdy 2007). Because of the relatively long operative lever arm and significant tipping force (which can not be equalized by the remaining teeth) a dento-musoccal support should be given to equalize masticatory forces also in class 2B. Therefore, an RPD with metal framework should be used also in class 2B. Similarly to cases of class 2A, the baseplate of RPD can be reduced significantly but saddle(s) should be used. Because of the edentulous areas between the remaining teeth, the possibilities for retention are in this class always quite favorable, therefore no indirect retainers are necessary (Fábián and Fejérdy 1979, 1981, Fábián 1980/b, Fábián 1983, 2001, Fejérdy 2007, Fejérdy et al. 2008).

There are four basic forms may be considered for the prosthetic restoration of cases of class 2B as follows: (1).: Edentulous areas between remaining teeth will be restored by a bridge with a subsequently fabricated RPD. In this case we in fact transform the case of class 2B into a case of class 2A. (2).: A bar (preferably of precision attachment) will be applied along the edentulous areas between the remaining teeth, which serves as both dental support and anchorage for an RPD. (3).: In case of multiple edentulous areas between the remaining teeth, these areas may be restored using a combination of both bridge(s) and bar(s) as well. Patients usually find it more convenient to have their missing front teeth restored by a bridge while bars (preferably of precision attachment) can be used for proper retention and anchorage in the lateral edentulous areas between remaining teeth (4).: Cases of class 2B may be restored with RPD without the use of any fixed prosthesis (but abutment crowns at best according to indication) (Fábián and Fejérdy 1979, 1981, Fábián 1980/b, Fábián 1983, 2001, Fejérdy 2007).

Treatment Planning for Cases of Class 3

In the case of class 3 few remaining teeth interrupt long edentulous areas, therefore two-directional rotary movements around these remaining teeth (interrupting long edentulous areas) occur which is difficult to control. Rotating around factual rotation axis the denture will tip upward on one side and downward on the other side. Therefore, mucosal support and a delayed type dental support achieved by resilient type telescopes should be used to control two-directional rotary movements. This will ensure primary settlement of the RPD thereby creating evenly distributed load on the underlying teeth and mucosa (Fábián and Fejérdy 1979, 1981, Fábián 1980/a,b, Fábián 1983, 2001, Fejérdy 2007). Accordingly, in case of class 3, removable partial denture is recommended, the baseplate of RPD can not be reduced and resilient type telescopes (i.e. delayed type dental support) should be used to achieve (delayed) dental support and to improve anchorage. The anchorage may also be improved via extension of the baseplate to an optimal dimension using functional impression. A baseplate of acrylic

resin without any metal framework will met the demands in cases of class 3 usually (Fábíán and Fejérdy 1979, 1981, Fábíán 1980/a,b, Fábíán 1983, 2001, Fejérdy 2007).

Conclusion

The prosthetic classification of partially edentulous dental arches (Fábíán and Fejérdy 1979, 1981) may enhance treatment planning and denture designing procedures for those partially edentulous patients being suitable for complete arch reconstruction (“full-arch” reconstruction) but not suitable for implant dentistry. However, simplification of treatment planning because of systemization should be avoided and all facets of a case should be considered to prevent treatment failure.

References

- Bae, KH; Kim, C; Paik, DI; Kim, JB. A comparison of oral health related quality of life between complete and partial removable denture-wearing older adults in Korea. *J. Oral Rehabil.* 2006 33, 317-322.
- Barone, A; Rispoli, L; Vozza, I; Quaranta, A; Covani, U. Immediate restoration of single implants placed immediately after tooth extraction. *J. Periodontol*, 2006 77, 1914-1920.
- Bartling, R; Freeman, K; Kraut, RA. The incidence of altered sensation of the mental nerve after mandibular implant placement. *J. Oral Maxillofac Surg.*, 1999 57, 1408-1410.
- Bensing, J. Bridging the gap. The separate worlds of evidence-based medicine and patient-centered medicine. *Patient Education Counseling*, 2000 39, 17-25.
- Biondi, M. and Picardi, A. Temporomandibular joint pain-dysfunction syndrome and bruxism: etiopathogenesis and treatment from a psychosomatic integrative viewpoint. *Psychother Psychosom*, 1993 59, 84-98.
- Brännström, M. Reducing the risk of sensitivity and pulpal complications after the placement of crowns and fixed partial dentures. *Quintessence Int*, 1996 27, 673-678.
- Comfort, MB; Chu, FC; Chai, J; Wat, PY; Chow, TW. A 5-year prospective study on small diameter screw-shaped oral implants. *J. Oral Rehabil.*, 2005 32, 341-345.
- Creech 3rd, JL; Walton, RE; Kaltenbach, R. Effect of occlusal relief on endodontic pain. *JADA*, 1984 109, 64-67.
- Cronström, R; Owall, B; René, N. Treatment injuries in dentistry - cases from one year in the Swedish Patient Insurance Scheme. *Int. Dent J.*, 1998 48, 187-195.
- Cunha-Cruz, J; Hujoel, PP; Kressin, NR. Oral health-related quality of life of periodontal patients. *J. Periodont Res.*, 2007 42, 169-176.
- Ekanayake, L. and Perera, I. Perceived need for dental care among dentate older individuals in Sri Lanka. *Spec Care Dentist*, 2005 25, 199-205.
- Elias, AC. and Sheiham, A. The relationship between satisfaction with mouth and number and position of teeth. *J. Oral Rehabil.*, 1998 25, 649-661.
- Fábíán, T. Wann und in welcher Form bewehrt sich die Teleskopprothese? *Stomatol DDR*, 1980/a 30, 841-843.

- Fábián T. The significance of prevention in prosthetic dentistry and its realization in practice. *Prothet Stomatol*, 1980/b 30, 4-5.
- Fábián T. *Treatment planning for partially edentulous patients*. (A részleges foghiányok pótlásának tervezése). Budapest: Fogtechnikai Vállalat; 1983; 26-80.
- Fábián T. Prosthodontic treatment of partially edentulous patients. (A részleges foghiányok pótlása). In Fábián T editor *Principles of prosthetic dentistry*. (A fogpótlástan alapjai). Budapest: Semmelweis Kiadó; 2001; 231-280.
- Fábián, T. and Fejérdy, P. Prosthetic classification of partially edentulous jaws. *Fogorv Szle*, 1979 72, 310-315.
- Fábián, T. and Fejérdy, P. Prothetische Einteilung der Lückengebisses. *Stomatol DDR*, 1981 31, 759-765.
- Fábián, T., Fejérdy P. Effect of partial denture design on the remaining teeth. In: Vogel G editor. *Protesi occlusione ATM*. Bologna: Monduzzi; 1991; III/3-9.
- Fábián, TK and Fejérdy, P. *Psychogenic denture intolerance. Theoretical background, prevention and treatment possibilities*. New York: Nova Science Publishers; 2010. 124-128.
- Fábián, TK. and Fábián, G. Dental stress. In: Fink G, editor in chief. *Encyclopedia of stress*. (vol. 1.). San Diego: Academic Press; 2000; 567-659.
- Fábián, T., Fejérdy, P., Simon, G., Gerle, J. Prothesenplanung bei der Klasse 1A. *Stomatol DDR*, 1981 31, 675-678.
- Fábián, T., Gerle, J., Simon, G., Fejérdy P. Planung des Zahnarztes bei Zahnlücken der prothetischen Klasse 1B. *Stomatol DDR*, 1983 33, 539-543.
- Fábián, TK; Fábián, G; Fejérdy, P. Dental stress. In: Fink G, editor in chief. *Encyclopedia of stress. Second edition*. (vol. 1.). Oxford: Academic Press; 2007; 733-736.
- Fejérdy, P. Prosthodontic treatment of partially edentulous elderly patients. (A részleges foghiányok ellátása idős korban). In Fejérdy P, Nagy G, Orosz M editors *Gerostomatology. Dentistry for elderly patients (Gerosztomatológia az idős kor fogászata)*. Budapest: Semmelweis Kiadó; 2007; 170-182.
- Fejérdy, P. and Orosz, M. Personality of the dentist and patient-assistant-dentist interrelationships (A fogorvos személyisége, a beteg-asszisztens-fogorvos kapcsolatrendszer) In: Vértes G, Fábián TK editors. *Psychosomatic dentistry (Fogorvosi Pszichoszomatika)* Budapest: Medicina; 2007; 22-31.
- Fejérdy, P; Tollas, ÖL; Fejérdy, L; Borbély, J; Hermann, P. Monitoring trends in fixed partial denture design based on national surveys. *Fogorv. Szle*, 2007 100, 279-288.
- Fejérdy, P; Borbély, J; Schmidt, J; Jáhn, M; Hermann, P. Removable partial denture design and its effect on remaining teeth, based on Hungarian national survey. *Fogorv. Szle*, 2008 101, 3-11.
- Goené, R; Bianchesi, C; Hüerzeler, M; Del Lupo, R; Testori, T; Davarpanah, M; Jalbout, Z. Performance of short implants in partial restorations: 3-year follow-up of Osseotite implants. *Implant. Dent.*, 2005 14, 274-280.
- Goodacre, CJ; Bernal, G; Rungcharassaeng, K; Kan, YKK. Clinical complications in fixed prosthodontics. *J. Prosthet. Dent*, 2003/a 90, 31-41.
- Goodacre, CJ; Bernal, G; Rungcharassaeng, K; Kan, YKK. Clinical complications with implant and implant prostheses. *J. Prosthet. Dent.*, 2003/b 90, 121-132.

- Graham, R; Mihaylov, S; Jepson, N; Allen, PF; Bond, S. Determining "need" for a removable partial denture: a qualitative study of factors that influence dentist provision and patient use. *Brit. Dent. J*, 2006 200, 155-158.
- Green, CS. and Laskin, DM. Temporomandibular disorders: Moving from a dentally based model to a medically based model. *J. Dent. Res.*, 2000 79, 1736-1739.
- Heft, MW; Gilbert, GH; Shelton, BJ; Duncan, RP. Relationship of dental status, sociodemographic status, and oral symptoms to perceived need for dental care. *Community Dent. Oral Epidemiol.*, 2003 31, 351-360.
- Hermann, P; Gera, I; Borbély, J; Fejérdy P; Madléna, M. Periodontal health of an adult population in Hungary: findings of a national survey. *J. Clin. Periodontol.*, 2009 36, 449-457.
- Jones, JA; Orner, MB; Spiro, A(3rd); Kressin, NR. Tooth loss and dentures: patients' perspectives. *Int. Dent J.*, 2003 53 (5 Suppl.), 327-334.
- Kaptein, MLA; Hoogstraten, J; Putter de, C; Lange de, GL; Blijdorp, PA. Dental implants in the atrophic maxilla: measurements of patients' satisfaction and treatment experience. *Clin. Oral. Impl. Res.*, 1998 9, 321-326.
- Kao, RT. Strategic extraction: A paradigm shift that is changing our profession *J. Periodontol.*, 2008 79, 971-977.
- Kern, M; Kleimeier, B; Schaller, HG; Strub, JR. Clinical comparison of postoperative sensitivity for a glass ionomer and a zinc phosphate luting cement. *J. Prosthet Dent.*, 1996 75, 159-162.
- Laney, WR. Considerations in treatment planning. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment in prosthodontics*. Philadelphia: Lea and Febiger; 1983; 157-181.
- Learreta, JA; Beas J; Bono, AE; Durst A. Muscular activity disorders in relation to intentional occlusal interferences. *Cranio*, 2007 25, 193-199.
- Levin, B. and Landesmann, HM. A practical questionnaire for predicting denture success of failure. *J. Prosthet. Dent.*, 1976 35, 124-130.
- Mazurat, NM. and Mazurat, RD. Discuss before fabricating: Communicating the realities of partial denture therapy. Part I: Patient expectations. *J. Can. Dent. Assoc.*, 2003 69, 90-94.
- Müller-Fahlbusch, H. and Sone, K. Präprotetische Psychagogik. *Deutsch Zahnärztl Z*, 1982 37, 703-707.
- Nedir, R; Bischof, M; Briaux, JM; Beyer, S; Szmukler-Moncler, S; Bernard, JP. A 7-year life table analysis from a prospective study on ITI implants with special emphasis on the use of short implants. Results from a private practice. *Clin. Oral Implant Res.*, 2004 15, 150-157.
- Piquero, K; Ando, T; Sakurai, K. Buccal mucosa ridging and tongue indentation: incidence and associated factors. *Bull. Tokyo Dent. Coll*, 1999 40, 71-78.
- Ringland, C; Taylor, L; Bell, J; Lim, K. Demographic and socio-economic factors associated with dental health among older people in NSW. *Aust. N. Z. J. Public Health*, 2004 28, 53-61.
- Rosenoer, LM. and Sheiham, A. Dental impacts on daily life and satisfaction with teeth in relation to dental status in adults. *J. Oral Rehabil.*, 1995 22, 469-480.
- Roumanas, ED. The social solution - denture esthetics, phonetics, and function. *J. Prosthodont*, 2009 18, 112-115.

- Schwartz-Arad, D; Laviv, A; Levin, L. Survival of immediately provisionalized dental implants placed immediately into fresh extraction sockets. *J. Periodontol.*, 2007 78, 219-223.
- Shor, A; Shor, K; Goto, Y. The edentulous patient and body image - achieving greater patient satisfaction. *Pract. Proced. Aesthet Dent.*, 2005 17, 289-295.
- Smith, PW. and McCord, JF. What do patients expect from complete dentures? *J. Dentistry*, 2004 32, 3-7.
- Stein, RS. Mutual protective complex of dental restorations. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment of prosthodontics*. Philadelphia: Lea and Febiger; 1983; 306-326.
- Szentpétery, AG; John, MT; Slaze, GD; Setz, JM. Problems reported by patients before and after prosthodontic treatment. *Int. J. Prosthodont.*, 2005 18, 124-131.
- Walton, JN. Altered sensation associated with implants in the anterior mandible: a prospective study. *J. Prosthet Dent.*, 2000 83, 443-449.
- Witter, DJ; De Haan, AF; Käyser, AF; Van Rossum, GM. A 6-year follow up study of oral function in shortened dental arches. Part II: Craniomandibular dysfunction and oral comfort. *J. Oral Rehabil.*, 1994 21, 353-366.
- Witter, DJ; van Palenstein Helderman, WH; Creugers, NH; Käyser, AF. The shortened dental arch concept and its implications for oral health care. *Community Dent. Oral Epidemiol.*, 1999 27, 249-258.
- Zitzmann, NU; Hagmann, E; Weiger, R. What is the prevalence of various types of prosthetic dental restorations in Europe? *Clin. Oral Implant. Res.*, 2007 18 (Suppl. 3), 20-33.
- Zlatarić, DK. and Celebić, A. Factors related to patients' general satisfaction with removable partial dentures: a stepwise multiple regression analysis. *Int. J. Prosthodont.*, 2008 21, 86-88.

Chapter II

Orthodontics for Patient of Prosthetic Dentistry. Possibilities and Complications of Psychosomatic Origin

***Tibor Károly Fábián^{*1}, Anita Beck^{1,2}, Boglárka Rencz³,
Pál Fejérdy¹, Péter Hermann¹ and Gábor Fábián³***

¹Clinic of Prosthetic Dentistry,

²Department of Oral Biology,

³Clinic of Pediatric Dentistry and Orthodontics,

Semmelweis University Budapest, Faculty of Dentistry, Budapest, Hungary, EU

Abstract

Various dental specialties have made such enormous advances that it has become impossible to absorb all that is new in all dental areas. Therefore a collaborative effort between orthodontist and prosthodontist is often necessary to achieve an optimal treatment result. Prosthodontists and restorative dentists rely on orthodontics as a means of providing more comprehensive care to their patients. Such collaboration is not only desirable, but mandatory if dentists are to provide highest level care for their patients. Preprosthetic orthodontics gains significance in both esthetics and functionality of the denture being under preparation. Therefore, an interdisciplinary approach may also be cost-effective for both patient and dentist from the standpoint of producing more stable and durable restorations of high esthetics. Orthodontics may contribute to the success of cases undergoing restorative dental procedures where preliminary minor tooth movement is necessary, to correct local malpositions and/or to move abutment teeth with simple appliances. Success of implant placement may also be improved via orthodontic forced extrusion of periodontally compromised teeth. Collaboration of orthodontist and prosthodontist may be even more demanding when correction of primary orthodontic anomalies would be needed. No wonder that, the integration of orthodontics and

* E-mail address: fab@fok.usn.hu.

prosthetic dentistry into a multidisciplinary rehabilitation is often indispensable for a proper dental care of the patients. A proper plaque control is also essential for a successful treatment in all cases. Along with mechanical and biological principles, unique psychological and social aspects of the combined orthodontic and prosthetic treatment should also be considered for a successful therapy. In this chapter the important role of orthodontics and its contribution to the success of prosthetic dentistry for adult patients is discussed.

Introduction

Various dental specialties have made such enormous advances that it has become impossible to absorb all that is new in all dental areas [LaSota 1988]. Therefore a collaborative effort between orthodontist and prosthodontist is often necessary to achieve an optimal treatment result [LaSota 1988]. Prosthodontists and restorative dentists rely on orthodontics as a means of providing more comprehensive care to their patients [LaSota 1988]. Such collaboration is not only desirable, but mandatory if dentists are to provide highest level care for their patients [LaSota 1988]. Orthodontics may contribute to the success of cases undergoing restorative dental procedures where preliminary minor tooth movement is necessary, to correct local malpositions and/or to move abutment teeth with simple appliances [Shapira 1978, Keesee et al. 2002]. Success of implant placement may also be improved via orthodontic forced extrusion of periodontally compromised teeth [Atherton 1970, Heithersay 1973, Ingber 1974]. Collaboration of orthodontist and prosthodontist may be even more demanding when correction of primary orthodontic anomalies would be needed [LaSota 1988]. No wonder that, the integration of orthodontics and prosthetic dentistry into a multidisciplinary rehabilitation is often indispensable for a proper dental care of the patients [Diedrich 1996]. A proper plaque control [Diedrich 1996, Fábíán et al 2000] as well as careful evaluation of relevant psychosocial and psychoemotional aspects [Fábíán et al 2000] are also essential for a successful treatment in all cases. In this chapter the important role of orthodontics and its contribution to the success of prosthetic dentistry for adult patients is discussed.

Correction of Drifting or Tilting towards the Adjacent Tooth

Caries or other approximal lesion may result in the drifting or tilting of teeth towards their neighboring teeth, which makes restoration difficult. The close root approximation and poor long-axis alignment, may lead to inability to restore proper contours and to establish esthetics [Keesee et al. 2002]. Protection of the adjacent tooth during preparation procedure and avoidance of the exposure of the pulp of a tipped tooth under preparation may also be difficult. In other cases lack of adequate mesiodistal space may not provide adequate space for fabrication insertion and maintenance of implant crowns, because of a drifting/tilting of the adjacent tooth [Rotella et al. 2011]. To solve all these problems, orthodontic separators have been used to create space for crown placement on misaligned premolars and molars

[Reagan 1988, VanderWeele et al. 1998, Keesee et al. 2002, Rotella et al. 2011]. Placement of an elastic orthodontic separator accomplished with 2 standards of dental floss used to stretch the separator and snap it in between the two teeth may be used as a simple technique to achieve minor tooth movement [VanderWeele et al. 1998, Keesee et al. 2002]. Similarly, orthodontic separator can be placed between the adjacent tooth and a provisional screw-retained temporary crown of an implant [Rotella et al. 2011].

Roughly 1-1.5 mm of separation can be reached in 1-2 weeks with this method [Keesee et al. 2002, Rotella et al. 2011]. If further separation is needed, the tooth contact can be restored with an interim restoration and the process can be repeated [Keesee et al. 2002, Rotella et al. 2011]. However, it should be considered that, bodily movement (translation) cannot be expected because the application of force is a tipping one. Further, the tipping force may be extrusive therefore the occlusion must be monitored during the treatment [Keesee et al. 2002]. It should be also considered that the procedure may lead to transient hypermobility of the involved teeth [Keesee et al. 2002], and that the procedure is just as likely to move both adjacent tooth if the anchorage of both of them is insufficient [Keesee et al. 2002]. Since osseointegrated implants do not move in response to orthodontic loads [Odman et al. 1994, Kato and Kato 2006] premised problems do not occur when the separator was placed between an adjacent tooth and an implant (i.e. implant with provisional crown certainly) [Rotella et al. 2011]

Correction of Drifting or Tilting into the Pontic Space

To set a tooth (tipped into the pontic space) upright and/or to decrease the span (diastema) between an abutment tooth (drifted into the pontic space) and the adjacent tooth may also be needed frequently [LaSota 1988, Claman et al. 2003, Lewinstein et al. 2003]. In such cases the position of the abutment tooth may be optimized using brackets, arch wire, and elastic chain [Claman et al. 2003] or activated coil-spring [Lewinstein et al. 2003]. Removable orthodontic appliance may also be used to close a diastema between teeth [Schmitz et al. 2001]. In certain cases orthodontic rubber band fixed with two composite buildups on the labial surface of two adjacent teeth may also be used to close a diastema between them [Schmitz et al. 2001]. In certain cases adjacent osseointegrated implants serving as abutments for permanent prosthetic constructions (i.e. after the completion of the orthodontic treatment) may also be used to set a tooth upright and/or to achieve bodily movement to decrease the span between the tooth and the implant [Odman et al. 1994]. If brackets are used, they should be placed onto all the abutment teeth of the planned bridge and also on the adjacent teeth (at least on the adjacent tooth of the drifted/tipped tooth). If activated coil-spring is used, it should be positioned in that way that the force hits the drifted/tipped abutment tooth from the pontic side certainly. The orthodontic movement may take few months.

Forced Eruption of Periodontally Compromised Teeth

The use of forced eruption followed by extraction of periodontally compromised teeth provide the opportunity to permanently increase the vertical height of hard and soft tissues at the site of the tooth undergoing this procedure (i.e. forced eruption followed by extraction) [Atherton 1970, Heithersay 1973, Ingber 1974,1989, Mantzikos and Shamus 1997, Capri et al. 2003]. However the bone that moves coronally is attached to the apical third of the root (because of the advanced periodontal disease of the tooth at issue), where the root diameter is relatively narrow [Capri et al. 2003]. Therefore, the coronal movement of the bone may produce the necessary height to the edentulous ridge for making esthetic pontic or insertion of an implant [Capri et al. 2003, Chambrone and Chambrone 2005, Holst et al. 2007, Kim et al. 2011]; but the width may remain deficient in many cases because of the narrow root diameter [Capri et al. 2003]. To increase the buccolingual bulk of the alveolar bone, a buccal root torque component may be applied concomitantly [Korayem et al. 2008], but a mucogingival surgery to widen the ridge may also be needed in certain cases to develop a desired ridge width [Capri et al. 2003].

For forced eruption of periodontally compromised teeth brackets should be bounded to the treated tooth and two-three adjacent neighboring teeth at both sides [Capri et al. 2003]. The orthodontic movement can be achieved by a 0.016 inch nickel titanium wire [Capri et al. 2003, Chambrone and Chambrone 2005] with the insertion of the bracket in a more apical position onto the erupted tooth [Capri et al. 2003, Erkut et al. 2007]. To avoid intrusion of the anchorage teeth, use of a 0.019 x 0.025 inch stainless steel overlay wire (auxiliary anchorage wire) is also recommended [Chambrone and Chambrone 2005, Korayem et al. 2008]. For tooth eruption, light constant extrusive forces are recommended (i.e. 15 g for anterior and 50 g for posterior teeth) [Korayem et al. 2008]. The extrusion rate is roughly 2.0 mm per month. A retention/stabilization phase of the same length as of the active extrusion phase is recommended prior to extraction [Korayem et al. 2008]. The desired movement can be usually achieved in roughly three months with another three month retention phase before extraction [Capri et al. 2003, Chambrone and Chambrone 2005].

Forced Eruption of Teeth with Advanced Hard Tissue Damage

Forced eruption generates alteration of the periodontal tissues resulted in a coronal shift of periodontal fibers, marginal gingiva and bone (coupled with a somewhat more pronounced coronal shift of the erupted tooth certainly). Forced eruption may be used for clinical crown lengthening of buried roots, teeth with non-restorable advanced coronal damage and teeth with subgingivally located hard tissue damage (i.e. deep approximal carious lesion) [Ingber 1976, Heithersay 1973, Shiloah 1981, Stein and Sidley 1997, Ziskind et al. 2000]. In such cases the coronal movement of the bone produces both the necessary height and width to the alveolar crest for making esthetic restoration (i.e. crown or post and core with crown).

Forced eruption of a buried root can be carried out with a simplified eruption procedure using a horizontal bar fixed between the neighboring teeth, a hook cemented into the root canal, and an elastic band responsible for the tooth movement [Stein and Sidley 1997, Ziskind et al. 2000]. Brackets bounded to two-three adjacent neighboring teeth at both side together with a 0.016 inch nickel titanium wire [Ziskind et al. 2000, Chambrone and Chambrone 2005] fixed to the hook cemented into the root canal may also be used certainly. Forced eruption of teeth with non-restorable coronal damage may be carried out either in the same way as described for buried roots (i.e. after initial preparation of the tooth for post and core) or after a temporary restoration of the crown with proper buildup for fixation of a bracket [Stein and Sidley 1997, Chambrone and Chambrone 2005]. In the case of teeth with subgingivally located hard tissue damage but with more or less intact coronal region (i.e. a tooth with deep approximal carious lesion), a bracket should be fixed onto an intact coronal surface area of the teeth at issue.

Since in the above cases an extrusion of few mm is usually enough, the orthodontic procedure may take roughly a month only. Care should be taken that placement of the horizontal bar (or orthodontic wire) the hook and the elastic band influence the direction of eruption [Ziskind et al. 2000]. Therefore a root may be unintentionally moved in buccolingual or mesiodistal direction as an accompanying side effect of the forced eruption procedure [Ziskind et al. 2000].

Intrusion of Supraerupted Teeth

Supra-erupted teeth may cause problems such as functional disturbances and occlusal interference and may also cause great difficulty during prosthetic reconstruction. However, prosthetic crown reduction often leads to root canal therapy. To prevent this complications an orthodontic intrusion of supraerupted teeth may be need [Rabie et al. 1998]. However it should be emphasized that, intrusion of teeth without a proper plaque control may lead to pocket formation [Diedrich 1997], therefore great care should be taken about patients' oral hygiene activities and proper professional plaque control should also be carried out regularly [Diedrich 1997]. In this way an increase of bone coverage of the root surfaces can be reached [Heravi et al. 2011], however it is not yet clear that whether an increase of periodontal attachment (i.e. increase of the surface of attachment with periodontal fibers) also occurs or not [Diedrich 1997]. It is very likely that a significant increase of periodontal attachment (i.e. significant increase of attachment with periodontal fibers) can only be achieved if the intrusion procedure is coupled with guided tissue regeneration.

Major problem from an orthodontic point of view of intrusion of elongated teeth is that the procedure may accompanied by the extrusion of the teeth used for anchorage [Kravitz et al. 2007, Heravi et al. 2011]. Orthodontic intrusion of molars requires anchorage reinforcement by incorporating multiple teeth and/or using extraoral devices [Kravitz et al. 2007, Heravi et al. 2011]. However, despite these efforts, efficient intrusion of molars is still difficult to accomplish, and the procedure needs high level compliance of the patient too [Kravitz et al. 2007, Heravi et al. 2011]. Since osseointegrated implants do not move in response to orthodontic loads [Odman et al. 1994, Kato and Kato 2006], orthodontic temporary anchorage devices (i.e. osseointegrated miniscrews) and/or implants inserted for

prosthodontic restorations provide a proper treatment alternative [Greekmore 1983, Costa et al. 1998, Kato and Kato 2006 Kravitz et al. 2007, Landes et al. 2008, Heravi et al. 2011].

Miniscrews should be inserted into a region with high bone density covered by thin keratinized soft tissue [Kravitz et al. 2007, Landes et al. 2008]. (Placement of screws into a region covered by nonceratinized soft tissue may lead to greater failure rate than insertion into attached soft tissue [Kravitz et al. 2007]). Placement of screws into the tuberal region of maxilla is not recommended because of the fine trabecular bone (i.e. low bone density) of this region [Kravitz et al. 2007]. The size of the screw should be selected according to the bulk of the available bone. Diameter of the screws is ranging from 1 mm to 2,3 mm, whereas length of the screws is ranging from 6 mm to 17 mm in the related literature [Chen et al. 2009]. Use of a pilot drill (of 0.2-0.5 mm less diameter than the miniscrew itself) is recommended, although self drilling methods (without pilot drilling and incision) may also be used [Chen et al. 2009]. The screws can be loaded immediately, and most miniscrews can withstand 100-200 g of horizontal immediate load [Chen et al. 2009]. One or two screws are usually inserted into the interradicular bone adjacent to the tooth for intrusion [Kravitz et al. 2007, Landes et al. 2008]. If inadequate interradicular bone is available, screws can be placed palatally [Kravitz et al. 2007]. In certain cases osseointegrated dental implants inserted as abutments for prosthetic restorations (i.e. for implant crowns or bridges) may also be used for anchorage (before making definitive prosthetic restoration) using provisional crowns with proper buildups for orthodontic anchorage [Kato and Kato 2006].

For tooth movement titanium-molybdenum springs [Heravi et al. 2011] as well as elastic chain or nickel titanium coil [Kato and Kato 2006, Kravitz et al. 2007, Landes et al. 2008] may be used. The duration of such an orthodontic procedure usually ranges from 4 to 8 months using 100-200 g force per tooth [Kravitz et al. 2007, Heravi et al. 2011, Landes et al. 2008], and an amount of intrusion between 1.5 – 4.5 mm can be achieved under such circumstances. In certain cases (i.e. for molars) 300 g force per tooth may also be needed for such efficient movement [Kato and Kato 2006]. Importantly, pressure from intrusive forces concentrates at the root apex, which may lead to compression and necrosis of the periodontal ligament and root resorption [Kravitz et al. 2007, Heravi et al. 2011]. However, the amount of the resorption was found to be roughly 0.3 – 0.4 mm which is clinically not significant [Heravi et al. 2011]. A small amount of crest resorption may also occur during active intrusion. However, it is not comparable to the amount of intrusion, therefore there is improved final root coverage at the end of the treatment [Heravi et al. 2011].

Abutment Positioning

Increase of the pontic space (enlargement of the interabutment span) towards an edentulous area located on distal end of the arch is frequently needed and may be advantageous in many cases for prosthetic treatment [LaSota 1988, Lewinstein et al. 2003, Arslan et al. 2006]. However, orthodontic intervention is often overlooked as a viable modality to solve such problems [Cohen 1995]. For such purposes brackets should be placed onto all the abutment teeth of the planned bridge as well as on some adjacent teeth of the mesial abutment tooth/teeth of the planned bridge [Lewinstein et al. 2003]. In some cases brackets are placed on all remaining teeth of the arch [Arslan et al. 2006]. An activated coil-

spring should be positioned in that way that the force hit the drifted/tipped abutment tooth from the pontic side certainly [Lewinstein et al. 2003, Arslan et al. 2006]. Coil-springs may be applied to both the lingual and vestibular surfaces of the distalized tooth to attain a more parallel distalization [Arslan et al. 2006]. Similarly to above, enlargement of the interabutment span (increase of the pontic space) towards an edentulous area located between remaining teeth may also be needed [LaSota 1988, Savadi et al. 2010]. In such cases insertion of brackets onto all remaining teeth of the arch is usually needed, and elastic chain can be used to achieve tooth movement [Savadi et al. 2010]. Similarly, osseointegrated implants serving as abutments for permanent prosthetic constructions (after completion of the orthodontic treatment) may also be used for bodily movement and positioning of abutment tooth utilizing provisional implant crown with proper buildup for orthodontic anchorage [Odman et al. 1994, Drago 1999]. The orthodontic movement for abutment positioning may take 4-5 months [Lewinstein et al. 2003, Arslan et al. 2006], followed by a restabilization of the teeth within roughly 6-10 weeks [Lewinstein et al. 2003, Arslan et al. 2006].

Correction of Primary Orthodontic Anomalies before Prosthetic Treatment

Correction of primary orthodontic anomalies may be even more demanding cases for the collaboration of orthodontist and prosthodontist [LaSota 1988, Fábíán et al. 2000]. Anterior crowding [LaSota 1988], open bite [Fábíán et al. 2000, Furuse et al. 2008] and crossbite [Sakar et al. 2004] as well as rehabilitation of patients with congenital abnormalities including hypodontia [Chaushu et al. 2001, Carter et al. 2003, Thilander 2008], ectopic eruption [Savage and Kokich 2002, Jahangiri et al. 2006], ectodermal dysplasia [Suri et al. 2004] and amelogenesis imperfecta [Siadat et al. 2007] may be mentioned at first. Cases treated with orthognathic surgical procedures including cleft lip and palate [Strong 2002, Moore and McCord 2004] as well as mandibular or maxillary prognathism [Sakar et al. 2004] should also not be forgotten. Recently, many patients with premised primary orthodontic anomalies are benefited from orthodontic and/or orthognathic surgical treatment and require little or no prosthodontic treatment. However, still many treatments of such patients are planned with a final prosthetic restoration. These cases are usually rather challenging presenting with their own multidisciplinary diagnostic and treatment problems. However, treatment planning (and execution) for these patients need rather wide spreading multidisciplinary considerations, which will not be detailed in this chapter being dedicated to “preprosthetic” orthodontics.

Psychosomatic Complications

A combined orthodontic-prosthetic treatment approach may invite unique and complex psychosomatic problems. Diagnostic and treatment decisions for these patients are based largely on objective morphological considerations with the goal of proper aesthetics and biomechanics of the denture. Although patients' decision-making also centers primarily on esthetics, they may vary greatly in their expectations and perceptions of the esthetic effects

and of the desirability and success of a treatment [Fox et al. 1982]. Moreover, little is known about the factors determining these various expectations and perceptions. Further, other subjective factors including self-image [Giddon 1995], self-esteem [Badran 2010], body-image [Polo 2011], treatment related unconscious expectations [Fábíán and Fábíán 2000, Fábíán et al. 2007] and other psychoemotional/psychosocial factors [Fábíán and Fejérdy 2010] also strongly influence patients' perceptions, decisions and satisfaction. Because of this complex mixture of treatment related psychological, psychoemotional and psychosocial factors a risk of treatment failure may be high from psychosomatic point of view [Fábíán et al. 2000].

Beside above, combined orthodontic-prosthetic treatments are usually time consuming and may be frustrating for the patients, which may also lead to psychogenic manifestations [Fábíán and Fábíán 2000, Fábíán et al. 2007]. Orthodontic pain should be considered first as the most frequent and discouraging complaint during treatment, and primary reason for wanting to discontinue care [Oliver and Knapman 1985, Patel et al. 2011]. Decrease of the oral health-related quality of life [Liu et al. 2011] may also be a cause of treatment related frustration especially in the early phase after the insertion of (fixed)orthodontic appliance [Liu et al. 2011]. Complaints related to esthetics and occlusion following orthodontic-prosthetic treatment may also occur, which may also lead to the appearance of psychosomatic symptoms, especially if the patient is frustrated due to the previous treatment procedure [Fábíán et al. 2005].

It should be also considered that, patient satisfaction with a suddenly improved esthetics may lead to a disadvantageous transitory hypomanic reaction in rare cases [Fábíán et al. 2000], which may invite disturbances in partner- and family- or workplace relations of the patient as a consequence [Fábíán et al. 2000]. The reason behind premised rarely occurring hypomanic reaction may be the sudden improvement of self confidence and social/psychological impact due to improved orofacial esthetics [Fábíán et al. 2000, Gazit-Rappaport et al. 2010]. Such reaction may occur after treatment of cases with primary orthodontic anomalies coupled with strongly compromised orofacial esthetics (before the treatment), or after the treatment of patients with strongly compromised esthetics due to edentulousness of the frontal region of the (upper) arch.

Because of above psychosomatic considerations it should be emphasized that, great psychological care should be taken about patients treated with combined orthodontic and prosthetic methods [Fábíán et al. 2000, 2005]. During the treatment, frustration of the patient may be decreased whereas mental health and compliance of the patient may be maintained using several psychotherapeutic approaches [Fábíán and Fejérdy 2010] advantageously combined with several mind-body type psychological techniques such as relaxation, hypnosis, self-hypnosis, photo-acoustic stimulation, biofeedback etc. [Fábíán et al. 2009/a,b, Fábíán and Fejérdy 2010]. Treatment induced pain may be reduced with pain killers like ibuprofen [Patel et al. 2011], with certain physiotherapies like pulsed electromagnetic field (PEMF) therapy [Fábíán et al. 2006, Fábíán and Söti 2007, Fábíán and Fejérdy 2010] and with premised mind-body techniques [Fábíán et al. 2009/a,b, Fábíán and Fejérdy 2010] or with their combination. Great care should also be taken about the after-care of these patients to prevent late appearance of manifestations [Fábíán et al. 2000]. In case of the appearance of psychogenic manifestation(s) an initial psychosomatic dental therapy should be introduced immediately [Fábíán and Fejérdy 2010].

Conclusion

Various dental specialties have made such enormous advances that it has become impossible to absorb all that is new in all dental areas [LaSota 1988]. Therefore a collaborative effort between orthodontist and prosthodontist is often necessary to achieve an optimal treatment result. [LaSota 1988]. Such collaboration is not only desirable, but mandatory if dentists are to provide highest level care for their patients [LaSota 1988]. Preprosthetic orthodontics gains significance in both esthetics and functionality of the denture being under preparation [Diedrich 1996]. Therefore, an interdisciplinary approach can be cost-effective for both patient and dentist from the standpoint of producing more stable and durable restorations of high esthetics [Cohen 1995]. A proper plaque control is also essential for a successful treatment in all cases [Diedrich 1997]. Along with mechanical and biological principles, unique psychological and social aspects of the combined orthodontic and prosthetic treatment should also be considered for a successful treatment [January 1951, Fábíán et al. 2000, 2005].

References

- Arslan, SG; Tacir, IH; Kama, JD. Orthodontic and prosthodontic rehabilitation of unilateral free-end edentulous space. *Australian Dent. J.*, 2006 51, 338-341.
- Atherton, JD. The gingival response to orthodontic tooth movement. *Am. J. Orthod.*, 1970 58, 179-186.
- Badran, SA. The effect of malocclusion and self-perceived aesthetics on the self-esteem of a sample of Jordanian adolescents. *Eur. J. Orthodontics*, 2010 32, 638-644.
- Capri, D; Albehbehani, Y; Smukler, H. Augmentation of an anterior edentulous ridge for fixed prosthodontics with combined use of orthodontics and surgery: A clinical report. *J. Prosthet Dent*, 2003 90, 111-115.
- Carter, NE; Gillgrass, TJ; Hobson, RS; Jepson, N; Meechan, JG; Nohl, FS; Nunn, JH. The interdisciplinary management of hypodontia: orthodontics. *Brit. Dent J*, 2003 194, 361-366.
- Chambrone, L; Chambrone, LA. Forced orthodontic eruption of fractured teeth before implant placement: Case report. *J. Can. Dent. Assoc.*, 2005 71, 257-261.
- Chausu, S; Becker, A; Zalkind, M. Prosthetic considerations in the restoration of orthodontically treated maxillary lateral incisors to replace missing central incisors: A clinical report. *J. Prosthet Dent*, 2001 85, 335-341.
- Chen, Y; Kyung, HM; Zhao, WT; Yu, WJ. Critical factors for the success of orthodontic mini-implants: A systemic review. *Am. J. Orthod. Dentofacial. Orthop.*, 2009 135, 284-291.
- Claman, L; Alfaro, MA; Mercado, A. An interdisciplinary approach for improved esthetic result in the anterior maxilla. *J. Prosthet Dent*, 2003 89, 1-5.
- Cohen, BD; The use of orthodontics before fixed prosthodontics in restorative dentistry. *Compendium*, 1995 16, 110-120.
- Costa, A; Raffini, M; Melsen, M; Miniscrews as orthodontic anchorage. *Int. J. Adult Orthod. Orthognath Surg.*, 1998 13, 201-209.

- Diedrich, P. Preprosthetic orthodontics. *J. Orofac. Orthop.*, 1996 57, 102-116.
- Diedrich, P. Parodontologie und Kieferorthopädie. In: Heidemann D editor. *Parodontologie*. München-Wien-Baltimore: Urban and Schwarzenberg; 1997; 344-362.
- Drago, CJ. Use of osseointegrated implants in adult orthodontic treatment: a clinical report. *J. Prosthet Dent*, 1999 82, 504-509.
- Erkut, S; Arman, A; Gulsahi, A; Uckan, S; Gulsahi, K. Forced eruption and implant treatment in posterior maxilla: A clinical report. *J. Prosthet Dent*, 2007 97, 70-74.
- Fábíán, G; Gáspár, J; Fábíán, TK. Some aspects of the orthodontic-prosthetic approach of adult patients. A case report. *Fogorv Szle*, 2000 93, 233-238.
- Fábíán, G; Bálint, M; Fábíán, TK. Psychology and psychosomatics of the orthodontic treatment. *Fogorv Szle*, 2005 98, 113-118.
- Fábíán, TK. and Fábíán, G. Dental stress. In: Fink G, editor in chef. *Encyclopedia of Stress*. Vol. 1. San Diego: Academic Press; 2000; 657-659.
- Fábíán, TK and Fejérdy P. *Psychogenic denture intolerance. Theoretical background, prevention and treatment possibilities*. New York: Nova Science Publishers; 2010. 9-28, 145-162, 168.
- Fábíán, TK. and Sőti, Cs. Low induction value pulsed electromagnetic field stimulation. [Alacsony indukció értékű mágneses impulzus tér stimuláció] In: Vértés G, Fábíán TK editors. *Psychosomatic dentistry [Fogorvosi Pszichoszomatika]* Budapest: Medicina; 2007; 201-210.
- Fábíán, TK; Sőti, Cs; Fejérdy, P. Low induction value pulsed electromagnetic field stimulation. A minimal-invasive possibility in the treatment of several idiopathic symptoms of denture intolerance. *Fogorv Szle*, 2006 99, 66.
- Fábíán, TK; Fábíán, G; Fejérdy, P. Dental Stress. In: Fink G, editor in chef. *Encyclopedia of Stress. 2-nd enlarged edition*, Vol. 1. Oxford: Academic Press; 2007; 733-736.
- Fábíán, TK; Gótai, L; Krause, WR; Fejérdy, P. Zahnärztliche Hypnoseforschung an der Semmelweis Universität. *Deutsch Z Zahnärztl Hypn*, 2009/a 8, 9-14.
- Fábíán, TK; Kovács, KJ; Gótai, L; Beck, A; Krause, WR; Fejérdy, P. Photo-acoustic stimulation: theoretical background and ten years of clinical experience. *Contemporary Hypnosis*, 2009/b 26, 225-233.
- Fox, RN; Albino, JE; Green, LJ; Farr, SD; Tedesco, LA. Development and validation of a measure of attitudes toward malocclusion. *J. Dent Res.*, 1982 61, 1039-1043.
- Furuse, AY; Franco, EJ; Mondelli, J. Esthetic and functional restoration for an anterior open occlusal relationship with multiple diastemata: A multidisciplinary approach. *J. Prosthet Dent*, 2008 99, 91-94.
- Gazit-Rappaport, T; Haisraeli-Shalish, M; Gazit, E. Psychosocial reward of orthodontic treatment in adult patients. *Eur. J. Orthodontics*, 2010 32, 441-446.
- Giddon, DB. Orthodontic applications of psychological and perceptual studies of facial esthetics. *Semin Orthod*, 1995 1, 82-93.
- Greekmore, TD; The possibility of skeletal anchorage. *J. Clin. Orthod.*, 1983 17, 266-269.
- Heithersay, GS. Combined endodontic-orthodontic treatment of transverse root fractures in the region of the alveolar crest. *Oral Surg Oral Med Oral Pathol*, 1973 36, 404-415.
- Heravi, F; Bayani, S; Madani, AS; Radvar, M; Anbiaee, N. Intrusion of supra-erupted molars using miniscrews: Clinical success and root resorption. *Am. J. Orthod. Dentofacial Orthop*, 2011 139, S170-S175.

- Holst, S; Hegenbarth, EA; Schlegel, KA; Holst, AI. Restoration of a nonrestorable central incisor using forced orthodontic eruption, immediate implant placement, and an all-ceramic restoration: A clinical report. *J. Prosthet. Dent.*, 2007 98, 251-255.
- Ingber, JS; Forced eruption. Part I. A method of treating isolated one and two wall infrabony osseus defects – Rational and case report. *J. Periodontol.*, 1974 45, 199-206.
- Ingber, JS. Forced eruption. Part II. A method of treating non-restorable teeth – Periodontal and restorative considerations. *J. Periodontol.*, 1976 47, 203-216.
- Ingber, JS. Forced eruption: alteration of soft tissue cosmetic deformities. *Int. J. Periodontics Restorative Dent*, 1989 9, 416-425.
- Jahangiri, L; Luu, L; Estafan, D. A multidisciplinary approach for treating congenitally transposed canines: A clinical report. *J. Prosthet Dent*, 2006 95, 413-416.
- January, JW. Psychosomatic in patient management. *Angle Orthod*, 1951 21, 155-162.
- Kato, S; Kato, M. Intrusion of molars with implants as anchorage: A report of two cases. *Clin. Implant Dent. Related Res.*, 2006 8, 100-106.
- Keesee, SM; Baty, DL; Cameron, SM; Lefler, TB; Morris, WJ. A technique for achieving prerestorative minor tooth movement with orthodontic separators. *J. Prosthet Dent*, 2002 88, 544-547.
- Kim, SH; Tramontina, VA; Papalexiou, V; Luczyszyn, SM. Orthodontic extrusion and implant site development using an interocclusal appliance for a severe mucogingival deformity: A clinical report. *J. Prosthet Dent*, 2011 105, 72-77.
- Korayem, M; Flores-Mir, C; Nassar, U; Olfert, K. Implant site development by orthodontic extrusion. A systematic review. *Angle Orthodontist*, 2008 78, 752-760.
- Kravitz, ND; Kusnoto, B; Tsay, TP; Hohlt, WF. The use of temporary anchorage devices for molar intrusion. *J. Am. Dent. Assoc.*, 2007 138, 56-64.
- Landes, CA; Glasl, B; Kopp, S; Sader, R; Ludwig, B. Microanchor mediated upper molar intrusion in deep posterior bite after long-term edentulousness for prosthetic reconstruction with dental implants. *Oral. Maxillofac Surg.*, 2008 12, 155-158.
- LaSota, EP. Orthodontic considerations in prosthetic and restorative dentistry. *Dent Clin North Am*, 1988 32, 447-456.
- Lewinstein, I; Ganor, Y; Pilo, R. Abutment positioning in a cantilever shortened dental arch: A clinical report and static analysis. *J. Prosthet Dent*, 2003 89, 227-231.
- Liu, Z; McGrath, C; Hägg, U. Changes in oral health-related quality of life during fixed orthodontic appliance therapy: An 18-month prospective longitudinal study. *Am. J. Orthod Dentofacial Orthop*, 2011 139, 214-219.
- Mantzikos, T; Shamus, I. Forced eruption and implant site development: soft tissue response. *Am. J. Orthod Dentofacial Orthop*, 1997 112, 596-606.
- Moore, D; McCord, JF. Prosthetic dentistry and the unilateral cleft lip and palate patient. The last 30 years. A review of the prosthodontic literature in respect of treatment options. *Eur. J. Prosthodont. Restor. Dent.*, 2004 12, 70-74.
- Odman, J; Lekholm, U; Jemt, T; Thilander, B. Osseointegrated implants as orthodontic anchorage in the treatment of partially edentulous adult patients. *Eur. J. Orthod.*, 1994 16, 187-201.
- Oliver, RG, Knapman, YM. Attitudes to orthodontic treatment. *Brit. J. Orthod.*, 1985 12, 179-188.
- Patel, S; McGorray, SP; Yezierski, R; Fillingim, R; Logan, H; Wheeler, TT. Effect of analgesics on orthodontic pain. *Am. J. Orthod. Dentofacial. Orthop.*, 2011 139, e53-e58.

- Polo, M. Body dysmorphic disorder: A screening guide for orthodontists. *Am. J. Orthod. Dentofacial Orthop.*, 2011 139, 170-173.
- Rabie, AB; Deng, YM; Jin, LJ. Adjunctive orthodontic treatment of periodontally involved teeth: case report. *Quintessence Int.*, 1998 29, 13-19.
- Reagan, SE. Correcting space loss caused by severe decay: report of case. *J. Am. Dent. Assoc.*, 1988 116, 878-879.
- Rotella, M; Geminiani, A; Ercoli, C. Simplified orthodontic tooth movement with dental implants and orthodontic elastics: A clinical report. *J. Prosthet Dent*, 2011 105, 363-366.
- Sakar, O; Beyli, M; Marsan, G. Combined prosthodontic and orthodontic treatment of a patient with Class III skeletal malocclusion: A clinical report. *J. Prosthet Dent*, 2004 92, 224-228.
- Savadi, RC; Savadi, AR; Kumar, VA; Shetty, VB. Modification of esthetics using a combined orthodontic and a prosthodontic approach: A case report. *J. Indian Prosthodont Soc.*, 2010 10, 128-131.
- Savage, RR; Kokich, VG. Restoration and retention of maxillary anteriors with severe root resorption. *J. Am. Dent. Assoc.*, 2002 133, 67-71.
- Schmitz, JH; Coffano, R; Bruschi, A. Restorative and orthodontic treatment of maxillary peg incisors: A clinical report. *J. Prosthet Dent*, 2001 85, 330-334.
- Shapira, Y. Orthodontics as an aid in adult restorative dentistry. *Refuat Hapeh Vehashinayim*, 1978 27, 23-27, 29-34.
- Shiloah, J. Clinical crown lengthening by vertical root movement. *J. Prosthet Dent*, 1981 45, 602-605.
- Siadat, H; Alikhasi, M; Mirfazaelian, A. Rehabilitation of a patient with amelogenesis imperfecta using all-ceramic crowns: A clinical report. *J. Prosthet Dent*, 2007 98, 85-88.
- Stein, E; Sidley, C. Orthodontic extrusion – a new and simplified aesthetic technique. *J. Dent. Assoc.*, 1997 52, 287-293.
- Strong, SM. Adolescent dentistry: multidisciplinary treatment for the cleft lip/palate patients. *Pract Proced Aesthet Dent*, 2002 14, 333-338.
- Suri, S; Carmichael, RP; Tompson, BD. Simultaneous functional and fixed appliance therapy for growth modification and dental alignment prior to prosthetic habilitation in hypohidrotic ectodermal dysplasia: A clinical report. *J. Prosthet Dent*, 2004 92, 428-433.
- Thilander, B. Orthodontic space closure versus implant placement in subjects with missing teeth. *J. Oral. Rehabil.*, 2008 35 (Suppl. 1), 64-71.
- VanderWeele, RA; Broome, JC; Ramer JP. Regaining space by using elastic orthodontic separators. *Gen Dent*, 1998 46, 454-456.
- Ziskind, D; Sharon, E; Hirschfeld, Z; Becker, A. Analysis of lateral tooth movement during forced orthodontic eruption. *J. Prosthet Dent*, 2000 84, 462-466.

Chapter III

Effects of Wearing Partial Denture Prosthesis on Jaw Functions and Associated Prefrontal Cortex Activity

Noriyuki Narita*

Department of Removable Prosthodontics,
Nihon University School of Dentistry at Matsudo,
Clinical Department of Neurological Dentistry,
Hospital of Nihon University School of Dentistry at Matsudo,
Matsudo, Japan

Abstract

Purpose: Denture wearing is considered to improve chewing ability, chewing comfort, and quality of life in elderly individuals. In an attempt to clarify the effects of prosthodontic treatment on activation of the prefrontal cortex involved in higher cognitive functions, we investigated the efficacy of wearing partial denture from the aspects of jaw functions such as muscle and movement activities and associated prefrontal cortex activity.

Methods: Twelve partially edentulous patients were enlisted as subjects. Functional near-infrared spectroscopy was used to detect changes of hemoglobin concentration in the prefrontal cortex area, while jaw muscle EMG and jaw motion activities were simultaneously examined. Masticatory scores were also determined to evaluate the subjective awareness of chewing ability and occlusal contact features were examined using a Dental Prescale Occluzer System. Statistical comparisons were performed to compare between data obtained while wearing and not wearing a partial denture prosthesis.

* E-mail address: narita.noriyuki@nihon-u.ac.jp.

Results: Significant increases were seen in chewing-related jaw functions such as jaw muscle EMG and jaw motion activities, as well as prefrontal cortex activity associated with wearing a partial denture prosthesis as compared with the not wearing condition. Furthermore, occlusal contact and masticatory scores were also significantly increased by wearing a partial denture prosthesis.

Conclusions: Occlusal reconstruction obtained by wearing a partial denture prosthesis improved prefrontal cortex activity and the awareness of chewing ability, as well as jaw functions, as shown by increased jaw muscle EMG and jaw motion activities. From these findings, we concluded that appropriate denture wearing might have an impact on not only jaw functions, but also cognitive functions closely related to masticatory performance. The relationship between masticatory function and cognitive ability is discussed from the standpoint of prosthodontic treatment in this chapter.

Keywords: Prefrontal cortex; partial denture prosthesis; near-infrared spectroscopy; masticatory score; jaw muscle activity; jaw motion; occlusal contact

Introduction

The relationships between chewing ability, and quality of life and activities of daily life have been reported, and chewing ability has been shown to be closely related to the quality of life of elderly individuals [1]. Findings from a study of an 80-year-old population showed that a greater variety of chewable foods and number of teeth were positively related to indicators of physical fitness, such as hand grip strength and leg extensor strength, which suggested the importance of preventative dental care to preserve chewing ability for enhancing the quality and activities of daily life of very elderly individuals [2]. In addition, edentulous subjects without a denture prosthesis showed significant deterioration in physical ability as compared with that of dentate subjects with 20 or more teeth, while the 6-year mortality rate of edentulous subjects without dentures was significantly higher than that of subjects with 20 or more teeth [3]. From those results, it is assumed that poor dentition status in aged individuals who do not use dentures might lead to deterioration in systemic health condition [4]. Furthermore, few remaining teeth might indicate increased risk of dementia, while edentulous individuals or those with as few as from 1 to 9 teeth may develop dementia later in life. These relationships between masticatory and cognitive functions in aged individuals based on the number of teeth present in cognitively normal subjects were significantly greater than in those who were cognitively impaired.

Maximum bite force, occlusal contact area, and masticatory scores for cognitively normal subjects were also significantly higher than those of cognitively impaired individuals [5]. Considering chewing ability from the standpoint of dental medicine, denture use may improve chewing activity and comfort, denture satisfaction, and dental quality of life in aged individuals, which might also contribute to the prevention of neurological manifestations such as dementia and cognitive impairment. In addition, eating satisfaction and the sense of comfort when chewing with dentures have been suggested to have influences on physiological and psychological wellbeing in elderly individuals in comparison to those who do not wear a denture prosthesis [6].

Based on the above background, we speculated that prosthodontic treatment contributes to higher brain functions and prevents neurological cognitive impairment in aged individuals.

Furthermore, the prefrontal cortex is one of the brain regions strongly susceptible to age-related changes in structure and function, and also has an impact on cognition and behavior [7].

Studies of masticatory function have been conducted using primates to define the detailed cortical neural mechanisms by means of physiological approaches [8-17], and advances over the past decade in brain imaging techniques, such as magnetoencephalography (MEG), functional magnetic resonance imaging (f-MRI), and functional near-infrared spectroscopy (f-NIRS), now allow investigation of neural mechanisms related to human behavior and cognition, such as perception, motor control, memory, and aspects of consciousness.

Recently, we reported the activation of prefrontal cortex by wearing partial denture prostheses with a functional near-infrared spectroscopy system [18]. In this chapter, the effects of wearing a prosthesis on activation of the prefrontal cortex and its functional relationships with reconstructed oral functions are discussed.

Methods

Subjects and Dental Treatments

Twelve partially edentulous patients (6 men, 6 women; average 63.1 years old) were enlisted as subjects, after obtaining informed consent. The Ethical Committee of Nihon University School of Dentistry at Matsudo approved the experimental protocol (EC06-008). During the 3 months prior to the study, the subjects received sufficient prosthodontic treatments in the form of a conventional removable partial denture prosthesis, including artificial teeth, a denture base, clasp retainers, major and minor connectors, and rest under a cross-arch stabilized condition. None of the subjects had complaints about the prosthesis, such as discomfort, pain, or difficulties in masticatory performance. Each had individual patterns of tooth loss, and consequently received examinations in order to determine the effects of wearing a dental prosthesis on higher brain and masticatory functions.

Measurements and Examinations

In this study, prefrontal cortex activity, jaw muscle EMG activity, and jaw movement tracking during masticatory performance, occlusal contact and masticatory scores were examined. Figure 1 presents the feature of simultaneous measuring of prefrontal cortex activity, jaw muscle EMG and jaw motion activities.

Measurement of Activity in Prefrontal Cortex

Functional near-infrared spectroscopy (f-NIRS system, ETG-100 optical topography, Hitachi Medical Co.) was used to measure the activity in the prefrontal cortex during masticatory performance while not wearing and wearing a partial denture prosthesis. This optical topographic system is capable of measuring changes in the concentrations of oxy-hemoglobin ([oxy-Hb]) and deoxy-hemoglobin ([deoxy-Hb]) with a time resolution of 0.1 seconds.




Figure 1. Devices used for simultaneous measuring of prefrontal cortex activity, masticatory score, jaw muscle EMG activities, and jaw motion tracking.




Figure 2. Locations of probe measuring points for near-infrared spectroscopy in dorsal prefrontal cortex area. The positions of the NIRS probes attached to the prefrontal region were standardized. The bottom lines of alignment of the NIRS probes were fit to FP1 and FP2 following EEG electrode placements using the international 10/20 method.

The probes of the optical topography system were set in symmetric order over the prefrontal areas in order to detect the changes in cerebral blood volume and near-infrared spectroscopy with 22 channels (3×5 arrays) was used. The NIRS probes were attached to the prefrontal region and standardized among the subjects, while the bottom lines of the NIRS probes were set according to the FP1 and FP2 electrode placements, with referral to the international 10/20 method used for electroencephalographic examinations (Figure 2).

Measurements of Jaw Muscle EMG, Jaw Motion Activity, and Features of Occlusal Contacts

EMG recording was simultaneously conducted to detect bilateral jaw muscle activities (masseter muscle: Mm, anterior part of the temporal muscle: Ta, anterior digastrics muscle: AD). EMG signals were monitored with a Bioelectric Amplifier (N5198, NEC Co.) and surface electrodes (SEE105, GE Medical Systems Co.). The subjects were seated in an upright position and fitted bilaterally with EMG electrodes, which were attached to the skin surface corresponding to the Mm, Ta, and AD positions, with a reference electrode attached to the ear lobe. In this study, exact EMG muscles activities for Mm and Ta were used for the data analyses. In addition, a jaw motion tracking device (Mandibular Kinesiograph *K6I*, *Myotronics*) was used to evaluate the differences in jaw motion activities between the not wearing and wearing conditions. Occlusal contact was also examined using a Dental Prescale (50H type R, Fuji Photo Film Co.) and Occluzer System (Fuji Photo Film Co.) to evaluate changes in occlusion with and without wearing a partial denture prosthesis.

Examination of Chewing Ability

Chewing ability was examined by the masticatory score based on a questionnaire reported by Hirai [19]. The masticatory score was calculated using 35 food items divided into 5 groups according to chewing difficulty level. In this study, masticatory scores were examined for both the not wearing and wearing conditions.

Stimulation task

A typical task performance consisted of chewing 5 times with each chewing trial conducted for 10 seconds. Each of the 5 trials was separated from the succeeding trial by a 40-second rest phase (Figure 3)




Figure 3. Sequence of task performance. The chewing task consisted of chewing 5 times and each trial lasted for 10 seconds, with a 40-second rest phase between each trial. The NIRS system used (ETG-100 optical topography) utilizes 2 near-infrared wavelengths of 780 and 830 nm.

For the stimulation chewing task, we used 1 piece of chewing gum (Freezone, LOTTE Co.), and the start and end of each chewing trial was indicated to the participant by verbal commands.

The subjects were instructed to chew on either the right or left side during a single session, and avoid intentional head movements while the task performance.

Data Analysis

During each session, data were recorded under 2 different oral conditions; while not wearing and wearing a partial denture prosthesis. Optical data were stored in the f-NIRS system and later transferred to a personal computer (Think Centre, IBM Co.), then quantitatively analyzed using the MATLAB software application (Topo Signal Processing Type G: Hitachi Medical Co.).

By observing changes in wave forms, the average [oxy-Hb] concentrations in all detected channels showed chewing-related changes during task performance under both the not wearing and wearing conditions (Figure 4). In contrast, the average [deoxy-Hb] concentrations in the detected channels did not show clear changes during the task performance under both conditions (Figure 4). Therefore, qualitative and quantitative analyses for the topographical changes in concentrations were conducted using [oxy-Hb] data obtained from all of the subjects (Figures 5, 6).

Jaw muscles EMG and jaw motion activities were quantitatively analyzed using a PC program (Multi-Scope, Medical TrySystem Ltd), as were cycle duration and burst duration. Amplitude activity was also revealed from the peak, area, and average of jaw muscles EMG activities of Mm and Ta, while jaw motion activities shown in the vertical, lateral, and




Figure 4. Hb maps of averaged Hb concentrations. Averaged [oxy-Hb] (red line) and [deoxy-Hb] (blue line) concentrations during the not wearing (a) and wearing (b) conditions are presented. Notably, the changes in wave forms and average concentrations of [oxy-Hb] in all detected channels clearly showed chewing-related differences during task performance between the not wearing and wearing conditions. Two blue arrows shown the onset and end of the chewing trial.

Color

Fig. 4




Figure 5. Hb topographies of averaged Hb concentrations. Topographical charts showing prefrontal cortex activation during the not wearing at 30 seconds (a) and wearing at 29.2s (b) conditions. Dorsal prefrontal cortex was markedly activated by wearing a partial denture prosthesis as compared with the not wearing condition.




Figure 6. Statistical comparison of prefrontal cortex activities between the not wearing and wearing conditions. (a) Statistical Hb map showing the highest significant t value at 26.7 seconds (red arrow). Red and blue horizontal dotted lines show the significance levels at 0.01 (t value: 3.105807) and 0.05 (t value: 2.200985), respectively. Two blue arrows show the onset and end of the chewing trial. (b) Statistical topography map showing a significant increase of [oxy-Hb] in the dorsal prefrontal cortex area while wearing a partial denture prosthesis as compared with the not wearing condition.

Features of occlusal contact was examined using a Dental Prescale (50H type R, Fuji Photo Film Co.) with 2 seconds of maximal biting, then total bite force and total occlusal contact area were qualitatively evaluated under the not wearing and wearing conditions using an Occluzer System (Fuji Photo Film Co.). Masticatory score determined with the questionnaire was calculated using the following formula: (total score of Group I + total score of Group II $\times$ 1.14 + total score of Group III $\times$ 1.3 + total score of Group IV $\times$ 1.52 + total score of Group V $\times$ 3) $\times$ 100/111.4 [19].

Statistical Analysis

A paired t-test was used to examine the NIRS data for [oxy-Hb] concentrations obtained under the not wearing and wearing conditions. Data were transferred to a PC (Precision T3400, Dell Co.), then calculated using MATLAB (Topo Signal Processing Type-G: Hitachi Medical Co.) in order to compare the prefrontal cortex activities between the not wearing and wearing conditions. A paired t-test was also used when the data samples were taken from a population in which the changes were normally distributed in order to compare jaw muscle EMG and jaw motion activities, total occlusal contact area and force, and masticatory scores between the not wearing and wearing conditions. On the other hand, if data samples were taken from a population in which the changes were not normally distributed, a Wilcoxon signed rank test was used for non-parametric statistical analysis. The statistical PC program SigmaStat (Jandel Science Inc.) was used for analyses of jaw muscle EMG and jaw motion activities, occlusal contact, and masticatory score. P values less than 0.05 were considered to be statistically significant.

Results

Prefrontal Activity during Chewing Task Performance

Averaged Hemoglobin Map Presentation

The averaged [oxy-Hb] concentrations in all detected channels showed chewing-related changes during task performance under both conditions of not wearing and wearing a denture prosthesis (Figure 4).

The onset of increase in [oxy-Hb] concentration coincided with the start of the chewing period and continued to increase gradually to the end of the period. Thereafter, the concentration returned to the baseline level within several seconds. In addition, wearing a partial denture clearly induced an increase in [oxy-Hb] concentration as compared to the not wearing condition (Figure 4).

Topographical Presentation of Chewing-Related Prefrontal Cortex Activities

Chewing-related topographical charts produced while not wearing and wearing a denture prosthesis are shown in Figure 5. Comparisons of the charts showed that wearing a partial denture markedly expanded the activation area and increased the intensity of the dorsal part of the prefrontal cortex as compared to the not wearing condition (Figure 5).

Statistical Comparison of Chewing-Related Prefrontal activity between Not Wearing and Wearing Conditions

Prefrontal cortex activity was significantly increased by wearing a partial denture prosthesis as compared with the not wearing condition (Figure 6). According to the results presented as statistical maps and topographies in Figure 6, significant activation by wearing a denture prosthesis was focused on the right lateralized dorsal prefrontal cortex area in accordance with the anatomical structures of the surface cortex (Figure 6-b).

Jaw Muscle EMG and Jaw Motion Activities While Not Wearing and Wearing Partial Dentures

Features of Jaw Muscle EMG and Motion Activities

Representative data showing jaw muscle EMG activity and jaw motion activity during masticatory performance under the not wearing and wearing conditions are shown in Figure 7. Wearing a partial denture prosthesis clearly presented the increased activity in the jaw closing muscles as compared with the features of the not wearing condition.







Figure 7. Representative features of jaw muscle EMG and jaw motion activities under the not wearing (a) and wearing (b) conditions. EMG activities of the jaw muscles were clearly increased while wearing a partial denture as compared with the not wearing condition.

Jaw Muscle EMG Activities

Cycle duration was not significantly changed by wearing a partial denture prosthesis as compared



Black & White

Fig. 8/c




Figure 8. EMG muscle activities. (a) Cycle duration was not significantly changed by wearing a partial denture prosthesis as compared with the not wearing condition. (b) Burst duration, (c) area, and (d) peak amplitude of EMG activity during the chewing task were significantly increased in Mm and/or Ta while wearing a partial denture as compared with the not wearing condition.

In contrast, statistical comparisons of jaw muscle EMG activities between the not wearing and wearing conditions revealed significant increases in the EMG findings for burst duration, area, and peak amplitude in regard to chewing-related jaw muscle EMG activities in the Mm and/or Ta muscles during partial denture wearing (Figure 8-b~d).

Jaw Motion Activities

Vertical and anterior/posterior components of the jaw motion activities, as well as jaw motion velocity were not significantly changed by wearing a partial denture prosthesis as compared with the not wearing condition.




Figure 9. (continued).





Figure 9. Jaw motion activities. (a) The vertical component of jaw motion activity was not significantly changed by wearing a partial denture prosthesis as compared with the not wearing condition. (b) The lateral component of jaw motion activity while wearing a partial denture prosthesis was significantly increased as compared with the not wearing condition.

In contrast, a significant increase was seen for the lateral component of jaw motion activity while wearing a partial denture as compared with the not wearing condition (Figure 9-b).

Occlusal Contact

Findings for total occlusal contact while not wearing and wearing a partial denture prosthesis are presented in Figure 10. Significant increases were seen for total occlusal force and total occlusal area when wearing a partial denture prosthesis as compared with the not wearing condition.



Black & White

Fig. 10/a




Figure 10. Occlusal contact. Significant increases of total (a) occlusal force and (b) occlusal area were seen while wearing a partial denture prosthesis as compared with the not wearing condition.

Black & White

Fig. 10/b

Masticatory Score

Significantly increased masticatory scores, which may represent subjective awareness of chewing ability, were seen when wearing a partial denture prosthesis as compared with the




Figure 11. Masticatory score. Significant increases in masticatory scores were seen while wearing a partial denture prosthesis as compared with the not wearing condition.

Black & White

Fig. 11

Discussion

Features of Prefrontal Activity when Wearing Partial Dentures and their Functional Significance

In the present study, we found that wearing a partial denture prosthesis significantly activated the dorsal prefrontal cortex as well as jaw functions, such as jaw muscle EMG and jaw motion activities. It has been reported that the dorsal prefrontal cortex is involved in various higher cognitive functions, such as movement planning and execution [20] [21] [22], sensory processing [23] [24] [25] [26] [27], learning and memory [28] [29], reward [30] [31] [32], and attention [33] [34]. According to data obtained in our previous study, which was presented in part at the IADR annual meeting in Toronto in 2008 [35], deafferentation of oral somatosensory input may cause modulated chewing behavior and deactivation of the dorsal prefrontal cortex during masticatory performance, which was especially predominant in the right lateralized prefrontal cortex area. Furthermore, chewing imagery also activated the dorsal prefrontal cortex area, bilaterally (unpublished data).

Taken together, we consider that the dorsal prefrontal cortex may be involved in execution, as well as somatosensory processing and sensorimotor integration during chewing performance.

Possible Relationships among Prefrontal Cortex Activity, Jaw Muscle EMG and Jaw Motion Activities, as well as Occlusal Contact Features, and Masticatory Score

According to our recent data analysis, there are significant correlations between mastication-related prefrontal cortex activity, feature in the occlusal contact, and masticatory score considered as the self-evaluation of the masticatory function by wearing partial denture prosthesis. These findings will be presented at the Neuroscience Meeting in 2011 (Abstract submitted) [36]. The prefrontal cortex plays a role in the cognitive process, such as working memory, and enables humans to hold the contents of conscious awareness by maintaining an active representation of information [37]. From these findings, it is considered that occlusal reconstruction obtained by wearing a partial denture prosthesis may contribute to improved conscious awareness of chewing ability and the related prefrontal cortex activity.

Clinical Implications

Stein et al. [5] have examined Oral Dis.ease as a potential risk factor in the development of dementia and reported that a low number of teeth increased the risk of developing certain forms of dementia. Miura et al. [6] also found that number of teeth, maximum bite force, and masticatory scores in cognitively impaired subjects were significantly lower than those in healthy elderly subjects. Recently, Weijenberg et al. [38] reviewed the causal relationship between mastication and cognition in animal and human experimental studies, and suggested that correlations exist between mastication and activities of daily living and nutritional status.

Furthermore, Hirai et al [39] also reported a relationship between masticatory score and cognitive deterioration, that is, masticatory score was significantly correlated to dementia score (Hasegawa test) in 80 elderly subjects.

Taken together, it is considered that wearing an appropriate denture prosthesis leads to improved conscious awareness of chewing ability, which might be related to prevention of neurological cognitive deterioration in aged individuals.

Conclusion

In the present study, occlusal reconstruction by wearing a partial denture prosthesis improved prefrontal cortex activity and the conscious awareness of chewing ability, as well as jaw functions as shown by increased jaw muscle EMG and jaw motion activities. From these, it is assumed that appropriate denture wearing might have an impact on not only jaw functions, but also higher brain functions closely related to masticatory performance.

Acknowledgments

This research was partially supported by a Grant-in-Aid for Scientific Research (C), No. 21592473, 2009~2011, from the Ministry of Education, Science, Sports and Culture of Japan.

References

- [1] Miura H, Miura K, Mizugai H, Arai Y, Umenai T, Isogai E. Chewing ability and quality of life among the elderly residing in a rural community in Japan. *J. Oral. Rehabil.*, 2000 27, 731–734.
- [2] Takata Y, Ansai T, Awano S, Hamasaki T, Yoshitake Y, Kimura Y, Sonoki K, Wakisaka M, Fukuhara M, Takehara T. Relationship of physical fitness to chewing in an 80-year-old population. *Oral. Dis.*, 2004 10, 44–49.
- [3] Shimazaki Y, Soh I, Saito T, Yamashita Y, Koga T, Miyazaki H, Takehara T. Influence of dentition status on physical disability, mental impairment, and mortality in institutionalized elderly people. *J. Dent Res.*, 2001 80, 340–345.
- [4] Stein PS, Desrosiers M, Donegan SJ, Yepes JF, Kryscio RJ. Tooth loss, dementia and neuropathology in the Nun study. *J. Am. Dent. Assoc.*, 2007 138, 1314–1322.
- [5] Miura H, Yamasaki K, Kariyasu M, Miura K, Sumi Y. Relationship between cognitive function and mastication in elderly females. *J. Oral. Rehabil.*, 2003 30, 808–811.
- [6] Koshino H, Hirai T, Ishijima T, Tsukagoshi H, Ishigami T, Tanaka Y. Quality of life and masticatory function in denture wearers. *J. Oral. Rehabil.*, 2006 33, 323–329.
- [7] Rajah MN, D’Esposito M. Region-specific changes in prefrontal function with age: a review of PET and fMRI studies on working and episodic memory. *Brain*, 2005 128, 1964–1983.

- [8] Huang CS, Hiraba H, Sessle BJ. Input-output relationships of the primary face motor cortex in the monkey (*Macaca fascicularis*). *J. Neurophysiol.*, 1989 61, 350-362.
- [9] Huang CS, Hiraba H, Murray GM, Sessle BJ. Topographical distribution and functional properties of cortically induced rhythmical jaw movements in the monkey (*Macaca fascicularis*). *J. Neurophysiol.*, 1989 61, 635-650.
- [10] Murray GM, Sessle BJ. Functional properties of single neurons in the face primary motor cortex of the primate. III. Relations with different directions of trained tongue protrusion. *J. Neurophysiol.*, 1992 67, 775-785.
- [11] Lin LD, Murray GM, Sessle BJ. Functional properties of single neurons in the primate face primary somatosensory cortex. I. Relations with trained orofacial motor behaviors. *J. Neurophysiol.*, 1994 71, 2377-2390.
- [12] Martin RE, Kempainen P, Masuda Y, Yao D, Murray GM, Sessle BJ. Features of cortically evoked swallowing in the awake primate (*Macaca fascicularis*). *J. Neurophysiol.*, 1999 82, 1529-1541.
- [13] Narita N, Yamamura K, Yao D, Martin RE, Sessle BJ. Effects of functional disruption of lateral pericentral cerebral cortex on primate swallowing. *Brain Res*, 1999 824, 140-145.
- [14] Narita N, Yamamura K, Yao D, Martin RE, Masuda Y, Sessle BJ. Effects on mastication of reversible bilateral inactivation of the lateral pericentral cortex in the monkey (*Macaca fascicularis*). *Arch. Oral. Biol.*, 2002 47, 673-688.
- [15] Yao D, Yamamura K, Narita N, Murray GM, Sessle BJ. Effects of reversible cold block of face primary somatosensory cortex on orofacial movements and related face primary motor cortex neuronal activity. *Somatosens Mot. Res.*, 2002 19, 261-271.
- [16] Yamamura K, Narita N, Yao D, Martin RE, Masuda Y, Sessle BJ. Effects of reversible bilateral inactivation of face primary motor cortex on mastication and swallowing. *Brain Res*, 2002 944, 40-55.
- [17] Sessle BJ, Adachi K, Avivi-Arber L, Lee J, Nishiura H, Yao D, Yoshino K. Neuroplasticity of face primary motor cortex control of orofacial movements. *Arch. Oral. Biol.*, 2007 52, 334-337.
- [18] Narita N, Kamiya K, Yamamura K, Kawasaki S, Matsumoto T, Tanaka N. Chewing-related prefrontal cortex activation while wearing partial denture prosthesis: pilot study. *J. Prosthodont Res.*, 2009 53, 126-35.
- [19] Hirai T, Anzai T, Kaneda K, Matai N, Tanaka O, Ikeda Y, Uchida T. Evaluation method for masticatory function of complete denture wearer-by questionnaire with 35 foods listings. *Nihon Hotetsu Shika Gakkai Zasshi*, 1988 32, 1261-1267.
- [20] Hunter MD, Green RD, Wilkinson ID, Spence SA. Spatial and tempOral. Dis.sociation in prefrontal cortex during action execution. *Neuroimage*, 2004 23, 1186-1191.
- [21] Jenkins IH, Jahanshahi M, Jueptner M, Passingham RE, Brooks DJ. Self-initiated versus externally triggered movements. II. The effect of movement predictability on regional cerebral blood flow. *Brain*, 2000 123, 1216-1228.
- [22] Lewis PA, Wing AM, Pope PA, Praamstra P, Miall RC. Brain activity correlates differentially with increasing temporal complexity of rhythms during initialisation, synchronisation, and continuation phases of paced finger tapping. *Neuropsychologia*, 2004 42, 1301-1312.

-
- [23] Bolton DA, Staines WR. Transient inhibition of the dorsal prefrontal cortex disrupts attention-based modulation of tactile stimuli at early stages of somatosensory processing. *Neuropsychologia*, 2011 49, 1928-1937.
 - [24] Anderson JS, Ferguson MA, Lopez-Larson M, Yurgelun-Todd D. Topographic maps of multisensory attention. *Proc. Natl. Acad. Sci. USA*, 2010 107, 20110-20114.
 - [25] Coghill RC, Gilron I, Iadarola MJ. Hemispheric lateralization of somatosensory processing. *J. Neurophysiol.*, 2001 85, 2602–2612.
 - [26] Stoeckel MC, Weder B, Binkofski F, Buccino G, Shah NJ, Seitz RJ. A fronto-parietal circuit for tactile object discrimination: an event-related fMRI study. *Neuroimage*, 2003 19, 1103–1114.
 - [27] Harada T, Saito DN, Kashikura K, Sato T, Yonekura Y, Honda M, Sadato N. Asymmetrical neural substrates of tactile discrimination in humans: a functional magnetic resonance imaging study. *J. Neurosci.*, 2004 24, 7524–7530.
 - [28] Abe H, Lee D. Distributed coding of actual and hypothetical outcomes in the orbital and dorsal prefrontal cortex. *Neuron*, 2011 70, 731-741.
 - [29] Celone KA, Thompson-Brenner H, Ross RS, Pratt EM, Stern CE. An fMRI investigation of the fronto-striatal learning system in women who exhibit eating disorder behaviors. *Neuroimage*, 2011 56, 1749-1757.
 - [30] Wallis JD, Miller EK. Neuronal activity in primate dorsal and orbital prefrontal cortex during performance of a reward preference task. *Eur. J. Neurosci.*, 2003 18, 2069–2081.
 - [31] Fallon S, Shearman E, Sershen H, Lajtha A. Food reward-induced neurotransmitter changes in cognitive brain regions. *Neurochem. Res.*, 2007 32, 1772–1782.
 - [32] Inoue M, Oomura Y, Aou S, Nishino H, Sikdar SK. Reward related neuronal activity in monkey dorsal prefrontal cortex during feeding behavior. *Brain Res.*, 1985 326, 307-12.
 - [33] Toichi M, Findling RL, Kubota Y, Calabrese JR, Wiznitzer M, McNamara NK, Yamamoto K. Hemodynamic differences in the activation of the prefrontal cortex: attention vs. higher cognitive processing. *Neuropsychologia*, 2004 42, 698–706.
 - [34] Tucha L, Simpson W. The role of time on task performance in modifying the effects of gum chewing on attention. *Appetite*, 2011 56, 299-301.
 - [35] Narita N, Kamiya K, Yamamura K, Funto M. Effects of oral deafferentation to chewing-related prefrontal cortex activity. (*Abstract of IADR Annual Meeting in Toronto*, 2008).
 - [36] Narita N, Kamiya K. Association between jaw functions and prefrontal cortex activity by wearing denture prosthesis. (*Abstract of Neuroscience Annual Meeting in Washington DC*, 2011).
 - [37] Courtney SM, Petit L, Haxby JV, Ungerleider LG.. The role of prefrontal cortex in working memory: examining the contents of consciousness. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 1998 353, 1819-1828.
 - [38] Weijenberg RA, Scherder EJ, Lobbezoo F. Mastication for the mind-the relationship between mastication and cognition in ageing and dementia. *Neurosci. Biobehav. Rev.*, 2011 35, 483-497.
 - [39] Hirai T, Ikeda Y, Yajima T, Ichioka N, Koshino H, Ishijima T, Tamura T, Tanaka O, Takeda H. Masticatory function and physical/mental activity in the Elderly, *Higashi Nippon Dental Journal*, 1995 14, 133–142.

Chapter IV

Inlay Retained Fixed Partial Dentures

Sherif Adel Mohsen *

Chairman of Department of Fixed Prosthodontics
Faculty of Dentistry, Minia University, Minia, Egypt

Abstract

Patients often refuse these options since the reasons of high cost and/or surgical intervention. The full coverage retainers for tooth supported fixed partial denture is a conventional approach to replace one or several missing teeth. Two types of full coverage retainers FPDs are used in dental clinics: porcelain fused to metal and all ceramic. Porcelain fused to metal provides great fracture strength and relatively good long-term clinical success. One of significant disadvantages of the porcelain fused to metal, as well as modern all ceramic FPDs is that large amount of sound tooth substance of the abutments needs be removed in FPD fabrication.

An inlay-retained FPD (IRFPD) is, however, a less-invasive treatment modality and a more conservative option for restoration of damaged teeth, because it requires minimal tooth reduction, preserves healthy tooth structure, and maintains the periodontal tissue's integrity. Inlay-retained FPDs are, therefore, alternatives to both anterior and posterior complete coverage conventional restorations. Inlay retained FPDs can be constructed by using dental alloys, ceramic materials, and fiber-reinforced composite. Several types of preparations for inlay retained FPDs are used: Occluso – proximal inlay, Tub-shaped , Proximal box, Rest seat on the occlusal surface, Lingual tooth reduction, and Retentive-slot preparations. The size of these preparation features depends on the size of the tooth.

Several researches were carried out about the effect of various designs, materials, connectors size on the fracture resistance, marginal integrity and retention of inlay retained FPDs.

* E-mail address: cherif.mohsen@gmail.com.

Introduction

Although implant-supported FPDs are high quality alternatives to the tooth supported FPDs, patients often refuse these options due to the reasons of high cost and/or surgical intervention [1]. The full coverage retainers for tooth supported fixed partial denture is a conventional approach to replace one or several missing teeth [2]. Two types of full coverage retainers FPDs using ceramics are used in dental clinics : porcelain fused to metal and all ceramic. Porcelain fused to metal provides great fracture strength and relatively good long-term clinical success. One of significant disadvantages of the porcelain fused to metal, as well as modern all ceramic FPDs is that large amount of sound tooth substance of the abutments needs to be removed in FPD fabrication [3].

Properties

An inlay-retained FPD (IRFPD) is a less-invasive treatment modality and a more conservative option for restoration of damaged teeth, as it requires minimal tooth reduction so it preserves healthy tooth structure. Inlay-retained FPD finish line is always supra-gingival maintaining the periodontal tissue's integrity [4-7]. Also, inlay-retained FPD is highly esthetic restoration when it is fabricated from ceramic or fiber-reinforced composite [8,9]. Recent, studies showed that inlay-retained FPD showed some drawbacks irrespective to the type of the FPD materials; decreased retention, high marginal discrepancies especially with machinable ceramics [10]. Inlay-retained FPDs are, therefore, alternatives to both anterior and posterior complete coverage conventional restorations, especially with the introduction zirconia ceramics which enables the construction of durable posterior FPDs [11,12]. However inlay-retained FPD is contra-indicated in mutilated, pulpless teeth, in above average functional occlusal forces and when control of moisture contamination is difficult.

Materials

Inlay retained FPDs can be constructed by using dental alloys, fiber-reinforced composite and ceramic materials [3,13]. Clinical results for metal inlay-retained FPDs have been favorable, however the visibility of the metal retainer and the change in natural tooth translucency are aesthetically unfavorable [12].

Inlay retained FPDs may be fabricated from fiber-reinforced composites (FRC). Physical data on reinforced composites suggest that these materials are best suited for conservative inlay FPDs [14]. With a carefully executed bonding technique, good results in marginal adaptation have been achieved with composite inlays. The stress-resistant marginal integrity of composite inlays has been attributed to their dentin-like elasticity modulus [15,16]. FRC inlay-retained FPDs are considered as minimal invasive, metal free, and good aesthetic restorations for replacing missing single or multiple anterior or posterior teeth [17]. Inlay-retained resin-bonded FRC FPDs can be used especially for patients having occlusal or proximal carious lesions or existing fillings in teeth adjacent to the edentulous space. The

abutment teeth are prepared for retaining the FPD without excessive removal of sound tooth substance. Several types of FRC frameworks were used and it was suggested that high-volume fraction frameworks provide better clinical success than low-volume fraction frameworks due to lack of support for the veneering composite of the pontics [18]. Although promising clinical results have been reported for such restorations, these suffer from the disadvantages of unstable aesthetics and problems with wear of the veneering composite. Also the pontic region is considered as a weak point for the fiber-reinforced composite inlay-retained FPD. This may be due to the fact that in occlusal loading conditions where the load is applied to the cusps of posterior teeth instead of central fossa, the delamination of veneering composite can occur at lower load levels [19].

Factors affecting the durability of FRC restorations include the properties of the fibers, matrix, and polymer, impregnation of fibers with the resin, adhesion of fibers to the matrix, the quantity of fibers, and the direction, orientation, location, construction, distribution, and position of the fibers [9,20]. The fiber content not only determines the fiber properties, but so does the matrix composition and the bond between fibers and the matrix [21]. Placement of fibre at or slightly away from the tensile side improved the flexural properties of the composite [1,22]. Fibers used for reinforced composites can be glass-fiber or polyethylene-fiber. Fracture resistance of restorations made of glass-fiber reinforced composite is higher than those of polyethylene fiber-reinforced composite restorations [1,9].

It has been reported that fiber-reinforced FPDs have the necessary fracture strength to withstand the average adult masticatory forces in the molar region, with recorded high fracture strength of approximately 700 N after thermo-cycling and mechanical loading [14]. The basic design of a FRC inlay-retained FPDs preparation is a box-shaped preparation used to achieve optimal extension of the fiber framework and retention [23]. A tub-shaped preparation may be indicated when there is insufficient space to prepare the box or to prevent irritation of the pulp of tipped teeth. Generally, FRC inlay-retained FPDs have been used for 3-unit FPDs replacing a single premolar or molar when the intra-abutment span does not exceed 15 mm [14].

Ceramics inlay-retained FPDs with different strong core are used. They were made either from glass infiltrated alumina ceramic (In-Ceram Alumina), pure alumina ceramic (Procera) or lithium-disilicate based glass-ceramic (IPS Empress 2, IPS e-max) [6]. Recently, another ceramic, yttrium-oxide partially stabilized zirconia (YPSZ) has been made available to dentistry through the CAD/CAM-technique (ice zircon, IPS e-max Zr CAD) or through pressing technique (IPS e-max Zr Press). Using this core material for all-ceramic frameworks, excellent mechanical performance and superior strength and fracture resistance, compared to other ceramics, have been shown [24-26].

Mode of Failure

Stress applied during mastication may range between 441 N and 981 N, 245 N and 491 N, 147 N and 368 N, and 98 N and 270 N in the molar, premolar, canine, and incisor regions, respectively. A restoration should be able to withstand stress to approximately 500 N in the premolar region and 500 N to 900 N in the molar region.

The mode of fracture is a good indicator of the path of crack propagation [27,28]. Mechanical failure of restorations may be due to several factors such as wrong treatment plane and case selection, the properties of the luting agent, the thickness of restoration, the preparation design and the thickness of connectors [8].

The mode of failure of the metal ceramic inlay-retained FPDs is ceramic chipping, which may be due to weak bond between metal and ceramic or due to excessive strength beyond the bearing capacity of the veneering ceramic materials [4,6].

The mode of failure of the fiber-reinforced inlay-retained FPD exhibits a 2-phase pattern, which consists of cracking and chipping of the veneer layer followed by adhesive failure between the veneer and fiber materials. The fracture line, running between the border of the fiber framework and the veneering material, also appeared in the facing material itself, with no evidence of fiber burst [29,30]. In general, stress concentrations within the resin and the interface are relieved by initiation of a crack and propagation of the crack through the resin until it meets the fiber, resulting in debonding of the resin composite. This means that the bond strength between the fiber and facing composite, rather than the strength of the fiber framework itself, appeared to have had the most influence on the fracture resistance. Therefore, it may be necessary to apply a silane-coupling agent to the fiber framework in order to achieve improved chemical bonding [30]. One of the reasons for the adhesive fracture of the restorations was the difference in the elastic modulus between the fiber framework and the veneering composite [31,32]. The flexure strength of fiber-reinforced restorations might be improved with the use of new polymer formulations with high filler particle distribution however, these materials are not suitable for chair-side use. The weakest parts of the restoration were in the cohesive strength of the veneering composite and the bond between the fiber composite framework. The direction of the failures was primarily in the MD direction, indicating that unidirectional fibers change the path of the crack [1,9].

The failure mode observed for all ceramic inlay-retained FPD is delamination and chipping of the veneering material. These observations indicate that the weak points ceramic inlay-retained FPD are the adhesion between the framework and the veneering materials and the strength of the veneering material itself. High fracture strength in case of zirconia ceramic core may be due to the fact that the core is a YPSZ consisting of partially stabilized zirconia particles densely sintered, resulting in a final microstructure in which voids, flaws, and cracks are reduced to a minimum. Also, the transformation toughening mechanisms increase the fracture strength of the material. On the other hand, adhesive failure between veneer and ceramic does not occur in the presence of a good bond between a compatible ceramic core and the veneering material [4-6,8-10].

Fracture of retainers were observed by some researchers, in case of 4x4 mm connector size, whereas in smaller connector size, fracture may happens not also at the retainer but also at the connector. So it is recommended to use a 4x4 mm connector size, in case of inlay-retained FPD [6].

The debonding rate of the long span FPDs was slightly higher than that of the short span FPDs. Debonding in the premolar location appeared to be caused by the smaller bonding area and the narrow connector dimension. When a load was applied to the center of the pontic, traction forces caused a lever effect on the terminal interface of the restorations. It is possible that the bending under load was the cause of debonding failure. It is likely that the box-shaped design is strong enough to resist the bending force due to the additional 2-mm wall

support of the proximal box. The increased debonding of the long span FPDs relative to the short span FPDs could be explained by their flexibility [4,9,10].

Preparations

Several types of preparations for inlay retained FPDs are used : Occluso – proximal inlay (figure 1-3), Tub-shaped (figure 4-6) , Proximal box (figure 7-9), Rest seat on the occlusal surface with or without Lingual tooth reduction and retentive-slot preparations and finally a palatal inlay with or without a proximal grooves. The size of these preparation features depends on the size of the tooth [1,3,4,8,9,12,14,33-35].




Figure 1. A schematic illustration for an inlay-proximal inlay retained partial denture.



Figure 2. An inlay-proximal inlay retained partial denture preparation.



Figure 3. An inlay-proximal inlay retained partial denture.




Figure 4. A schematic illustration for an tub design inlay retained partial denture.

The occluso-proximal inlay is composed of an occlusal inlay and a proximal box. The occlusal inlay must remove any carious or undermined enamel. For an intact tooth, the depth of the occlusal preparation must be of 2.0 mm for the ceramic. The occlusal preparation is 4 mm wide and extended 4 or 6 mm mesiodistally for the premolar or molar models, respectively.

The proximal box extends 1mm wide and had approximately 6° divergences, extending 2 mm apical to the isthmus floor.

The tub-shaped preparation consisted of an occluso-proximal inlay has the same dimension as the occlusal preparation of an occluso – proximal inlay. While the proximal box inlay retained FPDs has the same dimension of the proximal box in case of occluso-proximal inlay [9].



Figure 5. A tub shaped inlay retained partial denture preparation.



Figure 6. A tub inlay retained partial denture.




Figure 7. A schematic illustration for an proximal box design inlay retained partial denture.



Figure 8. A proximal box inlay retained partial denture preparation.




Figure 9. A proximal box inlay retained partial denture.

The rest seat is prepared on the occlusal surface of each abutment with minimal preparation on the proximal wall of the abutments. The drawback of this design is that the pontic may rotate along an axis formed by the 2 rest seats when occlusal force acts on the occlusal incline of the pontic [34].

Rest seat on the occlusal surface and lingual tooth reduction preparation can be performed by proximal and lingual minimal reduction of the abutments, one rest seat prepared on the occlusal surface of each abutment. Proximal boxes are also prepared. The facial and lingual walls of these proximal boxes are slightly convergent toward the occlusal surface to provide “undercut retention.” Reduction of the proximal surfaces was accomplished to ensure that the gingival and facial margins of the boxes were completely contained within the proximal reduction area. The grooves on proximal and palatal/lingual surfaces of abutment teeth serve 2 main functions: to define the path of insertion and to provide retention and resistance form to the retainer against the dislodging forces acting on the pontic. When designed with mesial and distal occlusal rests alone, the pontic may rotate along an axis formed by the 2 rest seats when occlusal force acts on the occlusal incline of the pontic [35].

Palatal inlay with or without a proximal grooves preparation is indicated in anterior teeth, the aim of the proximal grooves preparation is to enhance the retentivity of the preparation.

Fracture Resistance

New core materials in combination with new preparation designs have to be tested before they can be recommended for clinical use. It is not known what fracture resistance is needed to achieve a good long-term outcome of IRFPDs in the molar region. Numerous authors investigated the maximum bite forces during mastication. Mean values for the maximum bite force level varied from 98 to 981N. The highest bite force was found in the first molar region.

Reviewing the literature, Körber and Ludwig summarized that posterior FPDs should be strong enough to withstand a load of 500N [26,27]. Additionally, cyclic fatigue loading and stress corrosion fatigue caused by the oral environment has to be considered, because they can considerably weaken the fracture resistance of all-ceramic restorations [36]. Under the conditions of the oral environment, the inherent flaws of ceramic materials act as the origin of crack propagation and can grow to critical sizes [37,38]. The endurance limit for fatigue cycling that can be applied to dental ceramics is approximately 50% of the maximal fracture strength [39]. Therefore it seems reasonable to assume that an initial fracture resistance of 1000N should be required for a favorable clinical prognosis of posterior all ceramic IRFPDs. However, the results of quasi-static fracture tests should be related to dynamic fatigue in a chewing simulator combined with thermo-cycling. According to the literature, fatigue produced by 240,000 cycles in a chewing simulator corresponds to a period of 1 year of clinical service [40]. Therefore 1,200,000 chewing cycles were performed to simulate a service time of 5 years [41].

Material used in the construction of inlay retained FPD, specially the core materials, plays an important factor in the fracture resistance of the restoration as well as the veneering materials and its bonds with the core materials [26,42-45]. Zirconia ceramic core and ceramic veneering inlay retained FPD showed in several studies a significant higher fracture resistance than that of ceramo-metallic inlay retained FPD [4,6].

Connector size is an important factor of fracture resistance of inlay-retained FPD according to the type of core material used. In case of metal core or yttrium-oxide partially stabilized zirconia ceramic, a 3x3 mm or 4x4 mm connector size are clinically successful. As for lithium disilicate glass ceramic or fiber-reinforced composite a 4x4 mm connector size is indicated [6,46].

The occluso – proximal inlay shaped design provides greater surface area to resist the forces and a larger connector dimension of the framework, thereby resisting the bending forces better than the other inlay retained FPD designs. Also, in case of fiber reinforced composite, this design allows for a higher fiber content therefore, it might have provided a greater reinforcement effect near the tensile side and thus a higher fracture strength [5,6,9,19].

Studies in dental literature showed that in case of fiber reinforced composite, increasing span length correlated with the lower fiber-reinforced FPD strength. All FPDs flex slightly when subjected to a load. In other words, the longer the span is, the greater the flexing will be. However, the relationship between deflection and span length is not simply linear but varies with the cube of the span length. Excessive flexing under occlusal loads may cause

failure of a long-span FPD. It can lead to fracture of the veneering material, breakage of a connector, loosening of a retainer, or an unfavorable soft tissue response. Therefore, the application of a fiber-reinforced inlay FPD might be advisable for short pontic spans [9,11,19].

Marginal Adaptation

With a carefully executed bonding technique, good results in marginal adaptation have been achieved. In case of marginal imperfections, the adhesive functions as a second barrier against penetration of bacteria in dentin [11]. In case of fiber reinforced composite, although some authors concluded that with a carefully executed bonding technique, beveling of preparation finishing lines is not necessary, it is accepted that with the shrinking of composite materials, beveling of enamel decreases the risk of marginal gaps, microleakage, and enamel fractures. Beveling of enamel increases the loss of dental hard tissue; however, it may improve marginal adaptation in situations in which difficulties in accessing the margin in the gingival crevice are expected [47-49].

Adequate results in marginal adaptation are achieved with the use of the total bonding technique in case of canines. The geometry of the cavity preparation, which was unfavorable for selective bonding without beveling, may have provided the benefits of total bonding at insertion. However, the total bonding technique cannot be recommended for molars [11]. For these box-shaped cavity preparations, an increase of the bonded surface resulted in severely diminished flow. The contractive stresses can exceed adhesive strength and cause marginal imperfections after loading.

The marginal quality attained for molars with selective bonding can still be compared with the marginal quality of composite inlays. C-factor (ratio of bonded to free, un-bonded restoration surfaces) is reduced with the use of the selective bonding technique. After the final post-curing, there are only a few carbon double bonds available for the bond between the resin-cement and the crowns.

Therefore, all systems recommend roughening and silane-coating of the inner-crown surface before their adhesive cementation [23,50,51]. Some authors state that glass fibers bond better to silane coupling agents than to a composite surface. The composite surface contains inorganic fillers which bond to silane-coupling agents, but the extent of the bonding area depends on the filler content [11].

The marginal discrepancy of each restoration may vary greatly at different location, due to the difference in the preparation geometry, marginal configurations and the location and orientation of tubules. The inlay-shaped retained FPD shows a significant higher vertical marginal gap measurements than all the other inlay retained FPD designs due to the fact that it possess a much longer perimeter than them [10].

The technique of construction also plays an important role in the marginal adaptation of the inlay retained FPD. In case of ceramic, the machinable ceramic showed a less marginal adaptation than pressable.

This may be attributed to geometrical design of the restoration and difficulties regarding scanning, digitization, and the milling process of brittle ceramic material. Moreover, the

adaptation of restorations made out of milled ceramic blanks may be affected by the size of milling burs, and material conditions during the milling procedure [52].

Retention

The longevity of fixed prostheses depends on the retention and marginal integrity of restorations. Many factors have been demonstrated to influence the retention of fixed prosthetic appliances, such as the size and shape of prepared teeth, manipulation of cement, retentive properties of cement, cement film thickness, relieving space or venting for cement, cement application, roughness of dentinal surface, the convergence and the preparation height [4,8,9,12]. Dental literature is poor on studying the retentive strength of the inlay retained ceramic FPDs. As regard the forces needed to dislodge the inlay retained FPDs, a significant difference was found between the different designs. The inlay-shaped design was significant higher than the other inlay retained FPDs designs due to the long surface area of bonding between the restoration and the tooth. In a recent studies, a comparison was carried out between the inlay retained FPD and the full coverage FPD who recorded much higher retention. These difference in retention between the two tested FPDs may be attributed to the fact that longer preparations (full coverage FPDs) have more surface area and therefore are more retentive than the shorter preparations (inlay retained FPDs). Another reason for these differences may be due to the frictional resistance between a prepared tooth and ceramic crown is as important as the adhesive bond strength [10]. To increase the bond between the restoration and the tooth, the author recommends to subject the inner surface of the restoration to surface treatments, the same used for resin bonded retainers.

Clinical Survival Rate

Clinical survival rate of the different designs of inlay retained FPD is the main important factor to evaluate the success of this type of restoration. Strength and durability of inlay retained FPDs are highly dependent on the inlay designs, sizes, pontic span, connectors' size. All these variables are to be set according to the material used. Therefore, proper selection of the restorative materials and careful preparation designs are essential, especially in the posterior region. Ohlmann et al (2008) [12] in a preliminary clinical study on all-ceramic inlay-retained fixed partial dentures, reported that 6 out of 13 inlay retained FPDs were subjected to debonding.

Harder et al (2010) [53] in an eight-year outcome study of posterior inlay-retained all-ceramic fixed dental prostheses, reported that the percentage of debonding of this type of FPDs was 15%. In two clinical evaluations, Edelhoff et al. (2001) [3] and Edelhoff and Sorensen (2002) [7], reported that compared to crown retained FPDs, de-bonding of inlay retained FPDs appear to be much too high.

Conclusion

Inlay retained FPDs are conservative treatments for replacing a single missing tooth that can be constructed from metal, metal-ceramic, fiber-reinforced composites (FRC) and ceramic. With the evolution of ceramic and specially zirconia ceramic, it is possible to use ceramic inlay retained FPDs in the posterior quadrant. Although many preparation designs can be done for inlay retained FPDs, the inlay shaped preparation recorded the highest fracture resistance but it recorded the highest vertical marginal gap measurements. Retention of inlay retained FPDs is one of the drawbacks of this type of restoration, the author recommends to subject the inner surface of the restoration to surface treatments, the same used for resin bonded retainers. Few clinical survival rate studies were carried out showing that de-bonding of inlay retained FPDs is higher than crown retained FPDs.

References

- [1] Xie Q, Lassila LVP, Vallittu PK. Comparison of load-bearing capacity of direct resin-bonded fiber-reinforced composite FPDs with four framework designs. *J. Dent.*, 2007 35, 578-582.
- [2] Freilich MA, Karmaker AC, Burstone CJ, Goldberg AJ. Development and clinical application of a light-polymerized fiber-reinforced composite. *J. Prosthet Dent*, 1998 80, 311- 318.
- [3] Edelhoff D, Spiekermann H, Yildirim M: Metal-free inlay-retained fixed partial dentures. *Quintessence Int.*, 2001 32, 269-281.
- [4] Kilicarslan MA, Kedici PS, Kucukcemen HC, Uludag BC. In vitro fracture resistance of posterior metal-ceramic and all-ceramic inlay-retained resin-bonded fixed partial dentures. *J. Prosthet Dent*, 2004 9, 365-370.
- [5] Mohsen SA. Fracture resistance of three ceramic inlay retained fixed partial denture designs. An in-vitro comparative study. *J. Prosthodont*, 2010 19, 531-540.
- [6] Mohsen SA. Effect of preparation design and connector size on the fracture resistance of metal ceramic and all ceramic inlay retained fixed partial denture. *EDJ*, 2010 56, 699.
- [7] Edelhoff D, Sorensen JA. Tooth structure removal associated with various preparation designs for anterior teeth. *J. Prosthet Dent*, 2002 87, 503–509.
- [8] Wolfart S, Ludwig K, Uphaus A, Kern M. Fracture strength of all-ceramic posterior inlay-retained fixed partial dentures. *Dent. Mater.*, 2007 23, 1513-1520.
- [9] Song HY, Yi YJ, Cho LR, Park DY. Effects of two preparation designs and pontic distance on bending and fracture strength of fiber-reinforced composite inlay fixed partial dentures. *J. Prosthet Dent*, 2003 90, 347-353.
- [10] Mohsen SA. Vertical marginal gap and retention of ceramic full coverage and inlay retained fixed partial dentures. *EDJ*, 2011 57, in press.
- [11] Göehring TN, Peters OA, Lutz F. Marginal adaptation of inlay-retained adhesive fixed partial dentures after mechanical and thermal stress: An in vitro study. *J. Prosthet Dent*, 2001 86, 81-92.

-
- [12] Ohlmann B, Rammelsberg P, Schmitter M, Schwarz S, Gabbert O. All-ceramic inlay-retained fixed partial dentures: preliminary results from a clinical study. *J. Dent.*, 2008 36, 692-696.
- [13] Göehring TN, Mormann WH, Lutz F. Clinical and scanning electron microscopic evaluation of fiber-reinforced inlay fixed partial dentures: preliminary results after one year. *J. Prosthet Dent*, 1999 82, 662-668.
- [14] Behr M, Rosentritt A, Leibrock S, Schneider-Feyrer S, Handel G. In-vitro study of fracture strength and marginal adaption of fibre-reinforced adhesive fixed partial inlay dentures. *J. Dent.*, 1999 27, 163-168.
- [15] Krejci I, Lutz F, Gautschi L. Wear and marginal adaptation of composite resin inlays. *J. Prosthet Dent*, 1994 72, 233-44.
- [16] Braem MJ, Lambrechts P, Gladys S, Vanherle G. In vitro fatigue behavior of restorative composites and glass ionomers. *Dent. Mater.*, 1995 11, 137- 41.
- [17] Kolbeck C, Rosentritt M, Behr M, Lang R, Handel G. In vitro study of fracture strength and marginal adaptation of polyethylene-fiber-reinforced-composite versus glass-fiberreinforced-composite fixed partial denture. *J. Oral. Rehab.*, 2002 29, 668–74.
- [18] Freilich MA, Meiers JC, Duncan JP, Goldberg AJ. *Fiberreinforced composites*. Hong Kong: Quintessence Publishing Co. Inc.; 2000; Pp. 6, 16, 30–47.
- [19] Ozcan M, Breuklander MH, Vallitu PK. The effect of box preparation on the strength of glass fiber-reinforced composite inlay-retained fixed partial dentures. *J. Prosthet Dent*, 2005 93, 337-45.
- [20] Altieri JV, Burstone CJ, Goldberg AJ, Patel AP. Longitudinal clinical evaluation of fiber-reinforced composite fixed partial dentures: a pilot study. *J. Prosthet Dent*, 1994 71,16-22.
- [21] Behr M, Rosentritt M, Lang R, Handel G. Flexural properties of fiber reinforced composite using a vacuum/pressure or a manual adaptation manufacturing process. *J. Dent.*, 2000 28, 509-14.
- [22] Ellakwa AE, Shortall AC, Shehata MK, Marquis PM. The influence of fibre placement and position on the efficiency of reinforcement of fibre reinforced composite bridgework. *J. Oral. Rehab.*, 2001 28, 785-791.
- [23] Lutz F, Göhring TN. Fiber-reinforced inlay fixed partial dentures: maximum preservation of dental hard tissue. *J. Esthet Dent.*, 2000 12, 164-71.
- [24] Sjolín R, Sundh A, Bergman M. The Decim system for the production of dental restorations. *Int. J. Computerized Dent*, 1999 2, 197–207.
- [25] Sundh A, Molin M, Sjogren G. Fracture resistance of yttrium oxide partially-stabilized zirconia all-ceramic bridges after veneering and mechanical fatigue testing. *Dent. Mater.*, 2005 21, 476–82.
- [26] Tinschert J, Natt G, Mautsch W, Augthun M, Spiekermann H. Fracture resistance of lithium disilicate, alumina-, and zirconia- based three-unit fixed partial dentures: a laboratory study. *Int. J. Prosthodont*, 2001 14, 231–238.
- [27] Hikada O., Iwasaki M, Saito M, Morimoto T. Influence of clenching intensity on bite force balance, occlusal contact area and average bite pressure. *J. Dent Res.*, 1999 78, 1336-1344.
- [28] Korber KH, Ludwig K: The maximum biting force as a critical factor for fixed partial dentures. *Dent Labor*, 1983 31, 55-60.

- [29] Nohrström TJ, Vallittu PK, Yli-Urpo A. The effect of placement and quantity of glass fibers on the fracture resistance of interim fixed partial dentures. *Int. J. Prosthodont*, 2000 13, 72-78.
- [30] Behr M, Rosentritt M, Leibrock A, Schneider-Feyrer S, Handel G. In-vitro study of fracture strength and marginal adaption of fibre-reinforced adhesive fixed partial inlay dentures. *J. Dent.*, 1999 27, 163-168.
- [31] Koutayas SO, Kern M, Ferraresso F, Strub JR. Influence of design and mode of loading on the fracture strength of all-ceramic resin-bonded fixed partial dentures: an in vitro study in a dual-axis chewing simulator. *J. Prosthet Dent*, 2000 83, 540-547.
- [32] Kelly JR, Tesk JA, Sorensen JA. Failure of all-ceramic fixed partial dentures in vitro and in vivo: analysis and modeling. *J. Dent Res.*, 1995 74, 1253-1258.
- [33] Chow TW, Chung RW, Chu FC, Newsome PR. Tooth preparations designed for posterior resin-bonded fixed partial dentures: a clinical report. *J. Prosthet Dent*, 2002 88, 561-564.
- [34] El-Mowafy OM, Rubo MH. Retention of a posterior resin-bonded fixed partial denture with a modified design: an in vitro study. *Int. J. Prosthodont*, 2000 13, 425-431.
- [35] El-Mowafy OM. Posterior resin-bonded fixed partial denture with a modified retentive design: a clinical report. *J. Prosthet Dent*, 1998 80, 9-11.
- [36] Castellani D, Baccetti T, Giovannoni A, Bernardini UD. Resistance to fracture of metal ceramic and all-ceramic crowns. *Int. J. Prosthodont*, 1994 7, 149-54.
- [37] Fairhurst CW, Lockwood PE, Ringle RD, Twiggs SW. Dynamic fatigue of feldspathic porcelain. *Dent. Mater.*, 1993 9, 269-73.
- [38] Ribeiro CCC, Baratieri LN, Perdiago J, Baratieri LN, Ritter AV. A clinical, radiographic, and scanning electro microscopic evaluation of adhesive restorations on carious dentin in primary teeth. *Quintessence Int.*, 1999 30, 591-9.
- [39] Geis-Gerstorfer J, Faßler P. Evaluation of the endurance limit for fatigue cycling of dental ceramics. *Dtsch Zahnärztl Z*, 1999 54, 692-4.
- [40] Sakaguchi RL, Douglas WH, DeLong R, Pintado MR. The wear of a posterior dental composite in an artificial mouth. A clinical correlation. *Dent. Mater.*, 1986 2, 235-40.
- [41] Kern M, Strub JR, Lu X-Y. Wear of composite resin veneering materials in a dual-axis chewing simulator. *J. Oral. Rehabil.*, 1999 26, 372-8.
- [42] Al-Dohan HM, Yaman P, Dennison JB, Razzoog ME, Lang BR. Shear strength of core-veneer interface in bi-layered ceramics. *J. Prosthet Dent*, 2004 91, 349-355.
- [43] Stappert CFJ, Guess PC, Chitmongkolsuk S, Gerds T, Strub JR. All-ceramic partial coverage restorations on natural molars. Masticatory fatigue loading and fracture resistance. *Am. J. Dent.*, 2007 20, 21-26.
- [44] Ernst CP, Cohen U, Stender E, Willershausen B. In vitro retentive strength of zirconium oxide ceramic crowns using different luting agents. *J. Prosthet Dent*, 2005 93, 551-558.
- [45] Guazzato M, Albakry M, Simon P, Ringer SP, Swain MV. Strength, fracture toughness and microstructure of a selection of all-ceramic materials. Part II. Zirconia-based dental ceramics. *Dent. Mater.*, 2004 20, 449-456.

-
- [46] Fleming GJP, Nolan L, Harris JJ. The in-vitro clinical failure of all-ceramic crowns and the connector area of fixed partial dentures: the influence of interfacial surface roughness. *J. Dent.*, 2005 33, 405-412.
 - [47] Rosentritt M, Plein T, Kolbeck C, Behr M, Handel G. In vitro fracture force and marginal adaptation of ceramic crowns fixed on natural and artificial teeth. *Int. J. Prosthodont*, 2000 13, 387-91.
 - [48] Lösche GM, Neuerburg CM, Roulet JF. Adhesive restoration of conservative class-II-cavities [in German]. *Dtsch Zahnarztl Z*, 1993 48, 26-30.
 - [49] Opdam NJ, Roeters JJ, Kuijs R, Burgersdijk RC. Necessity of bevels for box only Class II composite restorations. *J. Prosthet Dent*, 1998 80, 274-9.
 - [50] Feilzer AJ, De Gee AJ, Davidson CL. Setting stress in composite resin in relation to configuration of the restoration. *J. Dent Res.*, 1987 66, 1636-9.
 - [51] Darbyshire PA, Messer LB, Douglas WH. Microleakage in class II composite restorations bonded to dentin using thermal and load cycling. *J. Dent Res.*, 1988 67, 585-7.
 - [52] Mohsen C. Corrosion effect on the flexural strength and micro-hardness of ips e-max ceramics. *OJST*, 2011 1, 29-35.
 - [53] Harder S, Wolfart S, Echbach S, Kern M. Eight-year outcome of posterior inlay-retained all-ceramic fixed dental prostheses. *J. Dent.*, 2010 38, 875-881.

Chapter V

Removable Partial Dentures with Double Crowns - Biological and Prophylactic Value and Potential Complications

*Ivica Stančić and Aleksandra Jelenković Popovac**

University of Belgrade, Faculty of Dentistry,
Clinic for Prosthetic Dentistry, Belgrade, Serbia

Abstract

Correctly indicated and accurately made, partial denture with double crowns is a dental restoration of the highest biological, prophylactic and aesthetic values. However, insufficient knowledge of double crowns systems, as well as errors and omissions in the clinical and laboratory work invalidate this type of restoration and can have negative effects on the supporting tissues. Double crowns may come in form of telescope crowns and conical crowns.

Double crowns as connecting elements should provide: retention, stabilization, physiological transmission of occlusal loads through the abutment teeth as well as the prophylactic value, comfort for the patient, durability and aesthetics. In order to achieve these qualities, it is necessary, among other things: to make proper indication, to execute adequate tooth preparation (grinding), to make the choice of materials for the construction, to establish proper connection of the outer crown and the metal framework and to choose a suitable aesthetic material for veneering.

It is necessary to understand the mechanism of double crowns retention so that the optimal value of the retention force which should be 5-9 N per each matrix can be established. Insufficient retention force will reduce the comfort for the patient because would make the separation too easy, whereas an excessive force may negatively affect the periodontium of the abutment teeth.

* E-mail address: jelenkovic.aleksandra@gmail.com.

Retention force and durability of restorations with telescope and conical crowns depend also on the characteristics of the material they are made of. The most frequently used are gold and palladium-silver alloys. In recent years, titanium and the combination of ceramic and galvanized gold are used.

Prophylactic effect of double crowns is also provided by a proper planning that will enable the physiological transfer of the forces on the abutment teeth and the health of free gingiva. A special challenge is to plan the connection of outer crown with the metal framework. The best results showed optimized approximal extensions.

Double crown production demands experience of a clinician and dental technician in the performance of clinical and laboratory phases. A removal of sufficient tooth tissue during grinding in order to avoid getting oversized artificial teeth and a formation of the appropriate demarcation line which will provide sufficient stability of the system in the gingival area represent a particular challenge. Errors in laboratory work can be avoided with a good knowledge of the milling proceedings, a selection of a suitable material for the external crowns modeling and acquisition of precision castings.

Introduction

Regardless of the expansive use of dental implants, removable partial denture retains an important role in prosthetic rehabilitation of partial edentulous patients. Partial dentures may have clasps as connecting elements, and as such they are called conventional partial dentures. When connecting elements are attachment or double crowns, we talk about complex partial dentures. In terms of function and aesthetics, complex partial dentures meet the high demands of the profession, as they represent the combination of fixed and removable restoration, in which, using tooth reduction methods, abutment teeth are prepared to accept the elements of the removable partial denture [1]. While constructing these restorations, it is necessary to choose the retention elements, and plan stabilization and transfer of dental chewing pressure. In addition to offering comfort and good aesthetics, adequate retention and stabilization also affect the patient's feeling of confidence, providing him/her with adequate function.

The system of double crowns is one of the possible solutions for retention and stabilization of removable partial dentures. This system comprises of the male component of attachment or patrix, cemented to the abutment tooth, and the female component of attachment or matrix, which is the removable part of the restoration.

Double crowns in dentistry began to apply more than 100 years ago, and with many improvements over the decades, still continue to represent an effective system for rehabilitation of partial edentulousness. The cylindrical telescope system was first mentioned in the literature by James Dexter, (New York, 1883) and Water and Starr, (Philadelphia, 1886). First clinical application of the telescope system was described by Pess and Goosle (1924). However, at that time it was not accurately made and did not have adequate retention power. Soon the double crown systems began to improve (Häupl and Reichborn-Kjennerud, 1927) and a few years later Gründler and Böttger started to apply the milling technique in the preparation of the telescope system. The conus double crown system was first introduced by K.H.Körber, in 1957 in order to improve the telescope system deficiencies [2].

However, regardless of the highest biological and functional value of partial dentures with double crowns, they are rarely made today in everyday practice. The aim of this chapter is to explain the basic functional, prophylactic and aesthetic characteristics of double crowns

as well as to point out their shortcomings and the reasons for the lack of their practical application based on literature review and clinical experience.

Double Crowns - Characteristics

Definition, Indication, Types

Regardless of the double crown type, it always consists of two parts, one representing the fixed part of the dental restoration and the other that becomes a piece of the mobile part (Figures 1 and 2). Merged together, they form a complex which should represent the natural shape and size of the tooth. They separate only in one direction, the direction of the import and export of the restoration's mobile part [1].

An indication for a double crown is a partial edentulous dental arc, including a case of a maximum partial edentulous arch in cases where the remaining teeth are in favorable arrangement and with a preserved biological value. Periodontal status may reduce the possibility of making double crowns, bearing in mind that the long-term toothless rarely leaves periodontal tissue of remaining teeth intact. However, there are cases where the conservative and/or surgical periodontal procedures can prepare the abutment teeth for double crowns production.




Figure 1. Inner crowns on retention teeth.




Figure 2. Outer crowns with partial denture.

This is particularly the case with the elderly patients whose periodontal disease is usually chronic with little functional problems and a low level of the tooth mobility [3]. In such cases, a properly made denture with double crowns will have a positive effect on the periodontium of the abutment teeth and can prolong their lifetime.

Also, there are situations when the specific arrangement of the abutment teeth directs to the partial denture with double crowns as the only possible solution for the dental rehabilitation. These are cases in which there are few remaining teeth with a wide span of edentulous fields. It is desirable that the abutments are on either side of median line because in that way it gets multiangular surface reliance. The most favorable abutment teeth for double crowns are premolars and molars, and the optimal number of crowns is 2-4 [4]. A clinical examination alone can not give a reliable answer to the question whether the position and shape of remaining teeth ensure the proper acceptance of partial denture elements. Therefore, the data obtained by clinical examination must be completed with an appropriate analysis of study model in the articulator and parallelometer. The analysis in parallelometer is done from the vertical direction and includes: the assessment of the amount of tooth tissue which will be removed by grinding, the estimation of parallelism of axial surfaces of abutment, the assessment of undercuts of residual alveolar ridge etc [1, 5].

There are two basic types of double crowns categorization. The first and most common is based on the types of the assemblies that the crowns form, classifying them on telescope and conus crowns. Telescope crowns represent cylindrical construction with parallel surfaces and they are characterized by equal gingival and occlusal diameter. Conus crowns are double cone-shaped assemblies in which the cone angles lie in the range of 4 to 8 degrees. Conical angle represents an angle which forms vertical surface of the cone with a central vertical axis of the teeth [1, 2, 6]. Because of the different type of the construction telescope and conus crowns have different ways of obtaining the retention for the partial denture and partially different characteristics and indications. Telescope systems are primarily indicated in maximum partially edentulous arch or in one with a small number of remaining teeth. Conus crowns can be applied in cases of more abutment teeth. In telescope crowns, due to a specific retention mechanism and method of production, the retention force is unpredictable until the delivery point of the prosthesis. Conus crowns, on the other hand, retain partial denture with a defined retention force. They also come to the basic position easily during the process of taking in the denture. Due to the specific cone assembly, it can be indicated in cases of unparallel or inclined abutment teeth [7]. Although there are many advantages of conus crowns over telescope systems described, each assemblies have their places in everyday practice if their characteristics and potential complications in the production process are known.

The other type of double crowns categorization is based on their morphological and functional characteristics with shell-type crowns as the most commonly applied. They are characterized by the inner crown reaching the abutment tooth demarcation line and outer crown which gives the tooth natural size and shape. Although telescope and conus crowns are the most used of all shell-type crowns, they can also come in form of resilient, open, partial crowns and crowns on the pins [1].

Resilient double telescope crowns are the crowns with a vertical horizontal space between the inner and outer crown. The vertical gap is about 3 mm high and corresponds to an average value of resilience of residual alveolar ridge mucosa. The horizontal space between the crowns is 0.03 mm in width and corresponds to an average tooth intrusion. The crowns are

parallel in the gingival area and slightly convergent in the occlusal area. Their function is reduced to the stabilization and leading the denture to the basic position. The retention is achieved as in complete dentures which means that the chewing pressure transmission mode is mucosal. As interocclusal gap is used for mucosa resilience compensation, an indication for resilient double crowns is a high resilience value and a small number of remaining teeth with a compromised state of periodontal tissue. They are indicated exclusively in the lower jaw. Although this system is well designed there are some open issues such as a lack of knowledge of the process in the abutment teeth periodontium under pressure if a tooth is axially pressured. Also, the size of the vertical interocclusal gap relating to the mucous resilience is not clearly defined, therefore making the primary indication for resilient double crowns questionable.

Open double crowns represent a system in which the inner crown is modeled with occlusal surface and the outer crown is in the form of a ring. They are indicated in the lateral region, where there is not enough space for two occlusal surfaces. The advantages of this system are a good stabilization, dental occlusal force transmission, lesser removal of tooth structure and a better hygiene. The lack of retention power is its weakness so that the existence of other retention elements is required.

Partial double crowns represent a system in which the inner crown is shaped as grinded teeth for partial crown with grooves on the axial and occlusal surfaces. The outer crown complements the inner to the shape of a partial crown. They have lost their importance and as such are no longer applied. Today, though rarely, a variant of partial double crown is made in which the inner crown is a veneered or a metal-ceramic crown with a space for outer partial crown made with milling technique.

Double crowns on pin are system in which the inner crown is in the form of pin cemented in the root canal and cap. The cap covers the occlusal surface of the root up to the demarcation line and has a wedge that is placed on the occlusal surface of the cap. The external crown, in the shape of a metal-ceramic or a veneered crown is set over the cap and pin. Frequency of different types of double crowns in everyday dental practice is not the same, with some rarely present and some not at all. As it is already mentioned, usually the telescope and conus crowns are indicated, with the other assemblies to much lesser extent. This is a result of the unfamiliarity with the precise mechanism of achieving the retention of certain double crowns types and the cases in which they should be indicated. In addition to this, lack of knowledge of some of the production phases of certain assemblies by a dental technician is quite common. The biggest criticism of the previously described circuits is the uncertainty in the realization of their fundamental roles [5]. All double crowns are expected to achieve the optimal retention, stabilization and guidance of the denture into its main position. However, regardless of the fact that when making most common double crowns (telescope and conus systems) problems related to the achieving optimal retention force can be experienced (as it will be explained in subsequent parts of the chapter), with the other assemblies it is logical to expect even larger errors.

Materials for Double Crowns

As the bonding mechanism of fixed and mobile part of restoration with double crowns is based on the friction and elastic properties of the material they are made of, it is

understandable that the functioning of double crowns, to a large extent, depends on the choice of the material. Dental alloys for double crown production should have the following characteristics:

- biocompatibility (non-toxic to oral and other tissues, not causing an immune response and not potentially carcinogenic);
- electrochemical stability and corrosion resistance to the living environment;
- good physical and mechanical properties (high strength, adequate hardness, abrasion resistance);
- good and easy workability and
- good bonding properties with veneering material [8, 9].

Until now, for double crown production, mainly the noble and base alloys were used as well as their combinations. Gold alloys which are used for double crowns are of type III and IV, characterized by a high content of gold (up to 88%) and platinum (up to 11%), and with a mechanical properties of very high strength, hardness and compressive strength. From the group of noble alloys, Ag-Pd alloys are also used. Their advantage over gold alloy is a much lower costs. They have physical and mechanical properties similar to the type III gold alloy. Their disadvantages compared to gold alloys are that the castings are generally less homogeneous and therefore less accurate [8]. That leads to the insufficient persistence in the mouth which contributes to a greater accumulation of dental plaque and soft tissue staining.

From the group of base alloys for making double crowns, an attempt has been made with Co-Cr-Mo alloys. However, they have many defects that affect the quality of the final product:

- low density and lack of fluidity in the molten condition and as a consequence less accurate castings than the noble alloys;
- greater contraction of castings during cooling;
- difficult processing of the castings;
- doubled values of modulus of the elasticity comparing to type IV Au alloy, which significantly increases the stiffness and results with very low retention and
- hardness two times higher than type IV Au alloy [1, 9].

In the case of both telescope and cone systems, the retention force depends on the materials that they are made of. According to research [6], conus crowns made in the same way but with different material have different retention force. Crowns made of Co-Cr-Mo alloy have lower coefficient of friction than those of Ag-Pd alloys and especially of Au-Pt. It leads to conclusion that the retention force of the conus crowns made of non-precious alloys is smaller. Bearing in mind these fact, it is necessary to adjust the milling angle according to type of alloy in order to achieve optimal retention force of 5-10N. More specifically, the optimal milling angle for Co-Cr-Mo alloy should be 5 °, for Ag-Pd is 5.5 °, and for Au-Pt 6 ° (Table 1.). The differences in static friction coefficient of Co-Cr-Mo alloy and gold alloy are showed by Ohida et al. [10]. Based on their results, a smaller cone angle of the conus crowns made of Co-Cr-Mo alloys is suggested due to a lower coefficient of friction.

Table 1. Conus angle depending on type of alloy

Type of alloy	Coefficient of friction	Conus angle
Co/Cr/Mo	0,17-0,19	5
Pd/Ag	0,20-0,22	5,5
Au/Pt	0,22-0,24	6

During the production process, the inner and outer crowns can be made of the same or different materials, making the frictional couple therefore homogeneous or heterogeneous. For homogeneous frictional couple, mainly noble gold alloy Type III and Type IV are used. Gold in these systems acts as a solid lubricant [8]. Base alloys are less desirable for homogeneous frictional couples due to the significantly higher modulus of elasticity, extremely hard alloy components and less precision castings. In heterogeneous frictional couples there is a rule that the internal crown should be made of base alloys due to the higher modulus of elasticity and the outer crown of gold alloy because of a lower modulus of elasticity. Heterogeneous frictional couples work better and longer than the homologous pairs of base alloys and worse than homologous pairs of noble alloys.

Recently, new materials have been introduced to the double crown production in order to improve desirable characteristics. Significant researches have been made investigating the combination of ceramic and gold. Weigl et al. conducted in vitro studies complemented by clinical studies on the subject of double conus where the inner crown was made of ceramic and outer crown of gold alloy obtained by galvanizing process. The results indicate that these materials have possibility to become materials of the future for double crowns production [11, 12]. The advantages of ceramics-galvanized gold system lay primarily in a simple laboratory production. The inner crown is made of CAD/CAM system and the external one using a galvanization device. The result is high accuracy and very precious fitting of inner and outer crown. The retention force is independent of the pressure that is to say, to the intensity of masticatory forces. Also, this system shows high resistance to abrasion (wear) which is very important in the function of time. In addition to reduced wearing, there is no phenomenon of "cold welding" in areas of intense contact and gold does not form a galvanic element with ceramics with no corrosion as a consequence. It is known that ceramic has small plaque susceptibility, resulting in good health of periodontal tissue of abutment teeth. Also, with these materials some shortcomings in aesthetics of conventional systems can be overcome. In the cases of the withdrawal of marginal gingiva the metal edge wouldn't be exposed, but the tooth-colored ceramic.

Retention of Double Crowns

Double crowns as connecting elements should provide: retention, stabilization, physiological transmission of occlusal loads through the abutment teeth as well as the prophylactic value, comfort for the patient, durability and aesthetics. However, regardless of

the same roles they exercise, the manner in which the roles are exercised in telescope and conus crowns is basically different.

Perhaps the most important and most studied role of the double crown is retention. The reason for this is due to the fact that in achieving optimal retention lies a prophylactic effect of double crowns as well. Optimal retention of double crown is 5-10N for a single anchor [1, 2, 5, 13]. Any force that deviates from this optimum can have negative consequences for the entire structure. It is essential to understand the mechanism of retention with double crowns in order to establish the optimal value of the retention force.

The Mechanism of Achieving Telescope and Conus Crowns Retention

Mechanisms of retention of telescope and conical crowns differ. While the retention of the crown telescope is based on the dynamic-static friction, the retention of the conical crown is based on the static friction.

Telescope crowns have a form of a cylindrical structure characterized by gingival and occlusal equivalent circumference. The inner surfaces of outer crowns are adapted to the inner crown. The accuracy of this construction depends on the ratio of the radius or the diameter of the inner and outer part of the assembly. The difference between diameter of external and internal crown forms a gap. Depending on the size of the gap there are three types of the assembly: construction under tension, the transitional assembly and a loose assembly [5]. A construction under tension and a loose assembly, due to the elastic deformation of the outer part, and, for easy separation of cylinders, are not applicable. The size of the gap should be as near as possible to zero so that optimal frictional contact can be obtained; this is called the transitional assembly (Figure 3). Depending on the accuracy of the making, the friction effect appears when connecting or separating the outer from the inner crown. From the moment of the first touch of the inner surface of the outer crown and the outer surface of the inner crown until the final position, including the reverse process, this assembly is subjected to the laws of friction. In essence, the friction represents the resistance that occurs between two bonded atoms, molecules or bodies in relative motion under the influence of mechanical forces. The wear of the surfaces in contact is proportional to the force of pressure. In other words, if the gap is smaller, the force is greater and therefore the greater is the wear. This leads to the conclusion that parallel cylindrical constructions don't have the constant shape, the gap is increasing over time, and the transitional construction becomes loose. The presence of liquid between contact surfaces also has impact on the wear and geometrical shape of the construction [13]. The friction varies considerably according to whether it is a friction of wet, half-wet or dry surfaces. The size of the wear depends on the properties of the materials which are in contact (their hardness and roughness), the pressure and the number of cycles of the merge and separation. The total separation (and merge) force of telescopic denture with more than one telescope crown equals the sum of all forces that correspond to each particular crown.

The outer surface of every solid body is rough and wavy. Even the ideally smooth surfaces of quartz crystals show elevations of $0.01\mu\text{m}$. The smoothest metal surface has bumps from 0.05 to $0.1\mu\text{m}$ [5].




Figure 3. A. Gap is difference between diameters of secondary and primary crowns. B. Assemblies categories depending on gap size.

Inaccuracy, roughness and waviness of the outer surface of the inner crown and the inner surfaces of the outer crown cause their contact with each other only in certain small areas. The real contact area is several hundred times smaller than the geometric surface. The size of the actual contact area is always the sum of the individual contact surfaces. At points of intense contact cold welding phenomenon occurs as a result of heat generated during deformation. The result is a loss of substance and increase of surface roughness, which essentially allows the progression of the process of wear.

The local wear in a thin surface layer is particularly intense in the first phase of the friction process (the “initial wear” or “running-in phase”) and is a result of the fatigue of this layer due to repeated elastic deformation and their transition into the plastic deformation. The consequence of the “initial wear” is that the final geometrical shape of telescopic crowns is accomplished only after a certain time of use [15]. The process of casting can not generally affect these processes, but the occurrence of reversible frictional connections can be influenced by choosing alloys with low elastic modulus for the inner crown and for the outer crown alloy with effect of solid lubricant (e.g. gold).

A system of double crowns with conical shape was developed with an aim to eliminate the disadvantages of cylindrical construction and retain its good qualities. Conus crowns have the following positive characteristics:

- the ability to retain the denture with defined retention force;
- a function based on the static coefficient of friction and
- the ability to allow the separation of the construction without the sliding friction, making them not subjected to wear [1, 16].

The conical construction is different from the cylindrical in the fact that it has a milling angle (conical angle or the angle of longitudinal surface treatment) other than 0° . The inner crown is made at the angle of 6° , so it takes the form of cone with a greater gingival diameter than the occlusal [2]. The outer crown with its internal surfaces follows the inner crown, completing it morphologically, forming a natural tooth shape and also becoming an integral

part of the partial denture. When placing external crown over the internal, the contact surfaces get into reaction, producing the posture force. According to Kerber, this represents a fine friction which occurs only in the final contact phase [14]. The maximum idle mode friction signifies the force which before the start of the movement must be overcome only by the force of separation during denture removal. During the chewing function, the pressure force is generated which acts perpendicular to the axis of the tooth. As a reaction to the pressure force, a new force is produced, of the same course but in the opposite direction, called the normal or reaction force. The normal force in conus crowns is a component localized at the contact surfaces, and it, the same as the chewing force, causes the idle mode friction (Figure 4.). Normal force can also be described as a force of a closed, ring-like spring, making it the force with which the individual parts of the outer conus exert the pressure on the inner conus. Parts of the outer cone, due to elastic deformation, exert pressure on the inner part, as if it was composed of many rings. The occlusal force acts in such manner that each ring, moving on an incline plane, is being elongated to some extent. The consequence of that is that the inner crown's analogue ring is being compressed (Figure 5.). Thus the occlusal force, which originates from the construction juncture and the appropriate occlusal forces originated during mastication, converts into a deformation and is stored as the force of friction mode [2, 5, 6].

There is a dependency between the friction force and the size of cone angle. For the cone angle of 6° the separation force is 5-10N. For standard conus crowns, the angle of 6° is recommended. A smaller cone angle gives a greater friction force and therefore stronger retention, which means that it will be harder to remove the denture [5, 7]. For example, for the angle of 2° , eight times bigger force is needed than for the angle of 16° . Having this in mind, it is possible to plan and make some double conus crowns with greater retention force, if the biological conditions (health of periodontal tissue) permit that. On the other hand, crown made with a larger cone angle will contribute more to the stabilization and transfer of chewing pressure. It is known that conus construction works if the outer part has a slightly larger diameter with a vertical interocclusal gap inside the closed cone construction. In order to achieve defined retention force it is necessary to have a strong touch of the internal lateral surfaces. Only then sufficient idle mode friction can be generated.




Figure 4. Schematic review of conus constitution forces: PS–pressure force, N–normal force, SR–disunion force, Tv–displacement force, To–maximum rest friction force.




Figure 5. Scheme of secondary crown elastic ring effect on primary crown: PS—pressure force, P—rest friction force.

Also, it is necessary to make a minimum dimension of the inter-crown gap during the influence of maximum force with adequate deformation of the outer conus. That makes the conus construction elastic. According to the literature, this gap should be 0.1-0.3mm in the case of an unloaded system. During the activation of conus system, the inter-crown occlusal space is reduced for a few tens of microns depending on the nature of material, the force power and the geometrical parameters. At the termination of this action, the newly formed configuration can not be returned to its original state, although it is an elastic system, but remains under tension and the system works – it retains the denture [6, 14]. If the crowns are in direct occlusal contact, the occlusal forces are transmitted directly to the tooth and no part of the force can be converted into the elastic deformation of the outer crown.

Due to the different ways of achieving retention, the opinions of scientists and clinicians differ in terms of preferences of a particular system. Some senior scientists have favored the telescope system, primarily because of a better aesthetic effect, considering that its settlement on the gingival edge is much larger due to the zero-point milling angle [1, 17]. That way gingiva covers the veneered part of the crown in the visible region of the oral cavity better. The disadvantage of the conical system lies in the presence of an occlusal inter-crown space slightly larger than in the telescope parallel system, due to the structural characteristics of conus system. From the clinical aspect, this space is the weak point of the system because of the plaque accumulation and less resistance to lacunar corrosion [5]. However, the conus system has its advantages because its function is determined only by static rather than dynamic-static friction coefficient. Due to wear, which is of a slight intensity, the conus system has a longer clinical exploitation. In addition, there is a possibility of self-activation by the reason of its settlement as a result of contact surfaces wear [7]. Another benefit is the savings of the material compared to the telescope parallel system. This fact is based on the constructional characteristics of the conus construction which has a form much closer to the grinded tooth compared to telescope construction one.

Problems in the Retention of Double Crown

As it was previously mentioned, the optimal retention is one of the basic prerequisite for functionality, prophylactic effect and longevity of the partial denture with double crowns.

Insufficient retention force will reduce the comfort for the patient making the separation too easy, whereas an excessive force may negatively affect the periodontium of the abutment teeth. The double crown retention factors are the following:

- Number of abutment teeth. It directly affects the total retention force of the partial denture with double crowns. It should be always taken into account and the force of a single construction adjusted to the number of constructions. This means that if more than a few constructions are present, which is more often the case with conus crowns, it is not recommended to insist on an optimal retention force of 5-10N for each anchor. In the case of the conus crowns, the adjustment of the retention force shouldn't be a great problem, because it depends on the cone angle. In the case of telescope crowns, the retention force remains unpredictable until the delivery of the denture to the patient, which will be discussed below in this section. Therefore, the most common error in the retention of telescope dentures with more than three constructions is the excessive retention force which causes difficulties to the patient during denture separation.
- Retention force of each double crown construction. The force of each construction has a direct impact on the total retention force of the denture.
- Selection of alloys.
- Secondary factors. They also directly affect the retention force of the denture, although to a lesser extent. Secondary factors include physical and physiological factors. Physical factors are the surface tension, viscosity of the saliva, the adhesion and cohesion and the reduced air pressure. Physiological factors include the muscles of the cheeks, lips and tongue.

When planning the denture with double crowns, clinician can not be conducted only by physical and mechanical laws. Although the optimal retention force is proved to be 5-10N, it must be taken into account to which class the abutment tooth belongs as well as its biological value. Although the premolars and molars are the most desirable abutments for double crowns, the teeth of other classes can also serve very well. For example, if a central incisor's periodontium is preserved, it can certainly be included as a carrier of a double crown. However, due to its less periodontal surface, it can not receive the same dose of force as canines or molars. Therefore, a special attention should be paid that the retention force in incisors does not exceed the optimum level. Also, if the abutments have undergone periodontal treatment, it should be prevented that they suffer excessive force, even if it is only in the "running - in" phase.

Too weak or too strong retention force is the consequence of the discord between the inner and outer crown. This disharmony is a result of errors and variations (decrease or increase) of gingival and occlusal diameter.

The separation of telescope denture from the inner crowns depends also on the roughness of the contact surfaces, so that the gap has to be even wider if these values are higher [1, 13]. If this fact is overlooked, the telescope construction will succumb to rapid wear and smooth sliding motion would not be possible. As a standard procedure of the casting gives large differences in the dimensions of molded objects, an object with a smaller inner diameter of the secondary crown is obtained. In that way, a construction under tension in which both

crowns suffer from mechanical deformation is created. If the gap between the crowns is too wide, whether as a result of the incorrect production or due to wear, it will perform the opposite effect – the denture could not be retained and it will be moving during function. In order to achieve good characteristics of telescope or conus construction, all working processes and materials, from impression to polishing, must be well coordinated.

During the telescope crown production, the size of the gap depends on technical process and the degree of its reliability. On the casted inner part, which can be correctly made by milling, the outer part of the construction is modeled, so that the gap actually depends on the possibility of precise transformation of the wax model into the final form. In this procedure, several factors affect the accuracy: the properties of modeling wax and mass for investment, the modeling process, the method of casting, the characteristics of material for the outer part of the construction, the procedure for processing the cast etc, concluding with the always present human factor [5].

In a cylindrical construction, there is no insight into the value of the retention force during the process. In the final phase of denture delivery to the patient, with the use of different procedures for adjusting the force of separation, it can only be concluded whether the retention force is satisfactory or not, but its real value remains unknown [18, 19]. This can be a problem in the function, because when a clinician notice the eventual error, it is often difficult to eliminate. As a result of these problems, the telescope crowns retention measurement is often the subject of research.

The authors of this chapter carried out a similar research with the aim to measure individual and total retention force of the telescope dentures [18]. The research sample included 50 cylindrical structures fabricated at the Clinic for Prosthetic Dentistry, Faculty of Dentistry, Belgrade University. Retention force was measured on each structure. In addition, further measurements were carried out: the total retention force on 12 removable partial dentures with two cylindrical structures, on five dentures with three structures on three and four structures with dentures was measured. Double crowns were made of noble alloy type IV, in other words, a homogeneous frictional couple. Cylindrical structures were constructed for all classes of teeth, with the largest number, 55% for canine teeth. Measurements were carried out with the Bredent dynamometer (friktionsmeßgerät fmg 20; Bredent, Senden, Germany) [21] (Figure 6.).




Figure 6. The separation of patrix from matrix by pulling it vertically.

Results of measuring the individual force retention showed that the average values of retention force were from the value of 1 to 10.7 N. In terms of tooth classification, the force value was largest for the canine tooth, amounting to 6.5 N, and smallest for the molar tooth - 3 N. The results were derived by comparing the data collected from the individual matrix-patrix components: the optimal value was present in 50% of the cases, the force was too strong in 15% of the cases and was inadequate in 35% of the cases. The advantage of the method described in this report is that it provides the opportunity to measure the total retention force of a removable partial denture with a cylindrical design. Such a method allows observation of the change of the force depending on the varying number of components of the denture. Of the 20 removable partial dentures for which retention force was measured, only six (36%) had values within the limits of optimal values, whilst a reduced force was not recorded for any in this group. In the largest number of cases (70%) retention force was too strong. The lowest value for the total force was 5 N, and the largest 32.5 N.

Bearing in mind that the measurement was made under in vitro conditions, one should take into consideration the lack of a lubrication effect, which is present in the oral cavity. However, Bayer et al. showed that there was no significant difference between the measurements of retention force of double crowns with and without saliva lubrication [20]. As previously discussed in the mechanism of retention of the double crowns, crown definite geometric shape is reached after "running-in" phase. This is evidenced by this study because it has been noted that, while separating inner and outer telescopic crowns, the largest force value was observed during the initial separation.

The results of this study suggest that retention force is not suitable in 50% of cases of individual force and 70% of cases where the collective force is involved. This shows that a large percentage of the telescope constructions are actually dysfunctional, and that is why it is necessary to introduce some system for checking and measurement of retention force during the telescope denture production. Bayer et al. examined the average retention force of telescope crowns [20]. The retention force of 140 telescopic crowns was measured in the study. The retention force was measured with and without lubrication using saliva substitute. The minimum retention force was 12.08 N and the maximum retention force was 29.98 N. For all the lubricated specimens a median retention force value of 1.93 N was reached and an interquartile distance of 4.35 N was calculated.

If the retention force is not of optimal range a correction must be undertaken. In the case of insufficient force, it is necessary to line the cylindrical matrix component with a composite material. If the force is too strong, it may be necessary to repeat the fabrication of the matrix or framework to polish some contact segments of the matrix and matrix.

Insufficient retention force is a serious problem because it produces movement of the denture in the function [19, 22]. Relining with composite materials is a good solution, because the process is simple and gives good results. The composite that is used for this purpose must have a minimum of wear. The procedure can be performed directly on the patient or in the laboratory. It is also possible to improve retention when the denture is in operation for a while reducing the retention due to excessive wear. Retention improvement with composite materials lies in the fact that in this way, a new type of construction is made. The insertion of composite between the outer and inner crown creates a metal-resin construction which replaces the metal-metal complex [21]. The metal on resin fit offers the benefit of a considerably greater coefficient of friction than the one of a pure metal fit. Consequently, increased resistance to wear and an extended service life are obtained.

The Advantages of Partial Denture with Double Crowns – The Prophylactic Effect and Longevity

The advantages of partial denture with double crowns are numerous. The very fact that these systems have survived in practice more than 100 years speaks about their effectiveness in the prosthetic rehabilitation of partially edentulous patients. Properly indicated and accurately produced partial denture with double crowns represents a dental restoration of the highest biological and aesthetic values. Conversely, a poor indication and certain shortcomings in work, devalues this restoration to the point in which not only there is no more preventive and biological value, but it represents a threat to its supporting tissues.

The main advantages of double crowns are the prophylactic effect on the supporting tissues, longevity (persistence) and comfort, with the precondition that the protocol is respected in the planning of this kind of dental restoration. Prophylactic effect is reflected in:

- abutment teeth - extending their life;
- alveolar bone - reducing its absorption in the function of time and
- free gingiva - maintaining its health by appropriate design of the outer crown and metal framework relationship.

If the double crowns are properly planned and designed, they physiologically transfer the accepted masticatory forces to the abutment teeth. It will extend the lifetime of abutment teeth even if they were periodontally compromised to a certain degree. This of course includes the cases where the conservative and/or surgical periodontal treatment has been performed and abutment teeth are stabilized. Double crowns continue the functional stimulation and have a positive effect on the recovery and further stabilization of the periodontal tissues of the abutment teeth [1, 4]. Therefore, this also creates the prophylaxis effect together with positive effects on oral mucosa in the contact with denture and alveolar bone. The principles of tooth preservation are based on a rigid connection between the abutment teeth and the denture with mainly axial loading, which reduces any existing mobility [7]. The positive effect on abutment teeth is a result of a lack of traction and other forces and tooth movement during functioning, which occurs in the case when the connecting elements are clasps. The loads are directed to the longitudinal axis of the tooth, i.e. in the direction in which large numbers of periodontal fibers are engaged. In addition, if the retention force is optimal, it also positively affects the supporting tissues as a functional stimulus.

Prophylactic effect of partial denture with double crowns is also provided by appropriate design of restoration that will allow the health of gingiva of abutment teeth and ability to maintain adequate hygiene. This design is very important to establish in the area of the connection of the outer crowns with metal framework of partial denture. However, this link can have both positive and very negative effect on periodontal tissue of abutment if it is not properly planned. The connection of the outer crown with the metal framework can be achieved in several ways:

- direct connection by soldering;
- connection through the approximal extensions and
- casting in one piece.

The connection of external crown with the metal framework through the collars of framework prepared on the oral side of abutments has been a method of choice for a long time. The collars are connected with the metal framework by soldering. Although this relationship is reliable from the mechanical aspects, free gingiva is mechanically injured due to dead spaces and denture sinking. Also, when this connection is done with a larger number of conus crowns, it can lead to changes in axial dimensions of the external surfaces of crowns, which reflects on the cone effect realization and retention. Elastic deformation of certain parts of the outer crowns is disabled as a result of the increase of the crown wall because of the soldering with metal framework [1, 23]. In addition to endangering the retention effect, this can lead to falling aesthetic veneers.

The best results in terms of prophylaxis shows a connection through the approximal extensions on the outer crowns over which riders of the metal framework are placed. The riders are analogue part of the metal framework which fit the approximal extensions accurately. They can be connected to the metal framework by soldering with acrylic resin or composites. Besides the prophylactic effect, bonding through the approximal extensions also has a positive effect on the preservation of retention because the dimensions of outer crowns are unchanged [24]. Also, in this way, the cast is kept from further heat exposure in additional warming, especially if it has already been thermally and mechanically processed.

However, the standard approximal extensions so far have shown some disadvantages, especially if they have not been well planned. The consequences of poor planning are sharp corners that cause high concentration of tension as a result of which the connection may break. If acrylic resin is used for bonding this type of connection it often leads to separation of mobile and fixed compensation. Also, during the check-ups, the inflammation and thickening of the free gingiva of the abutment teeth are observed especially if there are dead angles where there is an accumulation of dental plaque, as this makes the oral hygiene maintenance difficult [24].

Based on years of clinical experience, the advice of authors of this chapter is to take the registration impression over the internal crowns and to plan and model the outer crowns on that plaster cast.

Although in this way the procedure is extended for one more impression and casting, this is negligible comparing to the positive effect. The positive effect is that the exact position of free gingiva around the inner crowns is visible allowing a very accurate modelation of outer crown with approximal extensions. The approximal extensions should be in the slight touch with the free gingiva, brought maximally close to the interdental papilla and creating enough space for the metal framework and artificial teeth. All this requires the use of articulators in this particular segment.

Today, many more modern forms of approximal extensions appear in the market, in order to overcome the shortcomings of conventional ones. In this aspect, the term optimization of the approximal extensions appears in literature [24]. It indicates an individual approach in creating the geometrical shape of the crowns with denture connection (Figure 7). (Stančić I, patent solution No. P-0758/04, 2004, decision of the Institute of the Intellectual Property Office granted patent on RS no. 50 205).




Figure 7. Optimized approximal extensions of outer crowns.

Comfort creates a sense of stability to the patient, because the system cannot be separated by the action of surrounding muscles, sticky foods or gravity. Horizontal forces that lead to high stress of the periodontal tissue can be successfully stabilized by double crowns. Complete coverage of the abutment teeth and the direct contact of the vertical surfaces in the main position of the denture is the primary basis of stabilization of these systems. Of course, in addition to overall stabilization, it is necessary to stabilize the occlusal forces. There is no uniform occlusal scheme; however, some general principles must be followed:

- provision of stable contacts in central occlusion;
- undisturbed conduction in the eccentric positions and
- optimum interocclusal space in the physiological rest position [3, 25].

Also, the comfort is provided by the fact that the system can easily be connected and separated when necessary, whereas the patient feels that the fixed and mobile part are connected into the one whole.

Beside the sense of comfort, stabilization provides the prophylactic effect on the alveolar bone. Reduced movement in function will reduce bone resorption in time. Also, the previously mentioned stabilized and balanced occlusion will have stimulatory effect on the alveolar bone and osteoblasts function. Of course, in order to maintain this function, it is necessary to carry out regular controls of occlusion and to perform reocclusion or relining if there is need.

It is known that properly made denture with double crowns are reconstructive solutions with longevity. In everyday practice, patients who have successfully used these dental restorations over 20 years can be seen. The main reasons for partial dentures ruin in general are problems with abutment teeth or technical failures. Abutment teeth may face large periodontal problems, fractures or caries after a few years [26]. The mobile part of the restoration may suffer stress that can lead to fractures. This is usually a result of errors in work. Fractures are complicated to repair, depending on localization; some parts, such as an occlusal rest or minor connector, are very complex and expensive to repair. When a partial denture with double crowns is well made, the abutments will not be exposed to excessive forces as previously discussed, but instead, these forces will have a stimulating effect on the periodontal tissue. This causes abutment teeth longevity. Also, if the homologous frictional couple of precious alloys (primarily gold alloy) is used, the longevity will be also provided by

good qualities of these alloys such as biocompatibility, less plaque receptivity and good physical and mechanical properties. All this provides a greater asset of double crowns compared to the other connecting elements of the partial denture.

Longevity of partial denture with double crowns was the subject of many researches. Wöstmann et al. examined in a retrospective longitudinal study the long-term survival of telescopic partial denture, taking into consideration possible influencing factors as well as treatment needs within the functional period [27]. The observation period was 5.3 ± 2.9 years. The examinations were comprised of caries diagnoses, tooth sensitivity and mobility, friction of the telescopic crowns in congruency between denture saddles and soft tissues, the state of the facings, the acrylic denture base and the metal framework. During the observation period 4.7 % of telescopic dentures required replacement, 3.8% of the abutment teeth were extracted. The survival probability after 5 years was calculated as 95.1% for the telescopic dentures and as 95.3% for the abutment teeth. The estimated 5-year survival rate for dentures with only one abutment was only 70.9% compared to 90.4% (two abutments), 95.0% (three abutments) and 97.9% (four abutments). Denture with no more than four abutments had to be replaced during the observation period. The most frequent technical failures were damages or loss of facings (26.9%) and loss of cementation (20.6%).

In a similar study of Behr et al. [28], the longevity of telescope and conus systems was investigated. The average observation time of parallel-sided retained dentures was 4.6 ± 1.6 years, and for conical dentures retained 5.2 ± 1.3 years. The first technical failures observed were recorded, such as fracture of the denture base metal, fracture of the saddle resin, fracture of the soldering, the loss of abutment teeth, loss of cementation, loss of artificial teeth, or loss of facings. During the observation time, 48.8% of the patients with conical crown-retained dentures and 34.2% of those with parallel-sided crown-retained dentures experienced complications. Loss of cementation occurred most frequently, in 26% of the parallel-sided double crowns and in 18.6% of the conical crowns. Loss of cementation and loss of artificial teeth were observed most often. The loss of cementation was the other most frequent technical failure in conical crowns but it occurred more than five years after insertion of the dentures.

These studies demonstrate durability of partial denture with double crowns showing that with these restorations there is relatively small percentage of complications in a long period. The most common complications are a loss of cementation and facing, which are relatively easy and inexpensive to solve [29].

Failures and Problems in Double Crowns Production

Regardless the numerous advantages of double crowns, they are now less and less frequently used in rehabilitation of partially edentulous patients, even when they are the only possible solution left. Possible reasons for this are complex production, lack of familiarity with the making process, inadequate laboratory, absence of dental technicians' training and previous bad experiences. For the partial denture with double crowns to exhibit all its positive features all clinical and laboratory phases must be properly planned and performed. Errors can occur during the whole process, from the bad indication to the failure in the cementation.

At whatever stage they occur, omissions and errors may devalue the entire dental restoration and adversely affect the supporting tissues.

In addition to issues of retention and prophylactic effect of double crowns, as described in previous sections, potential problems can also occur in different clinical and laboratory stages. Problems with the veneering can often put clinicians in dilemma as well.

Potential Problems and Mistakes in Clinical Phases of Double Crowns Production

The main characteristic of the clinical phases of fixed part of denture with double crowns is significantly higher reduction of the tooth structure compared to conventional metal-ceramic crowns. This happens because enough space for the cement, inner crown, outer crown and veneering material must be provided. As a consequence, the extension of the clinical crown and endodontic treatment of abutment teeth is often indicated. Not removing enough tooth structure can cause aesthetic problems, since the artificial teeth will be larger than natural teeth [1].

Grinding and impression of the fixed part of restoration take place as usual, similarly to the process in other complexes dentures, but it is very important to establish a clear demarcation line of sufficient width to ensure the stability of both crowns in the gingival area [4]. This requires a certain amount of skill and experience of clinician.

A tryout of the inner crown includes the examination of the gingival edge leaning on the demarcation and control of the occlusal space. As previously stated, it is best to take the registration impression over the internal crowns in this phase in order to make the outer crowns in accordance to the requirements of prophylaxis, or without negative effect on the free gingiva. The impression for the denture is taken over the double crowns and it is important that they have a secure position on the grinded abutments. Therefore, it is sometimes necessary to fix them with a paste for temporary cementation or elastomer. For the impression making, elastomers are used and priority is given to "one-phase" technique. The deformation during the impression removal after the polymerization of elastomers should be particularly taken into account. The following clinical stages take place in the usual way.

The phase of denture delivery to the patient is similar to one in any complex denture. Before the cementation of the inner crowns it is important to check the path of denture input, denture boundaries and occlusion. It is also necessary to isolate well the contact areas of double crowns, so that mobile part could be separated later without the use of excessive force. The separation is done after 24 hours at which time the first control is also performed. The first separation is usually more difficult because of the "running-in phase" [18].

Potential Problems and Errors in the Laboratory Stages of Double Crowns Production

Before the start of the inner crown modeling, it is necessary to determine the direction of denture input. For direction of input, the most important is the position of the abutment teeth in the frontal and sagittal planes. The procedure includes the choice of the middle position of

abutment teeth in both planes, and the orientation of the model first in the frontal and then in sagittal plane. This procedure not only determines the direction of import and export of the prosthesis, but also the direction of paralleling the inner crowns. If there are no undercuts in the area of residual alveolar ridge, then only centre lines of abutments are taken into account. The inner crowns are made by milling process.

The precision of laboratory work in double crown production mainly depends on the accuracy obtained by casting. The contraction of casting is inevitable, but it can be directed, including by the expansion of investment mass. For accurate casting the following is important: a precise measurement of fireproof mass and expansion fluid, thermal removal of tension/relax of the wax model, the possibility of expanding investment mass, the thickness of casting canals, the object's position in the refractory block, the thickness of the investment mass around the object, the object's position in relation to the direction of casting, the vacuum investment of wax object, respect of the preheating time and the use of precise equipment for casting [5].

Casting accuracy affects the size of the gap. Inaccurate cast causes excessive gap and, in addition to the impact on retention, has a negative impact on inter-jaws relation and increased accumulation of plaque.

Gap depends on:

- properties of modeling wax and mass for investment;
- modeling process;
- way of casting;
- materials for the outer crown;
- proceedings in the casting processing and
- human factors.

Materials for the external crown modeling are:

- wax;
- two-component plastic materials (a combination of plastic and wax materials) – self-polymerized acrylates and
- heliopolimerized materials (acrylates as fillers based on wax - LIWA program).

Self-polymerized acrylate has many advantages such as quick and easy creation and exceptional dimensional stability. Also, during the polymerization, a contraction is negligible because there is no thermal reaction, resin burns without residue and is eliminated in the T of 480°C. Molded surfaces are smooth and precise. Photopolimerized acrylate has advantages in terms of constant plasticity which allows safe and easy operation without time limit. It is easy for use because the material is one-component in the form, ready for use immediately. It has minimal contraction and therefore ensures the optimum edge closure. It is dimensionally accurate, because there is no deformation. Also, transparency allows control of layer depth.

The outer surfaces of the inner crowns and the inner surfaces of the outer crowns occupy particular attention in the laboratory work. Since the outer crown exhibits elastic deformation during the setting over the inner crown, it is important to take into account the dimensions of its walls. The most vulnerable area is the area of gingival margin.

Finishing process involves the inner crown polishing to a high shine. The impact of quality of finish is great. Well-polished surface has as a result lower static and dynamic coefficient of friction. The inner surface of the outer crown is not polished, but its treatment ends with sandblasting. The degree of processing significantly affects the size of the friction and the wear volume and thus the retention as well.

Aesthetics of Partial Denture with Double Crowns - Problems with the Veneering

The aesthetics of partial denture with double crowns generally shows two problems. The first is the inadequate removal of tooth substance by grinding, resulting in oversized crowns, as described in the section related to the clinical stages. The second problem is related to the veneers. Usually, the type of veneering material and dental technician incompetence can result in poor aesthetics, but an additional problem is the lack of retention of the veneer. As noted earlier, the fallout of the veneer is one of the most common complications of the use of partial denture with double crowns. Although it's not complicated for correction, for patients it can be an aesthetic and financial problem if it is frequent. As veneering materials the following can be used:

- ceramic materials;
- acrylate materials and
- composite materials.

Ceramic materials can be used as veneering material for double crowns, but due to the very high modulus of elasticity of ceramics, loss of veneers are often. For this reason and because of the prices of ceramics, it is rarely used for veneering. The acrylic materials can be used, but it must be taken into account their poor physical properties:

- softness;
- low resistance to abrasion;
- low impact resistance;
- low modulus of elasticity and
- insufficient strength.

The connection of acrylic with the metal of exterior crown is purely mechanical and often insufficient. Part of the metal that is covered with veneer is sandblasted with aluminum oxide particles of 110-250µm at a pressure of 3 to 3.5 atmosphere, depending on the type of alloy. In this way the contact surface between acrylate and crown is increased and the mechanical connection gets stronger.

Nowadays, composite materials are mostly used for veneering in double crowns. Their benefits are more favorable mechanical and aesthetic properties:

- abrasion resistance;
- high degree of hardness;

- high impact resistance;
- modulus of elasticity harmonized with module of the alloy;
- coefficient of thermal expansion (CTE) harmonized with the CTE of the alloy and
- excellent aesthetic results [30, 31].

A good aesthetic effect of a composite is reflected in its ability to achieve a high gloss with polishing, wide range color choice and the possibility for the crown's individualization. In order to obtain the best possible resistance of veneers, it should be provided with a minimum thickness of 1.5 mm. Also, it should be noted that alloys with more than 90% of gold, palladium or platinum produce small amount of oxide for binding, thus making the retention smaller. Every day, composite materials are more developed to achieve better and better aesthetic and mechanical characteristics. Composites with increased filler content are more stable and convenient for veneers in double crowns. They improve double crowns' properties in terms of high resistance to shock, ultra-low brittleness, greater resistance to abrasion and outstanding aesthetics.

The importance of composite veneers has been tested in numerous studies. It was proved that the wear of composite materials for veneering is lesser comparing to the conventional light-cured composites but higher than the gold alloy and ceramic. Also, they showed excellent color harmonization and long-term stability [30]. Matsumura et al. compared the hybrid composites with an increased amount of inorganic fillers and composites with micro-fillers, where the hybrid composites showed significant improvement of mechanical properties [31].

Conclusion

In order to achieve functionality, biological and prophylactic effect of denture with double crowns, dental team who participate in the process of its making is subjected to many challenges. Years of clinical experience suggest that successful production of double crown depends on the following:

- adequate indication;
- adequate preparation of the tooth for fix part of the restoration;
- selection of materials for telescopic or conical construction;
- quality of outer crown and denture framework connection;
- prophylactic value of restoration (to avoid overlap and mechanical irritation of the free gingiva);
- ability to maintain adequate oral hygiene;
- quality of the connection of the aesthetic restoration and alloys and
- retention force of the construction.

Qualification and experience of a clinician and dental technician allow detecting mistakes and correcting them on time, because later reparations are usually unsuccessful and costly. If

the clinical and laboratory procedures are obeyed, the result is the maximum functional and aesthetic rehabilitation of the patient and his satisfaction.

References

- [1] Stamenković D. *Stomatološka protetika - parcijalne proteze. (1st edition)*. Beograd: Interprint; 2006.
- [2] Körber KH. *Konuskronen-Teleskope, (1st edition)*. Heidelber: Hüthig, 1969.
- [3] Budtz-Jorgensen E.: *Prosthodontics for Elderly: Diagnosis and Treatment*. Chicago: Quintessence 1999.
- [4] Rehmann P, Weber A, Wöstmann B, Ferger P. Clinical evaluation of teeth fitted with telescope crowns for retaining a partial denture. *Deutsch Zahnärztl Z* 2007 62, 99-103.
- [5] Stamenković D, Stančić I. Retencija parcijalnih proteza dvostrukim krunama. *Stom Glas J* 2004 51, 30–37.
- [6] Jakšić M: *Konusni teleskop u mobilnoj stomatološkoj protetici, Vojna knjiga*, Beograd: Magenta Z.I., 1995.
- [7] Gungor MA, Artunc C, Sonugelen M. Parameters affecting retentive force of conus crowns. *J. Oral. Rehabil.*, 2004 31, 271–277.
- [8] Stamenković D, Obradović Đuričić K, Beloica D. *Stomatološki materijali. (1st edition)*. Beograd: Zavod za udžbenike, 2003.
- [9] McCabe JF, Walls WG. *Applied Dental Materials. (9th edition)*. Blackwell Munksgard; 2008.
- [10] Ohida M, Yoda K, Nomura N. Evaluation of the static frictional coefficients of Co-Cr and Au alloys for cone crown telescope denture retainer applications. *Dental Materials J.*, 2010 29, 706-712.
- [11] Weigl P, Hahn L, Lauer HC. Advanced biomaterials used for a new telescopic retainer for removable dentures: ceramic vs. electroplated gold copings: Part I. In vitro tribology effects. *J. Biomed. Materials Res.*, 2000 53, 320-336.
- [12] Weigl P, Lauer HC. Advanced biomaterials used for a new telescopic retainer for removable dentures: ceramic vs. electroplated gold copings: Part II. Clinical effects. *J. Biomed. Materials Res.*, 2000 53, 337-347.
- [13] Becker H: Rasterelektronemikroskopische Untersuchungen Reibflschen teleskopierender Kronen nach verschiedenen Arbeitsporozessen, *Quintessenz. Zahntech.* 1983 12, 1325-1333.
- [14] Körber K.H, Ludwig K: Maximale Kaukraft als Berechnugs Faktor zahntechnischer Konstruktionen. *Dent Labor* 1983 31, 55-61.
- [15] Wenz HJ, Lehmann KM. A telescopic crown concept for the restoration of the partially edentulous arch: the Marburg double crown system. *Int. J. Prosthodont*, 1998 11, 541–50.
- [16] Shiba A: *The Conical Double-Crown Telescopic Removable Periodontic Prosthesis*. St. Louis – Tokyo: Ishiyaku EuroAmerica, Inc., 1993.
- [17] Widbom T, Lofquist L, Widbom C, Soderfeldt B, Kronstrom M. Tooth-supported telescopic crown-retained dentures: an up to 9-year retrospective clinical follow-up study. *Int. J. Prosthodont*, 2004 17, 29–34.

-
- [18] Stančić I, Jelenković A. Retention of telescopic denture in elderly patients with maximum partially edentulous arch. *Gerodontology*, 2008 25, 162-167.
 - [19] Shimakura M, Nagata T, Takeuchi M et al. Retentive force of pure titanium conus telescope crowns fabricated using CAD/CAM system. *Dent. Materials J.*, 2008 27, 211-215
 - [20] Bayer S, Stark H, Mues S, Keilig L, Schrader A, Enkling N: Retention force measurement of telescopic crowns. *Clin Oral Investig*, 2010 14, 607-11.
 - [21] Bredent: *Products for the Dental-Technical Laboratory*. Senden (Germany): Bredent, 2003-2004.
 - [22] Ohkawa S, Okane H, Nagasawa T, Tsuru H. Changes in retention of various telescope crown assemblies over long-term use. *J. Prosthet Dent*, 1990 64, 153.
 - [23] Akagawa Y, Seo T, Ohkawa S, Tsuru H. A new telescopic crown system using a soldered horizontal pin for removable partial dentures. *J. Prosthet Dent*, 1993 69, 228.
 - [24] Stančić I. *Teleskop proteze - veza krune i skeleta*. Beograd: Zadužbina Andrejević, Biblioteka disseratio, 2005.
 - [25] Versteegh PA, Polder BJ. Prosthetic dilemmas. Prosthetic decisions and planning for partially edentulous patients. *J. Ned Tijdschr Tandheelkd*, 2008 115, 580-7.
 - [26] Budtz-Jorgensen E, Isidor F. A 5-year longitudinal study of cantilevered fixed partial dentures compared with removable partial dentures in a geriatric population. *J. Prosthet Dent*, 1990 64, 42-7.
 - [27] Wöstmann B, Balkenhol M, Weber A, Ferger P, Rehmann P. Long-term analysis of telescopic crown retained removable partial dentures: survival and need for maintenance. *J. Dent.*, 2007 35, 939-945.
 - [28] Behr M, Hofmann E, Rosentritt M, Lang R, Handel G. Technical failure rates of double crown-retained removable partial dentures. *Clin. Oral Investig.*, 2000 4, 87-90.
 - [29] Hofmann E, Behr M, Handel G. Frequency and costs of technical failures of clasp- and double crown-retained removable partial dentures. *Clin. Oral. Investig.*, 2002 6, 104-8.
 - [30] Mimura S, Toyofuku T, Ryumon K, Yamamura O, Matsui H, Fujii T. The study on adhesive strength of mesh veneer plate with adhesive resin. *J. Gifu Dent Society*, 1988 15, 420.
 - [31] Matsumura H, Tanoue N, Atsuta M. Adhesive bonding of stainless steels and their component metals. *J. Oral. Rehabil.*, 2000 27, 1054-1059.
 - [32] Minagi S, Natsuaki N, Nishigawa G, Sato T. New telescopic crown design for removable partial dentures. *J. Prosthet Dent*, 1999 81, 684.

Chapter VI

Systemic Changes of Interest for Rehabilitation with Complete Dentures: Manufacture and Maintenance

***Leandro Dorigan de Macedo^{1,2,3}, Soraia Marangoni¹,
Maria do Carmo Favarin de Macedo³, Hilton Marcos Alves Ricz⁴
and Luciana Assirati Casemiro^{*,1}***

¹University of Franca, Dentistry School, Franca, SP, Brazil

²University of São Paulo at Ribeirão Preto, Medical School, Brazil
Hospital das Clínicas, Dentistry and Stomatology Service

³Fundação Hemocentro at Ribeirão Preto, Ribeirão Preto, SP, Brazil

⁴University of São Paulo at Ribeirão Preto, Medical School,
Hospital das Clínicas, Head and Neck Service, Brazil

Abstract

The aging population has profoundly transformed the profile of patients seeking oral rehabilitation. Care delivery to these patients, who present with multiple comorbidities and are usually being treated with various drugs, is becoming increasingly more frequent. Systemic changes prompted by the use of many medications may impose limitations to clinical practice with regard to indication, planning, materials, preparation techniques, and maintenance of prosthetic rehabilitation. Unfortunately, the dental literature is lacking in scientific papers reporting oral manifestations of systemic diseases, as well as in works describing the type of assistance that is necessary to treat such patients. The reported findings are limited to the description of techniques, which are sometimes adapted, for complete denture treatment in patients with neurological diseases, such as Alzheimer's and Parkinson's syndrome, and burning mouth in cases of allergy to dental materials. Several systemic conditions and different treatments should be taken into

* E-mail address: lucianacasemiro@hotmail.com.

account when it comes to the correct time for prosthetic rehabilitation completion, the materials and technique to be used, and the maintenance procedures to be employed. It is known that in the case of patients with severe immunosuppression, whose origin can be constitutional (senile), pathological (hematological, immune, or endocrine), or prompted by medication (use of immunosuppressants, corticosteroids, chemotherapy, among others), there is higher incidence of oral infections by fungi, especially opportunistic infections such as aspergillosis and mucormycosis, whose incidence may increase due to badly fitted and poorly cleaned dentures. Another group of autoimmune diseases, the so-called pemphigus vulgaris, may have repercussions in the oral mucosa. This disease predisposes the oral cavity to serious injury by lack of adherence of epithelial tissue and by infection due to the constant utilization of immunosuppressants, making the use of dentures very difficult. In addition, patients with renal failure and severe liver disease may present changes in both oral mucosa and salivary glands, which demands manufacture and use of dentures. Treatments such as radiotherapy and the chronic use of bisphosphonates imply in severe changes in both the bone and soft tissues. The common denominator of all these diseases or therapeutic protocols is that they have temporary or permanent contraindications for rehabilitation with conventional dentures or implant overdentures. Thus, on the basis of literature and a 10-year clinical group experience in a dental hospital service, this book chapter aims to describe major diseases and drug therapies that directly or indirectly reflect on the oral cavity, including the necessary adjustments in materials, manufacturing techniques, and maintenance of complete dentures.

Introduction

The development of technologies and methods in health care has become increasingly common the early diagnosis of diseases before unknown, besides of allowing a better monitoring of responses to treatment and disease evolution, often at the molecular level. Allied to these factors, the improvement of treatments using surgical and microsurgical techniques and more effective drugs are responsible for the increase in life expectancy worldwide. Faced with this scenario, it is increasingly common the dentist to meet, at the dentist office, patients with chronic diseases and comorbidities using different drugs [1-5].

Several pathologies and treatments may predispose to important changes in the stomatognathic system involving the oral mucosa, salivary glands, nerves, muscles and bone structure. These changes place the dental surgeon as a major agent in the early diagnosis of some diseases (such as tumors, diabetes, AIDS, among others), in the control of the effectiveness of treatment (pemphigus vulgaris and Behçet's disease, for example), and in the prevention and treatment of oral complications resulting from pathology or drugs, such as osteonecrosis by bisphosphonates and bleeding due to the use of anticoagulants.

Despite a worldwide trend in the reduction of edentulism, some of these chronic diseases and their treatments cause directly or indirectly the loss of dental elements in response to oral complications. These patients often have no systemic or psychological conditions for the rehabilitation with implants or have temporary or permanent contraindication for this type of rehabilitation due to the risks of failure and pathological complications in the bone tissue, not to mention the financial limitation that is imperative in deciding which type of prosthesis should to be used [6-8].

The rehabilitation with conventional complete dentures is often the best option for treatment in these patients; and local or systemic changes brought by these conditions require special care in the manufacture, installation and maintenance of prosthetic devices. Thus, this chapter aims to present the systemic changes and the most common side effects that interfere directly or indirectly in the indication for complete dentures, in its manufacture and maintenance. Unfortunately the literature is scarce in terms of dental management of these patients; the studies found are restricted to literature reviews, and to presentations of clinical case. Techniques and therapeutics showed here are based on the findings from the literature and 10 years of experience in hospital dentistry.

Immunosuppressive Changes

The microflora found in the oral cavity is known as one of the richest and most diverse in the human body. Although bacteria are the most abundant microorganisms in its composition, fungi are the most important causative agents of complications in immunosuppressed patients that make use of prosthetic devices [9-13]. Such microorganisms are organized in the form of biofilms; and, in healthy individuals, they rarely cause infectious complications because local barriers and immune system provide a protection that prevent the development of these infectious complications in the oral mucosa [14, 15].

Some fungi are common inhabitants of the oral cavity. The species most commonly found belong to the genus *Candida* and usually they are present in low concentration and do not cause disease [10, 16]. The identification and quantification of these pathogens have been routinely performed by conventional culture, but methods such as Polymerase Chain Reaction (PCR), REAL-time PCR, and Checkerboard CNA- DNA hybridization have made the recognition and quantification of microorganisms quickly and efficiently [9, 17, 18]. A literature review accomplished by Odds in 1988 found a prevalence of fungal colonization ranging from 20% to 71% in healthy patients [19], as confirmed by recent studies [20, 21]. The species most commonly found in descending order are: *C. albicans*, *C. glabrata*, *C. dubliniensis*, *C. tropicalis*, *C. Krusei*, *C. parapsilosis*, *C. famara*, and *C. guilliermondii*.

Changes that lead to local or systemic immunosuppression are associated with significant increase in oral fungal infections concomitant with the increase of these yeasts both in saliva and in tissues. Figure 1 presents a case of Aspergillosis infection in an immunosuppressed patient.

Several studies have shown a high incidence of *Candida* species infections in immunosuppressed patients, and the complications observed in these cases are closely related to the degree of immunosuppression, the integrity of the oral mucosa, and the amount of fungi present in the infected sites. The superficial candidiasis is the most frequent disease and, if untreated, it can raise to systemic infections mainly in severely immunosuppressed individuals, putting at risk the survival of these patients [22-27].

The emergence of fungi resistant to drugs has been reported in the literature [9, 28-31], and the factors responsible for this occurrence are: the recurrent use of low-doses of antifungal agents, changes in protein metabolism, and other genetic events. The biofilm formation has been appointed as an important factor in the increased emergence of resistance to azole drugs [11, 32-35]. In this context, the complete denture has an important role as

carrier of infectious agents because they act as reservoir of fungi in the form of biofilm [36, 37].




Figure 1. Aspergillosis infection in an immunosuppressed patient after allogenic bone marrow transplantation for leukemia.

The immunosuppressive conditions most commonly associated with the increase of oral candidiasis are: AIDS, hematological diseases, diabetes, use of immunosuppressants after transplantation, cancer treatments (surgery, radiotherapy, and chemotherapy), and chronic use of corticosteroids and of immunosuppressants to control autoimmune diseases, among others. The immunosuppression caused by disease will be discussed in this topic, and the drug immunosuppression will be discussed in each specific diseases.

Acquired Immune Deficiency Syndrome (AIDS)

The recurrent oral candidiasis without local causative factors and difficult to control are still considered one of the first signs of the AIDS. The dentist must be aware of this finding and request for serology and evaluation by a specialist in infectious diseases.

Although the incidence of oral candidiasis were significantly reduced after the advent of antiretroviral therapy (ART), its occurrence in HIV/AIDS patients is still frequent and well above that found in the general population.

In 2008, Earlandsen and collaborators evaluated 122 patients on ART and found colonization by species of the genus *Candida* in 81.1% of them. The authors also observed that 33.3% of the individuals examined had symptomatic infection, while 25.3% of the colonized patients had yeasts resistant to fluconazole. The resistance to azole was frequent among HIV/AIDS patients which made it difficult to control the disease. With the use of ART and reducing the use of azoles, the frequency of resistance of germs to drugs has reduced, but it remains in patients with advanced AIDS, in whom other species are often found (*C. glabrata*; *C. krusei*; *C. tropicalis*; *C. parapsilosis*; and *C. Dublinsers*) [38].

The risk of developing oropharyngeal candidiasis is closely related to the absolute count of CD4⁺T-lymphocyte <200cell/ μ L. However, current findings suggest that the occurrence and severity of candidiasis are most associated to the viral load; and values of count greater than 10000 copies/mL seems to expose the patient to increased risk of developing such infection [13, 39-43].



Figure 2. Terminal AIDS patient with oral and pulmonary infection (clinical aspect and tomography) caused by *Candida albicans*.

Diseases	Patient stable CD4+> 500 cells/ μ L and or VL<1000 copies/mL	Patient stable CD4+ from 200 to 500 cells/ μ L and or VL <5000 copies/mL	Patient stable CD4+ <200 cells/ μ L and or VL> 5000 copies/mL
Oral candidiasis	- Professional cleaning of CD and Immersion in 0.2% Clrx solution for 5minutes -Daily immersion in 0.2% Clrx for 2 minutes/14 days -Nystatin 100,000 IU 4 times/day for 14 days or oral gel of miconazole 4 times/day	- Professional cleaning of CD and Immersion in 0.2% Clrx solution for 5minutes -Daily immersion in 0.2% Clrx for 2 minutes/ 14 days -Fluconazole 100mg/day for 7-14 days	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes -Daily immersion in 0.2% Clrx for 2 minutes/ 14 days - Use of the CD only during the meals until the resolution of the disease -Fluconazole 100mg/day for 7-14 days
Oropharyngeal Candidiasis	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Fluconazole 100mg/day for 7-14 days	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Fluconazole 100mg/day for 7-14 days	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Fluconazole 100mg/day for 7-14 days
Failure after Fluconazole	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Itraconazole 200mg/day for 14-21 days	- Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Itraconazole 200mg/day for 14-21 days	-Cleaning of the CD and Immersion in 0.2% Clrx solution for 5minutes - Use of the CD only during the meals until the resolution of the disease -Itraconazole 200mg/day for 14-21 days
Refractoriness	- Change of the prosthesis - Hygiene Care -Use of Echinocandins	- Change of the prosthesis - Hygiene Care -Use of Echinocandins	- Change of the prosthesis - Hygiene Care -Use of Echinocandins

Figure 3. Treatment protocol of oral and oropharyngeal candidiasis in HIV/AIDS patients. (VL: viral loud; CD: complete denture; Clrx: chlorhexidine).

The dentist responsible for the manufacture and maintenance of complete dentures in these patients should be upgraded about the medical situation, adherence to treatment, use of medications such as antibiotics (which also predisposes to fungal infections), and results of laboratory exams (mainly viral load and count of CD4). Complete dentures can be made in a conventional way since the patient does not present lesion or infection in the oral mucosa. Figure 2 presents a clinical case of a patient with oral and pulmonary infections by yeast, and with terminal AIDS.

The adoption of an appropriate denture maintenance plan is very important for the success of the treatment. Auxiliary hygiene methods such as cleaning solutions and specific toothpastes should be used only for cases of poor hygiene denture or previous candidiasis [44-46]. If diagnosed candidiasis, the use of denture should be reduced as much as possible, until solving the disease. The use of rebasing material to reduce the trauma of oral mucosa can also be considered important, but it must be remembered that in the presence of candidiasis, it can serve as a reservoir of yeasts. Prophylaxis is not routinely used, but daily doses of fluconazole (100mg) can be used in cases of frequent or severe re-infection, until it reaches a count of CD4+ higher than 200 cells/ μ L [13, 47].

Figure 3 shows the protocol used for treatment of candidiasis in the Dentistry Service of the *Hospital das Clínicas* of the University of São Paulo at Ribeirão Preto Medical School. Other protocols can be found in the literature.

Every therapeutic choice should be discussed and decided together with the specialist in infectious disease who is responsible for the case. Cases with signs of dysphasia and doubts related to the esophageal involvement should be treated by the medical team. In all cases the denture relines is recommended when the denture does not fit, and this procedure should be done after a professional rigorous cleaning and chemical disinfection of the prosthetic device.

Hematologic Diseases

There are a variety of hematologic diseases that course with immunosuppression due to the lymphocyte commitment and/or severe neutropenia. Treatments of these pathologies often involve the use of chemotherapy agents highly myelotoxic and stomatotoxic, which can result in oral mucositis associated with severe neutropenia, thrombocytopenia and anemia [48,49]. Some of these diseases have oral manifestations as a primary signal (Figures 4 and 5).



Figure 4. Gingival lesion caused by Lymphoma – it was the unic signal of the disease in this patient.




Figure 5. Atrophy of tongue mucosa secondary to anemia.

There are no reports comparing the incidence of oral candidiasis in these patients and HIV/AIDS carriers. However, the presence of severe neutropenia exposes these patients to serious systemic infections associated with risk of death in a more critical way than in patients with AIDS [50-52]. Therefore, treatment should be more aggressive, and the use of removable dentures is entirely dispensed in the most severe cases.

The diagnosis of candidiasis is clinical and confirmed by direct identification of hyphae by exfoliative cytology or biopsy. Culture and assessment by PCR may facilitate the differential diagnosis among the types of *Candida*, but they are not routinely performed due to the large amount of surface contaminant found in the sample. Treatment depends on the extent of lesion, symptomatology and especially the degree of neutropenia (Figure 6).

The forms of superficial infection most commonly found in patients with onco-hematological diseases are pseudomembranous, which is characterized by white plaques that can be removed by scraping, and atrophic or erythematous, in which there is mucosal atrophy accompanied by local erythema. Disease progression is most often asymptomatic.




Figure 6. Flow chart showing the treatment for superficial oral candidiasis in patients with hematologic disease (Ne: Neutrophils, represented in $\times 10^3$ cel./ μ l; IU: International Units).



Figure 7. Bleeding in a patient with thrombocytopenia after chemotherapy to treat leukemia.

At our service, it has been observed an increased incidence of oral candidiasis caused by *C. glabrata*. Infections caused by these microorganisms generally have specific clinical manifestations, characterized by the presence of scraped membrane associated with necrotic tissue, local bleeding and pain. Treatment of choice is voriconazole and another option is to use Amphotericin B, since this yeast has been shown to be resistant to fluconazole [53].

Thrombocytopenia rarely brings bleeding complications in patients with complete denture, except in cases of trauma and laceration due to prosthesis associated with mucositis (Figure 7). In cases of platelet (PLQ) count greater than $50,000 \times 10^3 \text{ cells}/\mu\text{L}$, the bleeding is usually controlled with local methods, such as suturing and compression. It is indicated the transfusion and use of antifibrinolytic agents for refractory bleeding or in cases of $\text{PLQ} < 40,000 \times 10^3 \text{ cells}/\mu\text{L}$. This decision must be taken together with the hematologist.

Patients with acute hematologic diseases such as leukemia and lymphomas hardly seek a dentist for the manufacture or maintenance of prostheses, which is usually performed in hospitals. But those with long-term diseases such as myelodysplastic syndromes, aplastic anemia, Fanconi anemia, multiple myeloma, among others, may be submitted to the construction of conventional complete dentures, being essential to take care of using flexible molding materials such as silicones to reduce the risk of tissue damage. In addition, a full care should be taken with the disinfection of models, base plate, and other devices produced by the prosthetic.

Special attention should be given to patients with multiple myeloma because of the possibility of developing plasmacytomas, which are locally destructive bone lesions that can affect the mandible and jaw, causing risk of fracture. This situation requires change in hematologic treatment, with indication of radiotherapy in some circumstances. Thus, the panoramic X-ray should be taken in the beginning of treatment and whenever there is need. Moreover, it is common for these patients to take bisphosphonates which increase the risk of bone necrosis and contraindicate the use of the implant.

As for HIV/AIDS patients, the maintenance of prosthesis must be rigorous and the dentist must know in detail the patient's clinical condition. The dentist should be aware about laboratory tests (such as complete blood count), medication and chemotherapy treatment to predict the risk of complications in each moment. Patients who have neutrophil count less than $500 \times 10^3 \text{ cells}/\mu\text{L}$ should be treated by professionals.

Diabetes

The patient with diabetes, a metabolic syndrome with autoimmune feature, may have local or systemic immunosuppression and difficulties in the healing of oral lesions. These findings have been related to the progression of the disease and its control. Patients who are users of complete denture and with bad glycemic control may have increased risk for the development of oral candidiasis, especially in cases of failure in the adaptation of the prosthetic device [54].

Some studies show that *C. albicans* isolated from diabetic patients showed higher adherence to fibronectin and epithelial cells [55, 56]. This finding associated with a greater amount of hyphae in the oral cavity related to the higher concentration of glucose in saliva increases the risk of candidiasis in these patients [57-59].

The clinical framework, with rare exceptions, tends to be localized, less invasive, and should be treated with topical agents devoid of sugar such as miconazole oral gel (4 times/day for 14 days).

The use of nystatin should be avoided due to the large amount of sugar in its composition. The cleaning of the prosthesis by a qualified professional and the relining should be performed and, after the resolution of the infection, new prosthesis needs to be made if the old one is inadequate. In cases of maintenance of infection, after the seventh day of treatment with topical antifungals, the use of fluconazole is indicated (100 mg/day from 7 to 14 days).

It is important to highlight that the prosthesis may be the agent responsible for maintaining the fungal disease even after the rebasing. If the infection persists, the use of dentures should be discontinued until the resolution of the infection.

Immunosuppressive Drugs and Treatments

Several autoimmune diseases are treated with chronic corticosteroid regimens associated with or without immunosuppressive drugs. In all cases the care of the maintenance of prostheses and treatment of oral candidiasis should be performed similarly to that described for the diabetic patient.

Special attention should be given to the treatments that may cause neutropenia, such as chemotherapy treatments. Each case must be evaluated separately and the conduct should be taken depending on the degree of neutropenia and extent of the lesion, as shown in Figure 6.

Diseases with Direct Impact in the Oral Cavity

Several pathologies have direct repercussion in the oral cavity, and the first clinical symptoms often affect the oral mucosa. In the following, the diseases that most require special care or changes in treatment regarding the rehabilitation with complete dentures will be shown.

Phemphigus Vulgaris and Cicatricial Pemphigoid (Mucous Membrane Pemphigoid)

Despite having different histological features, clinical evolution and extent of involvement of tissues, the phemphigus vulgaris and cicatricial pemphigoid can cause serious lesions in the oral mucosa that often restrict the use of removable dentures. They are autoimmune blistering disorders characterized by the production of antibodies directed at different regions of the epithelium [57].

The mucous membrane pemphigoid or cicatricial pemphigoid is characterized by subepithelial bullous lesions with a predilection for the elderly, mean age 62 to 66 years [58-60]. The most commonly affected areas are the oral and ocular mucosa; but it may also presents lesions in genito-urinary, gastroesophageal and respiratory tracts. Severe complications such as upper airway obstruction, loss of vision, urinary, and sexual dysfunction can occur [61].

The dental surgeon is an important agent in the diagnosis and in the support for treatment of this pathology since the oral mucosa is involved in almost 100% of cases, being often the only affected area [62]. Cutaneous involvement can occur, though more rare. The oral cavity presents recurrent superficial exfoliative lesions, especially in areas of trauma, and the diagnosis can be confirmed by biopsy that presents subepithelial lesion associated with linear deposition of IgA, IgG and/or C3 along the epidermal basement membrane zone [61].

On the other hand, pemphigus vulgaris is characterized by intraepithelial blistering and affects the skin and oral mucosa. The initial lesions are often attenuated and attack the oral cavity, followed by an aggressive skin picture. The disease is precipitated by IgG autoantibodies binding to desmoglein 3. As the lesions are more superficial, the oral mucosa presents desquamative process with areas of scrub and, luckily, bullous lesions may be observed before its outbreak. From seventy to ninety percent of patients develop erosions in the oral cavity [57] (Figures 8, 9, and 10).

The rehabilitation of edentulous patients who have pemphigoid secondary oral lesions is a challenge in the sense that the conventional prostheses may worsen the case due to the trauma on the tissue. However, there are no studies reporting the peri-implant health in patients with these pathologies, since the cleaning of protocol-type prosthesis would be an important traumatic factor to the mucosa.



Figure 8. Tongue lesions in a patient with mucous membrane pemphigoid.



Figure 9. Pemphigus vulgaris: initial lesion.




Figure 10. Oral advanced involvement of pemphigus vulgaris.

In our clinic there are two patients who were using this type of prosthesis when the diagnosis of cicatricial pemphigoid was made. They had to stop the use of the prosthesis due to the recurrent bacterial infections and peri-implant lesions.

The attempt to manufacture conventional complete denture is suitable in those cases well controlled; in these situations, care must be taken in order to avoid trauma to the tissue during the manufacture of prostheses. The molding should be atraumatic and with the use of elastic materials, such as silicones (the materials of choice). The maximum adaptation should be obtained. The use of resilient reline seems to reduce lesions on the mucosa, and the occlusal adjustment should be carefully conducted. All possible interferences and overload to tissues should be adjusted. At our service, during the first 14 days, patients are reviewed daily for adjustments. In cases in which the patients do not use topical corticosteroids, we have initiated the use of a combination of hydrocortisone, neomycin, troxerrutin, and ascorbic acid in oral ointment administration, three times a day, on the surface of contact, 14 days before the installation of the prosthesis and with apparent success in reducing complications.

In cases of recurrent injuries which are difficult to control we have proposed the use of removable implant-prosthesis, in order to reduce the overload to tissues and to facilitate the cleaning. Unfortunately, there are no reports in the literature regarding the rehabilitation of these patients.

Whatever the case, the use of prosthesis should be suspended in the presence of extensive lesions and/or associated candidiasis. Local treatment consists of the use of topical corticosteroids in the form of pastes and gels. Orabase formulations are little employed because most patients report discomfort with their use. Refractory cases may be treated with

topical immunosuppressants such as tacrolimus, and in this case it is interesting the serum titration of drug concentration, since there are reports of immunosuppressive concentrations with the use in ulcerated mucosa. In cases involving other organs or poor control of lesions, the treatment becomes systemic with the use of corticosteroids and/or immunosuppressive drugs.

Behcet's Disease

Behcet's disease is an inflammatory alteration of probable autoimmune origin, characterized by recurrent oral ulcers and other systemic manifestations, which characterize the Behcet's Syndrome. This disease usually has calm and acute episodes and affects young adults between 25 and 30 years [63-64]. The oral ulcers tend to be deeper and more painful than those found in pemphigus and cicatricial pemphigoid, and produce scarring in up to 10 days [65].

Diagnosis is difficult and usually perilesional microscopic evaluation shows vasculitis. Formations of venous and arterial thrombosis may be occurs; and active lesions are infiltrated by neutrophils [63, 66]. The presence of oral, genital and ocular lesions associated with absence of typical changes of pemphigoid enables the diagnosis of disease. The treatment is controversial and there is no established protocol. The use of topical corticosteroids in oral lesions has shown disparate responses among patients [63].

Because the disease usually has periods of remission and the lesions are usually localized, the use of complete dentures is more peaceful than that described for cases of pemphigus and pemphigoid. However, constant acute lesions impose restrictions on the use of prosthesis due to the pain in the areas of lesions. The use of topical anesthetic lidocaine and vaseline in the area of lesion has shown adequate to our patients.

The manufacture of complete dentures can be performed routinely since the appearance of lesions is not associated with local trauma, and the care related to the adaptation and installation are the same as for patients with pemphigus and pemphigoid, with the exception of the use of corticosteroids prior to installation, which is not usually employed.

At our service, there is a patient with Behcet's disease who has important limitation of opening the mouth caused by fibrosis in the lip commissure, due to recurrent lesions in this area. In this case, the dental treatment is extremely committed and the possibility of lip commissurotomy should be discussed in a multidisciplinary team.

Sjögren's Syndrome

Sjögren's syndrome is a chronic inflammatory disease of probable autoimmune etiology, characterized by xerostomia (dry mouth), xerophthalmia and lymphocytic infiltration into fragments of minor salivary glands [67-71]. The impairment can affect other exocrine glands such as pancreas and sweat glands.

The degree of xerostomia may vary among patients but the symptoms usually described are dysphagia for solids, aphthous lesions in oral cavity, and secondary infections - candidiasis (especially in users of complete denture). Treatment is usually symptomatic with the replacement of saliva through artificial solutions and use of sugarless chewing gum. The

use of muscarinic agonists such as pilocarpine has been described as effective to stimulate saliva production, yet its benefit is still controversial [72-74].

The manufacture of complete dentures should be performed in a conventional technique, and in case of lesions or lack of retention of the prosthesis, the use of artificial saliva in the inner surface of the prosthetic device usually facilitates its use. Maintenance should be done carefully and frequently, repeating needed rebasing and reinforcing hygiene as a preventive method against fungal infection.

In case of oral candidiasis, the treatment may be conservative with the use of topical solutions of nystatin or miconazole oral gel, as described before, associated with the hygiene of the prosthesis. The manufacture of implant-supported prosthesis is not contraindicated and may be a good alternative to the difficulty in adapting to conventional prosthesis.

Chronic Renal Failure (CRF)

It has been described a number of secondary oral changes to disease of base and to treatments for patients with chronic renal failure [75]. The treatment in these patients may be expectant such as dietary changes and drug control of complications; or invasive, resulting in dialysis and renal transplantation. Some medications such as cyclosporine and calcium channel-blocker have been associated with gingival enlargement in patients with teeth; however, it is not found in complete edentulous, so this topic will not be addressed [76-81].

Xerostomia can arise in individuals receiving dialysis, probably due to the restriction of liquid intake or as a side effect of drugs. This change is associated with the difficulty in adapting to complete dentures, dysphagia for solids and dysgeusia [82-84]. Care in the manufacture and maintenance of complete dentures are the same presented for the cases of Sjögren syndrome. However, it seems that the life expectancy of the implants may be reduced in these patients due to changes in bone remodeling [85-89].

It is important to highlight that individuals with chronic renal failure may have an ammonia-like odor and important changes such as metallic taste [83]. Sometimes patients seek dental causes for such changes. Other lesions described in the oral cavity are ulcerations, lichen-planus-like disease and hairy leukoplakia. Uremic syndromes were associated with hairy leukoplakia without the presence of Epstein-Barr virus [90, 91].

The incidence of fungal infection in transplant patients is not higher than in the general population, probably by the use of secondary prophylaxis [90]. Special attention should be given to the prevention of lip cancer and Kaposi's sarcoma that seem to have an increased incidence probably due to immunosuppression [92, 93].

Coagulation Disorders

This item was placed in the chapter because it is frequent that dental surgeon does not accept the care of patients with coagulation disorders. Regardless of the presented disorder (hereditary bleeding disorders, hemophilia, deficiency of other factors, platelet diseases; use of anticoagulants or antiplatelet, among others), there are no restrictions or special care in the manufacture or maintenance of complete dentures in this patients.

When mucosal lesion occurs during the manufacture of dentures, it will hardly have major bleeding, since in most cases it is superficial. In this case, local compression with gauze soaked in antifibrinolytic solutions (such as ϵ -aminocaproic acid and tranexamic acid) for 5 minutes usually contains minor bleeding.

Neurological Disorders and Dementia

Among neurological diseases and dementia, Parkinson's and Alzheimer's will be treated separately due to their incidence, prevalence in the elderly, and direct and indirect impairment of the oral cavity. However, regardless of the disease, the dental treatment and care should be decided case by case depending on the degree of psychomotor impairment, family structure and evolution of the disease. In these cases, the good sense of the professional and the detailed knowledge of patient conditions are imperative in deciding about the best type of rehabilitation in complete edentulous.

Parkinson's Disease

Parkinson's disease is a neurodegenerative disorder characterized by changes in movement and muscle control due to loss of dopaminergic neurons in the substantia nigra [94]. Its incidence is estimated between 1% and 5% of the population over 65 years, second only to Alzheimer's disease [95]. Despite its predilection for age above 40 years, it can also affect younger individuals.

Initially symptoms usually are bradykinesia and rigidity, and even with the replacement of dopamine and other drugs, after years, the disease tend to progress to dyskinesia and motor fluctuations. Dementia can affect up to 20% of terminal cases [96-98].

The moderate and advanced phases of the disease show important changes to the maintenance of oral health, especially with regard to the rehabilitation of complete edentulous patients. In 2011, Bakke et al. evaluated and compared 15 patients who had a moderate or advanced disease with a control group presenting similar oral characteristics and age to the studied group, and found a significant deterioration in performance and masticatory efficiency, reduction of oral functions, reduction of mouth opening, and difficulty to perform oral hygiene [99]. Other changes have been reported such as dysphonia, dysphagia, drooling, bruxism, xerostomia and increased complications with complete dentures [100-102].

The rehabilitation of complete edentulous patients should be discussed with the patient and family and scheduled for the possible evolution of the disease. In the absence of significant motor impairment, there are no restrictions or special care in the use of conventional prosthesis. However, in the moderate or severe phases, the lack of muscle control precludes the use of this type of rehabilitation due to the difficulty of adaptation and the worsening of dysphagia, since the control of the oral phase of digestion is often precarious. In these cases, a viable option is the manufacture of implant-retained removable prosthesis because they offer greater safety in use and facilitate cleaning. The patient often has difficulty in cleaning, then it is essential the education of the caregiver.

Despite to deserve attention, upper prosthesis generally can be made through conventional techniques at any stage of the disease. During terminal phase, with the presence of dyskinesia, the manufacture and use of dentures is usually not possible.

Alzheimer's Disease

Alzheimer's disease is also a progressive neurodegenerative disorder; however, it is different from Parkinson's disease because, in the initial phase, Alzheimer's disease is characterized by memory failure, and progresses slowly to severe and disabling dementia. Neuromotor changes, when occur, are found in advanced phase of disease. The infectious pulmonary complications resulting from aspirations for dysphagia associated with poor hygiene, malnutrition and inanition are usually the causes of death in these patients [103-107].

Alzheimer's has been showed as the most common dementing disorder in the world with a prevalence of 3% between the ages of 65 and 74 years, 19% between 75 and 84 years, and 47% in individuals over 85 years [108]. The literature provides reports of serious complications due to the precariousness of oral cavity hygiene in more advanced dementia. In a prospective evaluation, Friedlander et al (2006) described frequent forgetfulness of the removal of the prostheses for up to 3 consecutive days, patients with food debris on the bottom of the vestibule, and higher incidence of prosthetic stomatitis and secondary lesions due to the use of conventional dentures [106].

Regarding the manufacture of new prosthesis and the type of rehabilitation to be used in complete edentulous, the decision should be taken, as well as for Parkinson's disease, based on the clinical situation of the patient, and social and family structure in which he/she is inserted, since in many cases the presence of a caregiver is essential. Usually during advanced phases, the manufacture of new prosthesis is impracticable due to the impossibility of patient's cooperation; in these cases, the maintenance of the old relined denture is a good choice. At any phase of the disease, the appointments to evaluate the adequacy of hygiene and cognitive and motor skills for the use of dental prostheses should be done frequently with returns every 30 or 60 days [106, 107, 109-115].

It is important to highlight that a variety of drugs used to control the disease have dangerous medication interactions with dental drugs such as local anesthetics. Of interest to the management of complete dentures is the possibility of unwanted interaction of itraconazole with donepezil, galantamine, rivastigmine, quetiapine, carbamazepine and divalproex; for this reason, its use should be avoided in the treatment of fungal infections. There are no reports on interactions with fluconazole [116-119].

Secondary Oral Changes due to Treatment

Radiotherapy of Head and Neck Regions

Among the acute complications of radiotherapy, oral mucositis (Figure 11) has been reported to be the most frequent, debilitating, dose-dependent and dose-limiting, being

considered an inevitable side effect of radiation in head and neck [120-126]. Studies show that the incidence varies from 50% to 97% in patients undergoing radiotherapy of head and neck, and doubling the incidence of grade 3 mucositis when chemotherapy is associated [127-134]. Besides increasing the cost of treating, this complication is associated with increased morbidity and mortality, being the main factor responsible for the interruption and alteration in the *radiotherapy management in head and neck* [135, 136].

Moreover, a high rate of oral infection by *Candida* associated with mucositis has been reported in the literature. Apparently, mucositis predisposes to infection due to the loss of epithelial barrier, while the installed candidiasis provides the appearance of ulcers, making the epithelial repair more difficult [137-141].

Assessments of oral microflora during and after radiotherapy have shown a "shift" in both the variety and amount of microorganisms [141-143]. Prospective studies have demonstrated a high incidence of colonization (more than 93%) and infection (between 30 and 52%) by *Candida* in the oral cavity of patients undergoing radiotherapy of head and neck [139, 142, 144]. An important participation of non-albicans species in the colonization and oral infections in these patients has been demonstrated [139, 143].

Xerostomia is one of the most frequent chronic complications and responsible for damaging the quality of life of these patients [129, 145-148]. Studies have shown lower levels of xerostomia in patients undergoing conformal radiotherapy and IMRT, with reduction of irradiation in contra-lateral parotid gland when compared to patients undergoing conventional treatment [129].

Related to what was exposed, the complete dentures can be a complicating agent in the development of oral mucositis and candidiasis during radiotherapy. The patient should receive adjustments in the prosthesis and oral hygiene instructions prior the beginning of radiotherapy with the aim of reducing the trauma and local contamination. Other important information is the requirement that the prosthesis is removed during the manufacture of the mask and during all radiotherapy sessions. In the cases which it is not possible to obtain well-adapted and decontaminated prostheses, their use should be discontinued prior the beginning of treatment.

The patient should receive weekly follow-up and, in the first sign of tissue injury, the use of the prosthesis should be reduced, since each session of radiotherapy is expected worsening of the fragility of the oral mucosa. Specific treatments for the prevention and control of mucositis can be used [149-152].



Figure 11. Oral mucositis after radiotherapy.

In the post-radiotherapy, patients can and should receive well-adapted complete dentures; at our service, the manufacture of new prosthesis is only initiated after the clinical resolution of secondary oral lesions due to radiotherapy. Care in the manufacture and maintenance of the prosthesis are the same used for the previously described cases of xerostomia.

Chemotherapy

The chemotherapy regimens used today are the most diverse possible depending on the histologic type, tumor location and clinical performance of patients. Thus, the oral complications presented in each regime is directly associated with the type of drug and dosage used in the treatment [22, 25].

Possible complications in the oral cavity are the development of mucositis and oral infections; the first one depends on the presence of stomatotoxic agents and the second of myelotoxic agents capable of causing neutropenia [48, 49].

The necessary care with the prosthesis in the prevention and control of mucositis are the same presented for radiotherapy. And the management of infectious complications and the risk of bleeding are the same presented for the hematological diseases.

Use of Bisphosphonates

The use of bisphosphonates, particularly intravenous, has increased because of its important role in controlling tumor bone lesions (tumors of the prostate, breast, multiple myeloma, among others). Unfortunately, there are reports of its widespread use for control of no severe osteoporosis due to the ease of dosing [153, 154].

These drugs have been associated with extensive *osteonecrosis* of the jaws, difficult to control, and with severe infectious complications (Figures 12 and 13). The pathogenesis of the disease is still not completely understood, but it seems to be associated with the failure in the activation of osteoblasts by reducing the number and activity of osteoclasts. In addition, its anti-angiogenic effect seems to develop an essential role in the process of necrosis [155-158].

Although the majority of osteonecroses are associated with triggering events, such as dental extractions, periodontal surgery, and prosthetic trauma, their occurrence may also happen spontaneously. The risk of necrosis is directly related to the type of bisphosphonate, application form, dosage, time of use, pathology of base, and oral conditions. However, some genetic component seems to play key role in its development [157, 158]. In general, the user of injectable presentations has significantly higher risk to develop a lesion than the user of oral formulations, due to its potency and bioavailability [153, 154].

The dental extractions should be avoided in patients who use or have used intravenous bisphosphonates and its indication and realization should be provided by professionals with experience in dealing with these patients. Regarding the conventional complete dentures, special care have to be taken in their manufacture and maintenance, aiming at the minimization of lesion in the oral mucosa, since its laceration can serve as a trigger for necrosis.




Figure 12. Biphosphonate-relate osteonecrosis of the jaw (from left to right: oral lesion, panoramic radiography, and bone removed after surgery).



Figure 13. Biphosphonate-relate osteonecrosis of the jaw (from left to right: panoramic radiography, oral lesion showing the exposed jawbone). This patient had myeloma and died due to infectious complication in necrotic area.

The manufacture of the prosthesis in areas operated by necrosis should be made with extra caution; and in the presence of any sign of suppuration or bone exposure, its use should be discontinued. Osseointegrated implants so far are contraindicated in users of injectable forms of the drug.

Conclusion

The world population is aging and, associated with this event, there is an increased incidence of chronic and autoimmune diseases. Dentists have a key role in early detection and treatment of such diseases. Due to this fact, it is important to performed a detailed medical history and learn about the pathophysiology of each case, as well as to dominate the therapy applied and its possible complications. This knowledge is necessary to indicate the best type of prosthetic rehabilitation and care need to prevent and treat possible complications of the treatment.

References

- [1] Hughes, BB; Kuhn, R; Peterson, CM; Rothman, DS; Solórzano, JR; Mathers, CD; Dickson, JR. Projections of global health outcomes from 2005 to 2060 using the International Futures integrated forecasting model. *Bulletin World Health Organ*, 2011 7, 478-486.
- [2] Musgrove, P. Relations between health and development. *Boletín de la Oficina Sanitaria Panamericana*, 1993 2, 115-129.
- [3] Franco, M; Cooper, RS; Bilal, U; Fuster V. Challenges and opportunities for cardiovascular disease prevention. *American Journal of Medicine*, 2011 2, 95-102.
- [4] Selassie, M; Sinha, AC. The epidemiology and etiology of obesity: a global challenge. *Best Practice and Research Clinical Anaesthesiology*, 2011 1, 1-9.
- [5] Wildgust, HJ; Beary, M. Are there modifiable risk factors which will reduce the excess mortality in schizophrenia? *Journal of Psychopharmacology*, 2010 4, 37-50.
- [6] Cooper, LF. The current and future treatment of edentulismo. *Journal of Prosthodontics*, 2009 18, 116-122.
- [7] Carlsson, GE; Osmar, R. The future of complete dentures in oral rehabilitation. A critical review. *Journal of Oral Rehabilitation*, 2010 37, 143-156.
- [8] Turkyilmaz, L; Company, AM; McGlumphy, EA. Should edentulous patients be constrained to removable complete dentures? The use of dental implants to improve the quality of life for edentulous patients. *Gerodontology*, 2010 27, 3-10.
- [9] Niimi, M; Firth, NA; Cannon, RD. Antifungal drug resistance of oral fungi. *Odontology*, 2010 98, 15-25.
- [10] Lalla, RV; Latortue, MC; Hong, CH; Ariyawardana, A; D'Amato-Palumbo, S; Fischer, DJ; Martof, A; Nicolau-Galitis, O; Patton, LL; Elting, LS; Spijkervet, FKL; Brennan, MT. A systematic review of oral fungal infections in patients receiving cancer therapy. *Support Care Cancer*, 2010 18, 985-992.
- [11] Williams, DW; Kuriyama, TK; Silva, S; Malic, S; Lewis, MAO. Candida biofilms and oral candidosis: treatment and prevention. *Periodontology*, 2000 55, 250-265.
- [12] Chen, SCA; Slavin MA; Sorrell, TC. Echinocandin Antifungal Drugs in Fungal Infections. *Drugs*, 2011 71, 11-40.
- [13] Thompson GR; Patel, PK; Kirkpatrick, WR. Oropharyngeal candidiasis in the era of antiretroviral therapy. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology*, 2010 109, 488-495.
- [14] Klein, RS; Harris, CA; Small, CB; Moll, B; Lesser, M; Friedland, GH. Oral candidiasis in high-risk patients as the initial manifestation of the acquired immunodeficiency syndrome. *The New England Journal of Medicine*, 1984 311, 354-358.
- [15] Fidel, PL Jr. Distinct protective host defenses against oral and vaginal candidiasis. *Medical Mycology*, 2002 40, 359-375.
- [16] Williams, DW; Lewis, MA. Isolation and identification of Candida from oral cavity. *Oral Dis.eases*, 2000 6, 3-11.
- [17] Sakamoto, M; Umeda, M; Benno, Y. Molecular analysis of human oral microbiota. *Journal of Periodontal Research*, 2005 40, 277-285.

-
- [18] Siqueira, JF Jr; Rocas, IN. Exploiting molecular methods to explore endodontic infections: Part 1. Current molecular technologies for microbiological diagnosis. *Journal of Endodontics*, 2005 31, 411-423.
- [19] Odds, FC. *Candida and candidiasis. (Second edition)*. London: Baillière Tindall; 1988.
- [20] Ligouri, G; Lucariello, A; Colella, G; De Luca, A; Marinelli, P. Rapid identification of *Candida* species in oral rinse solutions by PCR. *Journal of Clinical Pathology*, 2007 60, 1035-1039.
- [21] White, PL; Williams, DW; Kuriyama, T; Samad, SA; Lewia, MA; Barners, RA. Detection of *Candida* in concentrated oral rinse cultures by real-time PCR. *Journal of Clinical Microbiology*, 2004 42, 2101-2107.
- [22] Wingard, JR. Importance of *Candida* species other than *C. albicans* as pathogens in oncology patients. *Clinical Infectious Diseases Journal*, 1995 20, 115-125.
- [23] Pfaller, MA; Diekema, DJ; Jones, RN; Messer, SA; Hollis, RJ. Trends in antifungal susceptibility of *Candida* ssp. isolated from pediatric and adult patients with bloodstream infections: SENTRY Antimicrobial Surveillance Program, 1997 to 2000. *Journal of Clinical Microbiology*, 2002 40, 852-856.
- [24] Perezous, LF; Flaitz, CM; Goldschmidt, ME; Engelmeier, RL. Colonization of *Candida* species in denture wearers with emphasis on HIV infection: a literature review. *Journal of Prosthetic Dentistry*, 2005 93, 288-293.
- [25] Schelenz, S; Abdallah, S; Stubbings, H; Gow, I; Baker, P; Hunter PR. Epidemiology of oral yeast colonization and infection in patients with hematological malignances, head neck and solid tumors. *Journal of Oral Pathology and Medicine*, 2011 40, 83-89.
- [26] Clark, TA; Haijeh, RA. Recent trends in the epidemiology of invasive mycoses. *Current Opinion in Infectious Diseases*, 2002 15, 204-215.
- [27] Haijeh, RA; Sofair, AN; Harrison, LH; Lyon, GN; Arthington-Skaggs, BA; Mirza, SA; Phelan, M; Morgan, J; Lee-yng, W; Ciblak, MA; Benjamin, LE; Sanza, LT; Huie, S; Yeo, SF; Brand, ME; Wrnock, DW. Incidence of bloodstream infections due to *Candida* species and in vitro susceptibilities of isolates collected from 1998 to 2000 in a population-based active surveillance program. *Journal of Clinical Microbiology*, 2004 42, 1519-1527.
- [28] Loeffler, J; Stevens, DA. Antifungal drug resistance. *Clinical Infectious Diseases*, 2003 36, S31-41.
- [29] Withe, TC; Marr, KA; Bowden, RA. Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clinical Microbiology*, 1998 11, 382-402.
- [30] Perea, S; Lopez-Ribot, JL; Kirkpatrick, WR; Mcate, Rk; SAntillán, RA; Martínez, M; Calabrese, D; Sanglard, D; Patterson, TF. Prevalence of molecular mechanisms of resistance to azole antifungal agents in *Candida albicans* strains displaying high-level fluconazole resistance isolated from human immunodeficiency virus-infected patients. *Antimicrobial Agents and Chemotherapy*, 2001 45, 2676-2684.
- [31] Albertson, GD; Niimi, M; Cannon, RD; Jenkinson, HF. Multiple efflux mechanisms are involved in *Candida albicans* clinical fluconazole resistance. *Antimicrobial Agents and Chemotherapy*, 1996 40, 2835-2841.
- [32] Bartie, KL; Willians, DW; Wilson, MJ; Potts, AJ; Potts, AJ; Lewis, MAO. Differential invasion of *Candida albicans* isolates in an in vitro model of oral candidosis. *Oral Microbiology and Immunology*, 2004 19, 293-296.

-
- [33] LaFleur, MD; Kumamoto, CA; Lewis, K. *Candida albicans* biofilms produce antifungal-tolerant persister cells. *Antimicrobial Agents and Chemotherapy*, 2006 50, 3839-3846.
- [34] Leonhardt, A; Renvert, S; Dahlén, G. Microbial findings at failing implants. *Clinical Oral Implants Research*, 1999 10, 339-345.
- [35] Price, CI; Waters, MG; Williams, DW; Lewis, MAO; Sticker, D. Surface modification of an experimental silicone rubber aimed at reducing initial *Candida* adhesion. *Journal of Biomedical Materials Research*, 2002 63, 122-128.
- [36] Paranhos, HFO; Silva-Lovato, CH. Comparative study of methods for quantification of biofilm on complete dentures. *Brazilian Oral Research*, 2004 18, 215-223.
- [37] Paranhos, HFO; Silva-Lovato, CH; Cherubini, V; Macedo, LD; de Souza, RF. Distribution of biofilm on internal and external surfaces of upper complete dentures: the effect of hygiene instruction. *Gerontology*, 2007 24, 162-168.
- [38] Earlandsen, JEPP; Vallor, AC; Kirkpatrick, WR; Thompson GR 3rd; Herrera, ML; Wickers, BL. Current prevalence of *Candida* spp. colonization, oropharyngeal candidiasis (OPC) and fluconazole (FLU) resistance in HIV/AIDS patients using combined microbial and molecular methods of detection. In: *Programs and Abstracts of the 48th Annual IcaAC/IDSA 46-th Annual Meeting*, Washington, D.C., 2008. Poster M-718.
- [39] Lacasse, M; Fortier, C; Chakir, J; Cote, L; Deslauriers, N. Acquired resistance and persistence of *Candida albicans* following oral candidiasis in the mouse: a model of the carrier state in humans. *Oral Microbiology and Immunology*, 1993 8, 313-318.
- [40] Deslauriers, N; Cote, L; Montplaisir, S; de Repentigny, L. Oral carriage of *Candida albicans* in murine AIDS. *Infection and Immunity*, 1997 65, 661-667.
- [41] de Repentigny, L; Aumont, F; Ripeau, JS; Fiorillo, M; Kay, DG; Hanna, Z; Jolicoeur, P. Mucosal candidiasis in transgenic mice expressing human immunodeficiency virus type 1. *Journal of Infectious Diseases*, 2002 185, 1103-1114.
- [42] Mercante, DE; Leigh, JE; Lilly, EA; McNulty, K; Fidel, PL Jr. Assessment of the association between HIV viral load and CD4 cell count on the occurrence of the oropharyngeal candidiasis in HIV-infected patients. *Journal of Acquired Immune Deficiency Syndrome*, 2006 42, 578-583.
- [43] Cauda, R; Tacconelli, E; Tumbarello, M; Morace, G; De Bernardis, F; Torosantucci, A; Cassone, A. Role of protease inhibitors in preventing recurrent oral candidosis in patients with HIV infection: a prospective case-control study. *Journal of Acquired Immune Deficiency Syndrome*, 1999 21, 20-25.
- [44] Salles, AES; Macedo, LD; Fernandes, RAG; Silva-Lovato, CH; Paranhos, HFO. Comparative analysis of biofilm levels in complete upper and lower dentures after brushing associated with specific denture paste and neutral soap. *Gerodontology*, 2007 24, 217-223.
- [45] de Souza, RF; de Freitas, HOP; Lovato da Silva, CH; Abu-Naba'a, L; Fedorowicz, Z; Gurgan, CA. Interventions for cleaning dentures in adults. *Cochrane Database of Systematic Reviews*, 2009 4, CD007395.
- [46] Souza, RF; Regis, RR; Nascimento, C; Paranhos, HFO; Silva-Lovato, CH. Domestic use of a disclosing solution for denture hygiene: a randomized trial. *Gerontology*, 2009 27, 1-6.

- [47] Adult Prevention and Treatment of Opportunistic Infections Guidelines Working Group. *Guidelines for prevention and treatment of opportunistic infections in HIV-infected adults and adolescents*. Accessed January 25, 2010.
- [48] Sonis, ST; Oster, G; Fuchs, H; Bellm, L; Bradford, WZ; Edelsberg, J; Hayden, V; Eilers, J; Epstein, JB; LeVeque, FG; Miller, C; Peterson, DE; Schubert, MM; Spijkervet, FK; Horowitz, M. Oral mucositis and clinical and economic outcomes of hematopoietic stem-cell transplantation. *Journal of Clinical Oncology*, 2001 21, 384-388.
- [49] Chen, YK; Hou, HA; Chow, JM; Chen, YC; Hsueh, PR; Tien, HF. The impact of oral herpes simplex virus infection and candidiasis on chemotherapy-induced oral mucositis among patients with hematological malignancies. *European Journal of Clinical Microbiology and Infectious Diseases*, 2011 30, 753-759.
- [50] De Gregorio, MW; Lee, WM; Ries, CA. Candida infections in patients with acute leukemia: ineffectiveness of nystatin prophylaxis and relationship between oropharyngeal and systemic candidiasis. *Cancer*, 1982 50, 2780-2784.
- [51] Dreizen, S; Bodey, GP; Valdivieso, M. Chemotherapy associated oral infections in adults with solid tumors. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 1983 55, 113-120.
- [52] Micozzi, A; Cartoni, C; Monaco, M; Martino, P; Zittoun, R; Mandelli, F. High incidence of infectious gastrointestinal complications observed on patients with acute myeloid leukemia receiving intensive chemotherapy for first induction of remission. *Support Care Cancer*, 1996 4, 294-297.
- [53] Li, L; Redding, S; Dongari-Bagtzoglou. *Candida glabrata*, an emerging oral opportunistic pathogen. *Journal of Dental Research*, 2007 86, 204-215.
- [54] Mealey, BL; Ocampo, GL. Diabetes mellitus and periodontal disease. *Periodontology*, 2007 44, 127-153.
- [55] Darwazeh, AM; Lamey, PJ; Samaranayake, LP; MacFarlane, TW; Fischer, BM; Mac Cuish, AC. The relationship between colonization, secretor status and *in-vitro* adhesion of *Candida albicans* to buccal epithelial cells from diabetics. *Journal of Medical Microbiology*, 1990 33, 43-49.
- [56] Manfredi, M; Mccullough, MJ; Al-Karaawi, ZM; Vescovi, P; Porter, SR. *In vitro* evaluation of virulence attributes of Candida spp. isolated from patients affected by diabetes mellitus. *Oral Microbiology and Immunology*, 2006 21, 183-219.
- [57] Knudson, RM; Kalaaji, AN; Bruce, AJ. The management of mucous membrane pemphigoid and pemphigus. *Dermatologic Therapy*, 2010 23, 268-280.
- [58] Ahmed, AR; Hombal, SM. Cicatricial pemphigoid. *International Journal of Dermatology*, 1986 25, 90-96.
- [59] Laskaris, G; Sklavounou, A; Stratigos, J. Bullous pemphigoid, cicatricial pemphigoid and pemphigus vulgaris: a comparative clinical survey of 278 cases. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 1982 54, 656-662.
- [60] Mondino, BJ; Sutart, JB. Ocular cicatricial pemphigoid. *Ophthalmology*, 1981 88, 95-100.
- [61] Chan, LS; Ahmed, AR; Anhalt, GF; Bernauer, W; Cooper, KD; Elder, MJ; Fine, JD; Foster, CS; Ghohestani, R; Hashimoto, T; Hoang-Xuan, T; Kirtschig, G; Korman, NJ; Lightman, S; Lozada-Nur, F; Marinkovich, MP; Mondino, BJ; Prost-Squarcioni, C; Rogers, RS 3rd; Setterfield, JF; West, DP; Wojnarowska, F; Woodley, DT; Yancey,

- KB; Zillikens, D; Zone JJ. The first international consensus on mucous membrane pemphigoid. *Archives of Dermatology*, 2002 138, 370-379.
- [62] Fleming, TE; Korman, NJ. Cicatricial pemphigoid. *Journal of the American Academy of Dermatology*, 2000 43, 571-591.
- [63] Sakane, T; Takeno, M; Suzuki, N; Inaba, G. Behçet's disease. *The New England Journal of Medicine*, 1999 21, 1284-1291.
- [64] Yurdakul, S; Hamuryudan, V; Yazici, H. Behçet syndrome. *Current Opinion in Rheumatology*, 2004 16, 38-42.
- [65] Zoubolis, CC. Epidemiology of Adamantiades-Behçet's disease. *Annales de Médecine Interne*, 1999 150, 488-487.
- [66] Ehrlich, GE. Vasculitis in Behçet's disease. *International Reviews of Immunology*, 1997 14, 81-88.
- [67] Esch, TR. Pathogenesis factors in Sjögren's syndrome: recent developments. *Critical Reviews in Oral Biology and Medicine*, 2001 12, 244-251.
- [68] Jonsson, R; Moen, K; Vestrheim, D; Szodoray, P. Current issues in Sjögren's syndrome. *Oral Dis.eases*, 2002 8, 130-140.
- [69] Moutsopoulos, HM; Chused, TM; Mann, DL; Klippel, JH; Fauci, AS; Frank, MM; Lawley, TJ; Hamburger, MI. Sjögren's syndrome (Sicca syndrome): current issues. *Annals of Internal Medicine*, 1980 92, 212-226.
- [70] Rehman, HU. Sjögren's syndrome. *Yonsei Medical Journal*, 2003 44, 947-954.
- [71] Yamamoto, K. Pathogenesis of Sjögren syndrome. *Autoimmunity Reviews*, 2003 2, 563-570.
- [72] Nelson, JD; Friedlaender, M; Yeatts, RP; Yee, R; McDermott, M; Orlin, S; Gallagher, SC; Shrotriya, RC. Oral pilocarpine for symptomatic relief of keratoconjunctivitis sicca in patients with Sjögren's syndrome. The MGI PHARMA Sjögren's syndrome study group. *Advances in Experimental Medicine and Biology*, 1998 483, 979-983.
- [73] Ono, M; Takamura, E; Shinozaki, K; Tsumura, T; Hmano, T; Yagi, Y; Tsubota, K. Therapeutic effect of cevimeline on dry eye in patients with Sjögren's syndrome: a randomized, double-blind clinical study. *American Journal of Ophthalmology*, 2004 38, 6-17.
- [74] Petrone, D; Condemi, JJ; Fife, R; Gluck, O; Cohen, S; Dalgin, P. A double-blind, randomized, placebo-controlled study of cevimeline in Sjögren's syndrome patients with xerostomia and keratoconjunctivitis sicca. *Arthritis and Rheumatism*, 2002 46, 748-754.
- [75] Proctor, R; Kumar, N; Stein, A; Moles, D; Porter, S. Oral and Dental Aspects of Chronic Renal Failure. *Journal of Dental Research*, 2005 84, 199-208.
- [76] Ellis, JS; Seymour, RA; Steele, J; Robertson, P; Butler, TJ; Thomason, JM. Prevalence of gingival overgrowth induced by calcium channel blockers: a community based study. *Journal of Periodontology*, 1999 70, 63-67.
- [77] Ress, TD; Levine, RA. Systemic drugs as a risk factor for periodontal disease initiation and progression. *Compendium*, 1995 16, 20,22,26.
- [78] Seymour, RA; Smith, DG. The effect of a plaque control program on the incidence and severity of ciclosporina-induced gingival changes. *Journal of Clinical Periodontology*, 1991 18, 107-110.
- [79] Semour, RA; Thomason, JM; Ellis, JS. Oral and dental problems in the organ transplant patient. *Dental Update*, 1994 21, 209-212.

- [80] Somacarrera, ML; Hernandez, G; Acero, J; Moskow, BS. Factors related to the incidence and severity of ciclosporina-induced gingival overgrowth in trasplant patients. A longitudinal study. *Journal of Periodontology*, 1994 65, 671-675.
- [81] Pernu, HE; Oikarinen, K; Hietanen, J; Knuuttila, M. Verapamil-induced gingival overgrowth: a clinical histologic and biochemic approach. *Journal of Oral Pathology and Medicine*, 1989 18, 422-425.
- [82] Kao, CH; Hsieh, JF; Tsai, SC; Ho, YJ; Chang, HR. Decrease salivary function in patients with end-stage renal disease requiring haemodialysis. *American Journal of Kidney Diseases*, 2000 36, 1110-1114.
- [83] Kho, H; Lee, S; Chung, SC; Kim, YK. Oral manifestations and salivary flow rate, ph, and buffer capacity in patients with end-stage renal disease undergoing haemodialysis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 1999 88, 316-319.
- [84] Porter, SR; Hegarty, A; Scully, C. An update of the etiology and management of xerostomia. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2004 97, 28-46.
- [85] Craig, RG; Kotanko, P; Levin NW. Periodontal deseases – a modifiable source of systemic inflammation for the end-stage renal disease patient on haemodialyis therapy? *Nephrology Dialysis Transplantation*, 2007 22, 312-315.
- [86] Diakiewicz, M; Wojtowicz, A. Problems and algorithm of dental procedures in patients on renal replacement therapy. In: Rutkowski B, (ed.) *Dialysis in Medical Practice*, Makmedia, 2004.
- [87] Kukula, K; Wojtowicz, A; Diakiewicz, M. The linear radiometric analysis of the jaws in renal osteodystrophy: introduction to implant-prosthetic treatment. *Protetyka Stomatologiczna*, 2005 55, 336-343.
- [88] Nadimi, H; Bergagmini, J; Lilien, B. Uremic mixed bone disease. A case report. *International Journal of Oral and Maxillofacial Surgery*, 1993 22, 268-270.
- [89] Okada, H; Davies, JE; Yamamoto, H. Brown tumour of the maxila in a patient with secondary hyperparathyroidism: a case study involving immunohistochemistry and electron microscopy. *Journal of Oral and Maxillofacial Surgery*, 2000 58, 233-238.
- [90] King, GN; Healy, CM; Glover, MT; Kwan, JT; Williams, DM; Leigh, IM; Thornhill, MH. Prevalence and risk factors associated with leukoplakia, hairy leukoplakia, erythematous candidiasis, and gingival hyperplasia in renal transplant recipients. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 1994 78, 718-726.
- [91] Ross, WF 3rd; Salisbury, PL 3rd. Uremic stomatitis associated with undiagnosed renal failure. *General Dentistry*, 1994 42, 410-412.
- [92] Farge, D. Kaposi's sarcoma in organ transplant recipients. *The European Journal of Medicine*, 1993 2, 339-343.
- [93] Regev, E; Zeltser, R; Lustman, J. Lip carcinoma in renal allograft recipient with long-term immunosuppressive therapy. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 1992 73, 412-414.
- [94] Farrer, M; Wavrant-De Vrieze, F; Crook, R; Boles, L; Perez-Tur, J; Hardy, J; Johson, WG; Steele, J; Maraganore, D; Gwinn, K; Lynch, T. Low frequency of alpha-synuclein mutations in familial Parkinson's disease. *Annals of Neurology*, 1998 43, 394-397.

-
- [95] Bostantjopoulou, S; Katsarbu, Z; Papadimitriou, A; Veletza, V; Hatzigeorgiou, G; Less, A. Clinical features of parkinsonian patients with the alpha-synuclein (G209A) mutation. *Movement Disorders*, 2001 16, 1007-1013.
- [96] Hughes, AJ; Daniel, SE; Ben-Shlomo, Y; Lees, AJ. The accuracy of diagnosis of parkinsonian syndromes in a specialist movement disorder service. *Brain*, 2002 125, 861-870.
- [97] Marsden, CD; Parkes, JD; Quinn, N; Fahn, S. Movement disorders. In: Marsden, CD; Parkes, JD; Quinn, N; Fahn, S. *Neurology Part 2*. 1982, 96-122.
- [98] Quinn, NP. Classification of fluctuations in patients with Parkinson's disease. *Neurology*, 1998 51, S25-S29.
- [99] Bake, M; Larsen, SL; Lautrup, C; Karlsborg, M. Orofacial function and oral health in patients with Parkinson's disease. *European Journal of Oral Sciences*, 2011 119, 27-32.
- [100] Friedlander, AH; Mahler, M; Norman, KM; Ettinger, RL. Parkinson's disease: systemic and orofacial manifestations, medical and dental management. *Journal of American Dental Association*, 2009 140, 658-669.
- [101] Karlsson, S; Persson, M; Johnels, B. Levodopa induced ON-OFF motor fluctuations in Parkinson's disease related to rhythmical masticatory jaw movements. *Journal of Neurology, Neurosurgery and Psychiatry*, 1992 55, 304-307.
- [102] Packer, M; Nikitin, V; Coward, T; Davis DM; Fiske, J. The potential benefits of dental implants on the oral health quality of life with Parkinson's disease. *Gerontology*, 2009 26, 11-18.
- [103] Atri, A; Shaughnessy, LW; Locascio JJ, Growdon, JH. Long-term course and effectiveness of combination therapy in Alzheimer disease. *Alzheimer Disease and Associated Disorders*, 2008 22, 209-221.
- [104] Bacanu, SA; Devlin, B; Chowdari, KV; DeKosky, ST; Nimgaonkar, VL; Sweet, RA. Heritability of psychosis in Alzheimer disease. *American Journal of Geriatric Psychiatry*, 2005 13, 624-627.
- [105] Bird TD; Miller, BL. Alzheimer's disease and primary dementias. In: Kasper, D; Fauci, A; Branwald, E, et al. *Harrison's Principles of Internal Medicine, (17-th edition)*. McGaw-Hill, NY, 2393-2406.
- [106] Friedlander, AH; Norman, DC; Mahler, ME; Norman, KM; Yagiela, JA. Alzheimer's disease: Psychopathology, medical management and dental implications. *Journal of American Dental Association*, 2006 137, 1240-1251.
- [107] Hatipoglu, MG; Kabay, SC; Güven, G. The clinical evaluation of the oral status in Alzheimer-type dementia patients. *Gerontology*, 2010 10, 1-5.
- [108] Tang, MX; Cross, P; Andrews, H; Jacobs, DM; Small, S; Bell, K; Merchant, C; Antiguaa, R; Costa, R; Stem, Y; Mayeux, R. *Neurology*, 2001 56, 49-56.
- [109] Arai, K; Sumi, Y; Uematsu, H; Miura, H. Association between dental health behaviors, mental/psysical function and self-feeding ability among the elderly: a cross-sectional survey. *Gerodontology*, 2003 20, 78-83.
- [110] Friedlander, AH; Marder, SR. The psychopathology , medical management and dental implications of schizophrenia. *Journal of American Dental Association*, 2002 133, 603-610.
- [111] Limeback, H. Implications of oral infections on systemic in the institutionalized elderly with special focus on pneumonia. *Annals of Periodontology*, 1998 3, 262-275.

-
- [112] McEvoy, GK. *AHFS drug information 2005*. Bethesda, MD. American Society of Helth-System Pharmacists, 2005.
- [113] Ocasio, NA; Solomowitz, BH; Sher, MR. Dental management of the patients with Alzheimer's disease. *New York State Dental Journal*, 2000 66, 32-35.
- [114] Ostuni, E; Mohi, G. Communicating more effectively with the confuse or demented patients. *General Dentistry*, 1995 43, 264-266.
- [115] Priefer, BA; Robbins, J. Eating chances in mild-stage Alzheimer's disease: a pilot study. *Dysphagia*, 1997 12, 212-221.
- [116] Delfilippi, JL; Crismon, ML. Drug interactions with cholinesterase inhibitors. *Drugs Aging*, 2003 20, 437-444.
- [117] Goff, DC; Baldessari, RJ. Drug interactions with antipsychotic agents. *Journal of Clinical Psychopharmacology*, 1993 13, 57-67.
- [118] Haas, DA. Adverse drug interactions in dental practice: interactions associated with analgesics, part III in a series. *Journal of American Dental Association*, 1999 130, 397-407.
- [119] Hersh, EV; Moore, PA. Drug interactions in dentistry: the importance of knowing your CYPs. *Journal of American Dental Association*, 2004 135, 298-311.
- [120] Rose-Ped, AM; Bellm, LA; Epstein, JB; Trotti, A; Gwede, C; Fuchs, HJ. Complications of radiation therapy for head and neck cancers. The patient's perspective. *Cancer Nursing*, 2002 25, 461-467.
- [121] Talmi, YP. Quality of life issues in cancer of the oral cavity. *Journal of Laryngology and Otology*, 2002 116, 785-790.
- [122] Vissink, A; Jansma, J; Spijkervet, FK; Burlage, FR; Coppes, RP. Oral sequelae of head and neck radiotherapy. *Critical Reviews in Oral Biology and Medicine*, 2003 14, 199-212.
- [123] Epstein, JB; Emerton, S; Kolbinson, DA; Le, ND; Phillips, N; Stevenson-Moore, P; Osoba, D. Quality of life and oral function following radiotherapy for head and neck cancer. *Head and Neck*, 1999 21, 1-11.
- [124] Scully, C; Hodgson, T. Recurrent oral ulceration: aphthous-like ulcers in periodic syndromes. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2008 106, 845-852.
- [125] Sonis, ST; Eilers, JP; Epstein, JB; LeVeque, FG; Liggett Jr, H; Mulagha, MT; Peterson, DE; Rose, AH; Schubert, MM; Spijkervet, FKJ; Wittes, P. Validation of a new scoring system for the assessment of clinical trial research of oral mucositis induced by radiation or chemotherapy. Mucositis Study Group. *Cancer*, 1999 85, 2103-2113.
- [126] Sonis, ST; Peterson, DE; McGuire, DB; Williams, DA. Prevention of mucositis in cancer patients. *The Journal of National Cancer Institute Monographs*, 2001 29, 1-2.
- [127] Cooper, JS; Pajak, TS; Forastiere, AA; Jacobs, J; Campbell, BF; Saxman, SB; Kish, JA; Kim, HE; Cmelak, AJ; Rotman, M; Machtay, M; Ensley, JF; Chao, KS; Schultz, CJ; Lee, N; Fu, KK. Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *The New England Journal of Medicine*, 2004 350, 1937-1944.
- [128] Epstein, JB; Beaumont, JL; Gwede, CK; Murphy, B; Garden, AS; Meredith, R; Le, QT; Brizel, D; Isitt, J; Cella, D. Longitudinal evaluation of the oral mucositis weekly questionnaire-head and neck cancer, a patient-reported outcomes questionnaire. *Cancer*, 2007 109, 1914-1922.

-
- [129] Gomez, DR; Zhung, JE; Gomez, J; Chan, K; Wu, AJ; Wolden, SL; Pfister, DG; Shaha, A; Shah, JP; Kraus, KH; Wong, RJ; Lee, NY. Intensity-Modulated Radiotherapy in treatment of Oral Cavity Cancers. *International Journal of Radiation Oncology Biology Physics*, 2009 73, 1096-1103.
- [130] Madan, PD; Sequeira, PS; Shenoy, K; Shetty, J. The effect of three mouthwashes on radiation-induced oral mucositis in patients with head and neck malignancies: a randomized control trial. *Journal of Cancer Research Therapy*, 2008 4, 3-8.
- [131] Murphy, CK; Fey, EG; Watkins, BA; Wong, V; Rothstein, D; Sonis, ST. Efficacy of superoxide dismutase mimetic M40403 in attenuating radiation-induced oral mucositis in hamsters. *Clinical Cancer Research*, 2008 14, 4292-4297.
- [132] Rubenstein, EB; Peterson, DE; Schubert, M; Keefe, D; McGuire, D; Epstein, J; Elting, LS; Fox, PC; Cooksley, C; Sonis, ST. Clinical practice guidelines for the prevention and treatment of cancer therapy-induced oral and gastrointestinal mucositis. *Cancer*, 2004 100, 2026-2046.
- [133] Trotti, A; Bellm, LA; Epstein, JB; Frame, D; Fuchs, HJ; Gwede, CK; Komaroff, E; Nalysnyk, L; Zilberberg, MD. Mucositis incidence, severity and associated outcomes in patients with head and neck cancer receiving radiotherapy with or without chemotherapy: a systematic literature review. *Radiotherapy and Oncology*, 2003 66, 253-262.
- [134] Vuyyuri, SB; Hamstra, DA; Khanna, D; Hamilton, CA; Markwart, SM; Campbell, KC; Sunkara, P; Ross, BD; Rehemtulla, A. Evaluation of D-methionine as a novel oral radiation protector for prevention of mucositis. *Clinical Cancer Research*, 2008 14, 2161-2170.
- [135] Elting, LS; Cooksley, C; Chambers, M; Cantor, SB; Manzullo, E; Rubenstein, EB. The burdens of cancer therapy. Clinical and economic outcomes of chemotherapy-induced mucositis. *Cancer*, 2003 98, 1531-1539.
- [136] Sonis, ST; Oster, G; Fuchs, H; Bellm, L; Bradford, WZ; Edelsberg, J; Hayden, V; Eilers, J; Epstein, JB; LeVeque, FG; Miller, C; Peterson, DE; Schubert, MM; Spijkervet, FK; Horowitz, M. Oral mucositis and the clinical and economic outcomes of hematopoietic stem-cell transplantation. *Journal of Clinical Oncology*, 2001 19, 2201-2205.
- [137] Thaweboon, S; Thaweboon, B; Srithavaj, T; Choonharuangdej, S. Oral colonization of *Candida* species in patients receiving radiotherapy in the head and neck area. *Quintessence International*, 2008 39, e52-e57.
- [138] Beumer, J; Curtis, T; Harrison, RE. Radiation therapy of the oral cavity: sequelae and management, part 1. *Head and Neck Surgery*, 1979 1, 301-312.
- [139] Redding, SW; Zellars, RC; Kirkpatrick, WR; McAtee, RK; Caceres, MA; Fothergill, AW; Lopez-Ribot, JL; Bailey, CW; Rinaldi, MG; Patterson, TF. Epidemiology of oropharyngeal *Candida* colonization and infection in patients receiving radiation for head and neck cancer. *Journal of Clinical Microbiology*, 1999 37, 3896-3900.
- [140] Sixou, JL; Medeiros-Batista, O; Bonnaure-Mallet, M. Modifications of the microflora of the oral cavity arising during immunosuppressive chemotherapy. *European Journal of Cancer: Part B - Oral Oncology*, 1996 32B, 306-310.
- [141] Adamietz, AI; Rahn, R; Bottcher, HD; Schafer, V; Reimer, K; Fleischer, W. Prophylaxis with povidone-iodine against induction of oral mucositis by radiochemotherapy. *Support Care Cancer*, 1998 6, 373-377.

- [142] Jham, BC; Franca, EC; Oliveira, RR; Santos, VR; Kowalski, LP; Freire, ARS. Candida oral colonization and infection in Brazilian patients undergoing head and neck radiotherapy: a pilot study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2007 103, 355-358.
- [143] Leung, WR; Dassanayake, RS; Yau, JY; Jin, LJ; Yam, WC; Samaranayake, LP. Oral colonization, phenotypic, and genotypic profiles of Candida species in irradiated, dentate, xerostomic nasopharyngeal carcinoma survivors. *Journal of Clinical Microbiology*, 2000 38, 2219-2226.
- [144] Adelstein, DJ. Systemic chemotherapy for squamous cell head and neck cancer. *Expert Opinion on Pharmacotherapy*, 2003 4, 2151-2163.
- [145] Hey, J; Setz, J; Gerlach, R; Janich, M; Sehlleier, S; Schaller, HG; Gernhardt, CR; Kuhnt, T. Parotid-gland-sparing 3D conformal radiotherapy in patients with bilateral radiotherapy of the head and neck region - Results in clinical practice. *Oral Oncology*, 2009 45, e11-17.
- [146] Kielbassa, AM; Hinkelbein, W; Hellwig, E; Meyer-Luckel, H. Radiation-related damage to dentition. *The Lancet Oncology*, 2006 7, 326-335.
- [147] Lin, SC; Jen, YM; Chang, YC; Lin, CC. Assessment of xerostomia and its impact on quality of life in head and neck cancer patients undergoing radiotherapy, and validation of the Taiwanese version of the xerostomia questionnaire. *Journal of Pain and Symptom Manage*, 2008 36, 141-148.
- [148] Rosenthal, DI; Mendoza, TR; Chambers, MS; Asper, JA; Gning, I; Kies, MS; Weber, RS; Lewin, JS; Garden, AS; Ang, KK; Wang, S; Cleeland, CS. Measuring head and neck cancer symptom burden: the development and validation of the M. D. Anderson symptom inventory, head and neck module. *Head and Neck*, 2007 29, 923-931.
- [149] Arora, H; Pai, KM; Maiya, A; Vidyasagar, MS; Rajeev, A. Efficacy of He-Ne Laser in the prevention and treatment of radiotherapy-induced oral mucositis in oral cancer patients. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*, 2008 105, 180-186.
- [150] Bensadoun, RJ; Franquin, JC; Ciais, G; Darcourt, V; Schubert, MM; Viot, M; Dejou, J; Tardieu, C; Benezery, K; Nguyen, TD; Laudoyer, Y; Dassonville, O; Poissonnet, G; Vallicioni, J; Thyss, A; Hamdi, M; Chauvel, P; Demard, F. Low-energy He/Ne laser in the prevention of radiation-induced mucositis. A multicenter phase III randomized study in patients with head and neck cancer. *Support Care Cancer*, 1999 7, 244-252.
- [151] Genot-Klastersky, MT; Klastersky, J; Awada, F; Awada, A; Crombez, P; Martinez, MD; Jaivenois, MF; Delmelle, M; Vogt, G; Meuleman, N; Paesmans, M. The use of low-energy laser (LEL) for the prevention of chemotherapy- and/or radiotherapy-induced oral mucositis in cancer patients: results from two prospective studies. *Support Care Cancer*, 2008 16, 1381-1387.
- [152] Maiya, GA; Kumar, P; Rao, L. Effect of low intensity helium-neon (He-Ne) laser irradiation on diabetic wound healing dynamics. *Photomedicine and Laser Surgery*, 2005 23, 187-190.
- [153] King, AE; Umland, EM. Osteonecrosis of jaw in patients receiving intravenous or oral bisphosphonates. *Pharmacotherapy*, 2008 28, 667-677.
- [154] Lo, JC; O’Ryan, FS; Gordon, NP; Yang, J; Hui, RL; Martin, D; Hutchinson, M; Lathon, PV; Sanchez, G; Silver, P; Chandara, M; McCloskey, CA; Staffa, JA; Willy, M; Selby, JV; Go, AS. Prevalence of osteonecrosis of the jaw in patients with oral

- bisphosphonate exposure. *Journal of Oral and Maxillofacial Surgery*, 2010 68, 243-253.
- [155] Sedghizadeh, PP; Kumar, SK; Gorur, A; Schaudinn, C; Shuler, CF; Costerton, JW. Identification of microbial biofilms in osteonecrosis of the jaw secondary to bisphosphonate therapy. *Journal of Oral and Maxillofacial Surgery*, 2008 66, 767-775.
- [156] Stockmann, P; Vairaktaris, E; Wehrhan, F; Seiss, M; Schwartz, S; Spriewald, B; Neukam, FW; Nkenke, E. Osteotomy and primary wound closure in bisphosphonate-associated osteonecrosis of the jaw: a prospective clinical study with 12 months follow-up. *Support Care Cancer*. 2010 18, 449-460.
- [157] Such, E; Cervera, J; Terpos, E; Bagan, JV; Avaria, A; Gomez, I; Margaix, M; Ibanez, M; Luna, I; Cordon, L; Roig, M; Sanz, MA; Dimopoulos, MA; de la Rubia, J. CYP2C8 gene polymorphism and bisphosphonate-related osteonecrosis of jaw in patients with multiple myeloma. *Haematologica*, 2011 17 in press.
- [158] Wehrhan, F; Hyckel, P; Guentsch, A; Nkenke, E; Stockmann, P; Schlegel, KA; Neukam, FW; Amann, K. Bisphosphonate-associated osteonecrosis of jaw is linked to suppressed TGFbeta-1-signaling and increased Galectin-3 expression: A histological study on biopsies. *Journal of Translational Medicine*, 2011 4, 9:102.

Chapter VII

Laboratory Improvement of the Clinical Performance of Complete Dentures – A Review

***Luciana Assirati Casemiro¹, Soraia Marangoni¹,
André Marcelo Peruchi Minto¹
and Leandro de Dorigan de Macedo^{1,2,*}***

¹University of Franca, Dentistry School, Franca, SP, Brazil

²University of São Paulo at Ribeirão Preto, Medical School,
Hospital das Clínicas, Dentistry and Stomatology Service, Brazil

Abstract

Laboratory processing techniques have been refined in an attempt to achieve complete dentures with better clinical performance. Regardless of the type of denture base material (microwave-, light-, or heat-polymerized polymethyl methacrylate, for example), materials as well as handling and polymerization methods have been continuously modified with a view to obtaining more resistant dentures. In an attempt to improve the mechanical properties of base materials and to try to overcome the low impact and flexural strength of currently available dentures, which often results in clinical failure, various laboratory treatments have been proposed. The incorporation of reinforcing structures such as glass fiber, carbon fiber, Kevlar, aramid, and fiber-reinforced composite, which can be found in different sizes and shapes and are sometimes impregnated with substances such as silanes, has been successfully tested in some cases. However, incorporation of such compounds can culminate in undesirable effects such as larger amount of residual monomer, difficult polishing, and unsatisfactory aesthetics. Metal wires have also been used over time for this purpose, as well as substances such as methyl acetate and methyl formate. The increased bond strength between artificial teeth and the denture base material has also been relentlessly pursued.

* E-mail address: lucianacasemiro@hotmail.com.

New positive results are being obtained by means of techniques that promote interpenetration of the polymer networks of the two materials and make use of substances such as dichloromethane. To improve the biological properties of dentures, simple procedures such as immersion in water for defined time periods has been evaluated, in order to reduce their toxicity. Studies have shown that dibenzoyl peroxide, used as initiator in polymethyl methacrylate resin and associated with allergic processes, is released from conventional dentures after polymerization, but this event can be substantially reduced with completion of an additional polymerization cycle. Still, in order to reduce dentures contamination, several antimicrobial agents such as silver have been incorporated into the base material, which has led to positive effects with regard to their antimicrobial action. However, decrease in flexural and impact strength has sometimes been observed. In this context, this book chapter intends to update the reader on materials and laboratory techniques that can optimize the physical, mechanical, and biological properties of dentures that are considered to be the most relevant to the clinical performance of complete dentures.

Introduction

The best clinical performance of dentures has been pursued relentlessly over time. Among the factors responsible for the success of the rehabilitation treatment can be highlighted the materials selection and processing laboratory techniques, that can be controlled by professionals involved in the construction and repair of these prostheses.

Regardless of the material selected, the working parameters must be strictly defined and directed to obtain a higher conversion of monomers into polymers, since the presence of residual monomer affect the clinical performance of dentures by changing its physical, mechanical and biological properties [1-6]. Unsatisfactory results of flexural strength, for example, may be related to recurrent fractures of the middle line, as well as low impact strength values increase the likelihood of prosthesis fracture when it collides with hard surfaces [4-7]. High values of sorption and solubility can compromise the dimensional stability of the prosthesis, submit the material to an internal stress and decrease the values of mechanical strength [8], depending on the plasticizer effect of the polymer matrix [9]. In addition, the solubility releases toxic substances to the environment, which may cause negative reactions in oral tissues [4,10]. Thus, in addition to improving the working parameters for each material, researchers have evaluated the performance of reinforcing structures for acrylic resin to optimize the values of resistance and reduce the chances of fracture. Moreover, some alternatives to improve the adhesion of artificial teeth to the base material have been tested successfully. Given the above, some considerations on the denture laboratory techniques recently presented in scientific literature and its clinical consequences have been briefly reviewed below.

Reinforcement of Acrylic Resin

The use of reinforcement structures to improve the mechanical properties of denture base resins is not new. Over time, different materials were developed and put into practice, but until today questions about which material is most effective still remain.

Glass fibers, aramid and nylon were tested as reinforcement materials of acrylic resins and their benefits were proven through the improvement of flexural strength; however, the first one still promote the best results [11]. Reinforcement steel structures have also been used with success for a long time in order to reinforce the mechanical resistance of acrylic resins [12]. However, due to the predictability of clinical performance and the ease of application, glass fibers have been widely used than other types of reinforcement structures. Therefore, some considerations on the use of glass fibers are made in the following.

It is known that glass fibers promote the improvement of the mechanical properties of acrylic resins because their modulus of elasticity is high, which allows that the stress of chewing is absorbed with less deformation of the denture [11]. As a result, the flexural strength of resins reinforced by glass fibers is from 1.54 to 1.75 times higher than the unreinforced resins [13]. It was also reported that the use of glass fibers is able to promote a mechanical reinforcement of complete denture in a significant way (with average values of 919 N compared to 677 N for unreinforced materials) [14].

However, the positive clinical results are dependent on factors such as the stable union in the interface resin/fiber. Silane contributes notoriously to increase the adhesion between the resin and reinforcement fibers and promotes surface protection of these fibers, preventing damage during the handling [15]. Like the silane, the monomers also increase the adhesive strength of fibers, but this substance may remain without reacting inside the resin, having a negative influence on the mechanical properties of the material [16].

Fibers pre-impregnated by the manufacturer with polymers and silanes are available in the commercial market; others called pure or not impregnated fibers require a previously surface treatment made by the dental surgeon or dental prosthesis technician [17]. It is highlighted that in the latter case should have the maximum of care related to the complete wetting of the fibers by the bonding agent [18]. Contamination of the fiber reactive surface during handling, especially in the case of impregnated fibers, should also be carefully prevented.

The preference for pre-impregnated systems is because they eliminate the clinical stages which can result in errors, besides being easier to manipulate due to the decrease of their elastic memory and minor fraying of the fibers in the handling [19,20]. The pre-impregnated fibers also provide the highest values of flexural strength [21].

There are reports of clinical follow up for up to 5 years which showed that polymer pre-impregnated E-glass unidirectional fibers and pre-impregnated E-glass fiber nets promote successfully the reinforcement of acrylic resin complete dentures, especially for patients with high occlusal load [22]. Regarding the type of fiber glass, there is no scientific consensus on the subject, however some authors affirm that the unidirectional fibers promote a structural reinforcement more efficient than the braided ones [23,24].

The ideal positioning of the reinforcement fibers in the denture base resins diverges in the literature. A recent study demonstrated that the location (anterior region, middle region, anterior and posterior regions, and under the ridge lap) of the glass fibers reinforcement does not seems to influence in the flexural strength of acrylic resins [13]. However, a previous study revealed that the orientation of reinforcement fibers has significant influence on the prognosis of rehabilitation treatment with complete dentures, being an angle of 90° its ideal position in relation to the potential area for fracture and closer to the surface of the prosthesis [20,25]. There is also indication of placement of the reinforcement fibers at polished side to reduce the risk of fractures due to stress concentration in this region [26]. The amount of

fibers may also affect the structural reinforcement, having reports that the application of 1.0% of fibers promotes ideally the obtained effect [27].

The success of the use of reinforcement fibers in the manufacture of complete dentures is established; however, it depends on the strict following of the guidelines recommended for each material, both in clinical and laboratory stages.

Denture Repair

Fractures in complete denture bases are a relatively common clinical occurrence. Therefore, the need to develop effective repair techniques is justified. However, what is seen clinically is that this region is weaker and subject to re-fracture, which occurs predominantly between the base material and that used to repair, setting a fracture of adhesive nature predominantly. Factors such as the type of material used for repair and its surface contour as well as the treatment given to the denture base resin before the repair material application influence on the clinical results [28].

The desirable requirements of a material used for repair of complete denture bases are: color stability, flexural strength, dimensional stability, ease to work and accessible cost [29,30]. Among the materials indicated for the repair of denture base, the glass fibers are highlighted. However, regardless the materials selected for repair, it is known that repaired resins have lower flexural strength than intact resins, without fractures or repairs [31].

In the repair of heat-polymerized acrylic resins, which are the most commonly used for making removable dentures bases, a study showed that the use of autopolymerized resin reinforced with unidirectional and woven glass fibers is the most effective method to restore the mechanical properties of these materials [32].

Some alternatives to increase the bond strength between the repair and the base material have been evaluated such as the application of acetone, methylene chloride, or methylmetacrylate for periods from 30 to 180 seconds [33].

The continuity of the search for an effective and durable mechanism of denture repair is justified since the complete denture will be submitted with intensity to occlusal loads and their variations.

Bond Strength between Tooth and Base

The bond of artificial teeth to the denture base resins is a recurring problem in dental clinic. There is information in the literature that certain chemical [34-36] or mechanical treatment [37] in the ridge-lap artificial tooth surface may increase its adhesive strength to denture base materials.

The application of dichloromethane in artificial teeth, in the area of contact with the base resin, potentiates its bond with auto- and heat-cured resins [34,35]. The use of an acrylate bonding agent to enhance the shear bond strength was also recommended as a resource to promote the bond of highly cross-linked artificial teeth and light-polymerized urethane dimethacrylate [36]. However, there is information in the literature that the realization of

mechanical retention potentiates the shear bond strength values of resin and artificial tooth bond more than the surface treatment with monomers [37].

The parameters of the polymerized cycles also influence in the results of adhesion, being the shortest responsible for the lowest bond strength values [38,39]. Regarding the characteristics of laboratory manipulation, it should take special care in removing the wax residue, so there is no reduction in the values of adhesion between artificial tooth and base material [40].

Regarding the type of complete denture base resin, it is known that cross-linked artificial teeth (IPN), widely used nowadays due to their enhanced physical and mechanical properties; adheres easier with the heat-cured polymethylmethacrylate than with the microwave-polymerized materials [35].

Another aspect that has been highlighted as a factor that contributes to the adhesion of artificial teeth to the base material refers to the selection of both from the same manufacturer [41]. Ideally, manufacturers should indicate the compatibility of the denture base materials with the artificial teeth as well as the laboratory treatments indicated to improve the adhesive strength, then more favorable clinical results would be obtained.

Biocompatibility

The polymers used for the manufacture of denture base have shown different degrees of cytotoxicity, caused by their unreacted components and byproducts remaining after the polymerization process [42-44]. Clinically, the signs and symptoms most often associated with this fact are erythema, erosion of the oral mucosa and burning sensation of the mucosa and tongue [45].

In the prosthetic laboratories, the acrylic resin is the material that presents the occurrence of the highest number of adverse reactions, being responsible for developing contact dermatitis in the hands of 61% of the dental technicians [46].

Among the components released into the oral environment and which have relations with adverse effects, the dimethylaminoethyl methacrylate (DMAEMA) and its degradation products, such as dimethylethanolamine, are vasoactive and induce relaxation and contraction of blood vessels, exerting an important role in homeostasis [47]. Methyl methacrylate, the main component of acrylic resins, is associated with dermal sensitization and respiratory irritation; and besides this substance, the formaldehyde is also released into the environment, mainly from autopolymerized resins [43,48].

The biocompatibility of different types of denture base resins has been evaluated over time, in an attempt to define the best operating parameters for each material [49, 50]. In an attempt to reduce aggression to the tissues of the oral cavity, the hypoallergenic resins were developed. These materials have lower amounts of residual methyl methacrylate than conventional heat-polymerized acrylic resins [51]. Trials show that hypoallergenic resins have smaller sorption and solubility values than conventional acrylic resins, which is a significant advantage of these materials since these properties influence on the dimensional stability of the material, among other [52]. However, the poor availability of information on their mechanical properties does not give security to be definitely used in place of conventional polymethylmethacrylate.

It is well known that the handling characteristics have fundamental influence on the biological properties of base material. The polymer/monomer ratio is one of the variables that significantly influence the cytotoxicity of acrylic resins, and the more monomer added to the mixture, the greater the amount of its residual effect and cytotoxicity. However, the correct establishment of this relationship which should be proposed by the material manufacturer also takes into account the conditions of clinical work [53].

Polymerization method is decisive in the cytotoxicity of polymeric materials, since, from them, result in different residual monomer concentrations [53]. The literature shows that chemically-activated resins have lower biological compatibility than microwave-activated and heat-polymerized resins, since they have lower conversion ratio of monomers into polymers and, consequently, greater amount of residual monomer [54-56].

Depending on the characteristics of the polymerization cycle, such as temperature and time, the different amounts of residual monomer are maintained in the polymer, resulting in various degrees of cytotoxicity. Thus, in general, a longer polymerization cycle results in a lower cytotoxic effect [53,57].

Given the above, various post-polymerization treatments have been evaluated and recommended in order to minimize the cytotoxicity reactions, such as those that aim primarily to increase the degree of polymer conversion [44,58-61].

Additional microwave polymerization cycles may provide a reduction of the residual monomer content of up to 25% [58], are technically easy and quick to run, and produce little heat [59]. Due to the high conversion of monomer to polymer, these cycles potentiates the mechanical strength of these materials [44,58,59,60].

Water-bath post-polymerization also contribute to the reduction of cytotoxicity of denture base and reline acrylic resins, since in this early period occurs the biggest release of toxic components [44]. Besides the significant reduction of the amount of residual monomer, this treatment increases the hardness of the rigid reline resins [62]. Several protocols have been devised for this purpose with success, such as the permanence of the prosthesis for 10 minutes at 55°C [62] and 24/48 hours at 37°C [54].

Reduction of Prosthesis Contamination by the Addition of Antimicrobial Substances

Complete dentures act as microbial niches [63-65] and may accumulate respiratory pathogens that favor the onset of pneumonia; this disease is the largest geriatric problem [66,67]. It is possible that complete dentures collaborate to re-infection of their users, especially in cases which the hygiene of the prosthesis is not performed properly. This condition becomes alarming when considered that many patients with immune deficiency are users of these prosthetic devices [64]. The accumulation of microorganisms in dentures is favored even by the own characteristics of acrylic resin whose porosity, roughness and sorption facilitate the deposition of biofilm [64,68].

Mechanical and chemical methods are traditionally employed in order to remove the biofilm of dentures [69]. The former are essential for microbial displacement; however, there is evidence that this action alone is not sufficient for a perfect hygiene, especially for elderly

patients with reduced motor skills, being necessary its association with chemical methods, among which the antimicrobials are highlighted [70,71].

Currently, the application of antimicrobial agents in the complete dentures can be accomplished in the form of solutions for immersion, either conventional or specific to prostheses toothpastes [72,73]. However, in search of greater effectiveness and functionality, other forms of the use of antimicrobials have been evaluated in scientific research.

For the convenience provided to users, the incorporation of these agents on the materials by manufacturers is presented as a current trend [64,74,75]. With this method, regardless of the antimicrobial agent selected, it is possible to manipulate the variables so that the release is controlled and occurs over time, which is desirable due to the longevity of the accomplished works. This condition would be also favorable for acrylic resins for the reasons given above. However, the feasibility of the inclusion of antimicrobials in dental materials also depends on the preservation of their physical, mechanical and biological properties. Unfortunately, these incorporations have resulted not only in benefits but also in significant changes in these properties [76].

Among the materials tested as antimicrobial additives of acrylic resins, those based on silver, which act on cell walls of microorganism, interrupt the replication process of its RNA and block the cellular respiration of the microorganism are highlighted [77]. In addition, the active silver has low toxicity to human cells and broad antimicrobial activity, which explains its widespread use at this time [78].

In the last years, products impregnated with silver which has been more incorporated into denture base materials are zeolites [64,79,80]. These crystalline structures of aluminosilicate have thermal resistance and do not suffer disintegration even at high temperatures, enabling the success of its association with heat-polymerized materials [64].

The effects of this association have revealed that the viscoelastic properties of tissue conditioners are not changed compared to those presented at the pure materials [79]. Regarding the toxicology, which is essential due to the constant contact between the tissue conditioner and the oral mucosa, the addition of silver zeolite does not affect the cell viability when compared to the material without antimicrobial agents [80]. A scientific study showed that the incorporation of 2.5% of silver- and zinc-containing zeolite potentiates the action of heat-polymerized acrylic resins activated by microwave energy against *Candida albicans* and *Streptococcus mutans*; however, it reduces negatively the flexural strength and the impact of these materials [64]. Among the possible explanations for the reduction of resistance values of the resins which in general contain antimicrobial agents are the reduction of the degree of conversion and the consequent increase in the amount of residual monomer [81], dilution of components in the liquid and the molecular interaction [82].

Quaternary ammonium at different percentages (2-50%) has also been added to heat-polymerized acrylic resins in an attempt to reduce its contamination by different microorganisms, and it was observed the action of the compost formed for up to four weeks [83]. Nevertheless, the reduction of the hardness and elastic modulus of a cold-cured acrylic resin added of 10% w/w of chlorhexidine acetate was again reported by researchers [76].

Another attempt made to reduce the contamination of denture resins occurred through the coverage of these materials with titanium dioxide, that make the adhesion of *Candida albicans* and *Streptococcus sanguinis* more difficult [84].

Monomers have also been used for the purpose of conferring the antimicrobial action in resins for denture base. The effect of adding more than 1.75% of 2-tert-butylaminoethyl

methacrylate in these materials decreased their flexural strength values, possibly by causing the softening of resin. Clinically, this incorporation could result in the increase of the incidence of fractures in complete dentures bases [85].

The microbial adhesion to the resins can also be reduced by the incorporation of polar radicals, such as methacrylic acid, with the aim of increasing its hydrophilic. It was observed a significantly reduction of *Candida albicans* adhesion compared to control group (resin without treatment) [86].

Permanently there is an attempt to incorporate antifungal agents such as nystatin to tissue conditioners. It was observed that the addition of concentrations of 10% of the agent confers action against micro-organisms and preserves the mechanical properties of materials [87].

Surface treatments have proven to be promising alternatives to reduce the microbial adhesion to denture base materials. For example, the application of plasma on the surface of acrylic resin to change its hydrophobicity results in the reduction of the adhesion of *Candida albicans* to the material. However, this treatment does not have limitations resulting from the incorporation of antimicrobial agents in these materials [88], as already showed. Another proposed surface treatment was the application of a layer of light-cured glaze on the surface of acrylic resins; nevertheless, this protection was offered only for a limited period of time when its integrity was maintained [89].

Thus, the search for new materials or treatments that enable the reduction of contamination, which not cause adverse effects to oral tissues and have appropriate physical and mechanical properties should still continue. What should it have done while the problem is not solved is to control the variables of laboratory manipulation to reduce the porosity of resins, such as providing adequate monomer/polymer relation, perform the pressing and polymerization following the manufacturers' recommendations, and perform prosthesis polishing of good quality.

Conclusion

For better clinical performance of the complete dentures, it is suggested that clinical dentists and dental prosthesis technicians seek to manipulate the dental materials according to parameters set by their manufacturers. Additionally, the constant updating of these professionals should occur, since new materials and working techniques have been present often in literature.

References

- [1] Lefebvre, CA; Knoernschild, KL; Schuster, GS. Cytotoxicity of eluates from light-polymerized denture base resins. *Journal of Prosthetic Dentistry*, 1994 72, 644-650.
- [2] Hayden, WJL. Flexural strength of microwave-cured denture baseplates. *General Dentistry*, 1986 34, 367-371.
- [3] Vallittu, PK; Ruyter, IE; Buykuilmaz, S. Effect of polymerization temperature and time on the residual monomer content of denture base polymers. *European Journal of Oral Sciences*, 1998 106, 588-593.

-
- [4] Jerolimov, V; Huggett, R; Brooks, CS; Bates, JF. The effect of the variations in the polymer/monomer mixing ratios on residual monomer levels and flexural properties of denture base materials. *Quintessence International*, 1985 9, 431-434.
 - [5] Hayden, WJL. Flexural strenght of microwave-cured denture baseplates. *General Dentistry*, 1986 34, 367-371.
 - [6] Sato, S; Cavalcante, MRS; Orsi, IA; Zaniquelli, O. Assessment of flexural strength and color alteration of heat-polymerized acrylic resins after simulated use of denture cleansers. *Brazilian Dental Journal*, 2005 16, 124-128.
 - [7] Kanie, T; Kadokawa, A; Arikawa, H; Fujii, K; Ban, S. Flexural properties of ethyl or methyl methacrylate-UDMA blend polymers. *Dental Materials Journal*, 2010 29, 575-581.
 - [8] Takahashi, Y; Chai, J; Kawaguchi, M. Effect of water sorption on the resistance to plastic deformation of a denture base material relined with four different denture reline materials. *International Journal of Prosthodontics*, 1998 11, 49-54.
 - [9] Arima, T; Murata, H; Hamada, T. Properties of highly crosslinked auto polymerizing reline acrylic resin. *Journal of Prosthetic Dentistry*, 1995 73, 55-59.
 - [10] Baker, S; Brooks, SC; Walker, DM. The release of residual monomeric methyl methacrylate from acrylic appliances in the human mouth: an assay for monomer in saliva. *Journal of Dental Research*, 1988 67, 1295-1299.
 - [11] John, J; Gangadhar, SA; Shah, I. Flexural strength of heat-polymerized polymethyl methacrylate denture resin reinforced with glass, aramid, or nylon fibers. *Journal of Prosthetic Dentistry*, 2001 86, 424-427.
 - [12] Kostoulas, I; Kavoura, VT; Frangou, MJ; Polyzois, GL. The effect of length parameter on the repair strength of acrylic resin using fibers or metal wires. *General Dentistry*, 2008 56, 51-55.
 - [13] Takahashi, Y; Yoshida, K; Shimizu, H. Effect of location of glass fiber-reinforced composite reinforcement on the flexural properties of a maxillary complete denture *in vitro*. *Acta Odontologica Scandinavica*, 2011 69, 215-221.
 - [14] Hedzelek, W; Gajdus, P. Mechanical strength of an acrylic resin palatal denture base reinforced with a mesh or bundle of glass fibers. *International Journal of Prosthodontics*, 2007 20, 311-312.
 - [15] Yun, SH; Cho, D; Kim, J; Lim, S; Lee, GW; Park, M; Lee, SS. Effect of silane coupling agents with different organo-functional groups on the interfacial shear strength of glass fiber/Nylon 6 composites. *Journal of Materials Science Letters*, 2003 22, 1591-1594.
 - [16] Hamza, TA; Rosentiel, SF; Elhosary, MM; Ibraheem, RM. The effect of fiber reinforcement on the fracture toughness and flexural strength of provisional restorative resins. *Journal of Prosthetic Dentistry*, 2004 91, 258-264.
 - [17] Keyf, F; Uzun, G. The effects of glass fiber reinforcement at different concentrations on the transverse strength, deflection and modulus of elasticity of a provisional fixed partial denture resin. *Journal of Biomaterials Applications*, 2001 16, 149-156.
 - [18] Vallitu, PK. Flexural properties of acrylic resin polymers reinforced with unidirectional and woven glass fibers. *Journal of Prosthetic Dentistry*, 1999 81, 318-326.
 - [19] Goldberg, AJ; Freilich, MA. An innovative pre-impregnated glass fiber for reinforcing composites. *Dental Clinics of North America*, 1999 43, 127-133.

- [20] Vallittu, PK. Prosthodontic treatment with glass fiber reinforced composite resin bonded fixed partial denture. *Journal of Prosthetic Dentistry*, 1999 82, 132-135.
- [21] Bertassoni, LE; Marshall, GW; de Souza, EM; Rached, RN. Effect of pre- and postpolymerization on flexural strength and elastic modulus of impregnated, fiber-reinforced denture base acrylic resins. *Journal of Prosthetic Dentistry*, 2008 100, 449-457.
- [22] Goguță, LM; Bratu, D; Jivănescu, A; Erimescu, R; Mărcăuțeanu, C. Glass fibre reinforced acrylic resin complete dentures: a 5-year clinical study. *Gerodontology*, 2011 22, *in press*.
- [23] Aydan, C; Yilmaz, H; Caglar, A. Effect of glass fiber reinforced on the flexural strength of different denture base resins. *Quintessence International*, 2002 33, 457-463.
- [24] Meiers, JC; Duncan, JP; Freilich, MA; Goldberg, AJ. Preimpregnated, fiber-reinforced prostheses. Part II. Direct applications: splints and fixed partial dentures. *Quintessence International*, 1998 29, 761-768.
- [25] Vallittu, PK. Glass fiber reinforcement in repaired acrylic resin removable dentures: Preliminary results of a clinical study. *Quintessence International*, 1997 28, 39-44.
- [26] Cheng, YY; Li, JY; Fok, SL; Cheung, WL; Chow, TW. 3D FEA of high-performance polyethylene fiber reinforced maxillary dentures. *Dental Materials*, 2010 26, 211-219.
- [27] Stipho, HD. Effect of glass fiber reinforcement on some mechanical properties of autopolymerizing polymethyl methacrylate. *Journal of Prosthetic Dentistry*, 1998 79, 580-584.
- [28] Goiato, MC; Pesqueira, AA; Vedovatto, E; Santos, DM; Gennari Filho, H. Effect of different repair techniques on the accuracy of repositioning the fractured denture base. *Gerodontology*, 2009 26, 237-241.
- [29] Polyzois, GL; Andreopoulos, AG; Lagouvardos, PE. Acrylic resin denture repair with adhesive resin and metal wires: effects on strength parameters. *Journal of Prosthetic Dentistry*, 1996 75, 381-383.
- [30] Ward, JE; Moon, PC; Levine, RA; Behrendt, CL. Effect of repair surface design, repair material, and processing method on the transverse strength of repaired acrylic denture resin. *Journal of Prosthetic Dentistry*, 1992 67, 815-820.
- [31] Arioli-Filho, JN; Butignon, LE; Pereira, RP; Lucas, MG; Mollo-Junior, FA. Flexural strength of acrylic resin repairs processed by different methods: water bath, microwave energy and chemical polymerization. *Journal of Applied Oral Science*, 2011 19, 249-253.
- [32] Kostoulas, I; Kavoura, VT; Frangou, MJ; Polyzois, GL. Fracture force, deflection, and toughness of acrylic denture repairs involving glass fiber reinforcement. *Journal of Prosthodontics*, 2008 17, 257-261.
- [33] Sarac, YS; Sarac, D; Kulunk, T; Kulunk, S. The effect of chemical surface treatments of different denture base resins on the shear bond strength of denture repair. *Journal of Prosthetic Dentistry*, 2005 94, 259-266.
- [34] Takahashi, Y; Chai, J; Takahashi, T; Habu, T. Bond strength of denture teeth to denture base resins. *International Journal of Prosthodontics*, 2000 13, 59-65.
- [35] Korkmaz, T; Dogan, A; Dogan, OM; Demir, H. The bond strength of a highly cross-linked denture tooth to denture base polymers: a comparative study. *Journal of Adhesive Dentistry*, 2011 13, 85-92.

-
- [36] Fletcher-Stark, ML; Chung, KH; Rubenstein, JE; Raigrodski, AJ; Mancl, LA. Shear bond strength of denture teeth to heat- and light-polymerized denture base resin. *Journal of Prosthodontics*, 2011 20, 52-59.
- [37] Consani, RL; Carmignani, MR; Mesquita, MF; Correr-Sobrinho, L; Guiraldo, RD. Effect of microwave treatment on the shear bond strength of different types of commercial teeth to acrylic resin. *Gerodontology*, 2010 27, 236-242.
- [38] Barbosa, DB; Barão, VA; Monteiro, DR; Compagnoni, MA; Marra, J. Bond strength of denture teeth to acrylic resin: effect of thermocycling and polymerisation methods. *Gerodontology*, 2008 25, 237-244.
- [39] Dalal, A; Juszczuk, AS; Radford, DR; Clark, RK. Effect of curing cycle on the tensile strength of the bond between heat cured denture base acrylic resin and acrylic resin denture teeth. *European Journal of Prosthodontic and Restorative Dentistry*, 2009 17, 146-149.
- [40] Cunningham, JL; Benington, IC. An investigation of the variables which may affect the bond between plastic teeth and denture base resin. *Journal of Dentistry*, 1999 27, 129-135.
- [41] Patil, R; Juszczuk, AS; Radford, DR; Clark, RK. Is the bond between acrylic resin denture teeth and denture base resin stronger if they are both made by the same manufacturer? *European Journal of Prosthodontic and Restorative Dentistry*, 2010 18, 39-41.
- [42] Kalipçılar, B; Karaagaçlıoğlu, L; Hasanreisoglu, U. Evaluation of the level of residual monomer in acrylic denture base materials having different polymerization properties. *Journal of Oral Rehabilitation*, 1991 18, 399-401.
- [43] Tsuchiya, H; Hoshino, Y; Tajima, K; Takagi, N. Leaching and cytotoxicity of formaldehyde and methyl methacrylate from acrylic resin denture base materials. *Journal of Prosthetic Dentistry*, 1994 71, 618-624.
- [44] Jorge, JH; Giampaolo, ET; Machado, AL; Vergani, CE. Cytotoxicity of denture base acrylic resins: A literature review. *Journal of Prosthetic Dentistry*, 2003 90, 190-193.
- [45] Weaver, RE; Goebel, WM. Reactions to acrylic resin dental prostheses. *Journal of Prosthetic Dentistry*, 1980 43, 138-142.
- [46] Scott, A; Egner, W; Gawkrödger, DJ; Hatton, PV; Sheriff, M; Van Noort, R; Yeoman, C; Ggrummitt, J. The national survey of adverse reactions to dental materials in the UK: a preliminary study by the UK Adverse Reactions Reporting Project. *British Dental Journal*, 2004 196, 471-477.
- [47] Abebe, W; Maddux, WF; Schuster, GS; Lewis, JB. Vascular responsiveness to dimethylaminoethyl methacrylate and its degradation products. *Journal of Biomedical Materials Research Part A*, 2003 66, 155-161.
- [48] Borak, J; Fields, C; Andrews, LS; Pemberton, MA. Methyl methacrylate and respiratory sensitization: A Critical review. *Critical Reviews in Toxicology*, 2011 41, 230-268.
- [49] Jorge, JH; Giampaolo, ET; Vergani, CE; Machado, AL; Pavarina, AC; Carlos, IZ. Biocompatibility of denture base acrylic resins evaluated in culture of L929 cells. Effect of polymerization cycle and post-polymerization treatments. *Gerodontology*, 2007 24, 52-57.

- [50] Barron, DJ; Schuster, GS; Caughman, GB; Lefebvre, CA. Biocompatibility of visible light-polymerized denture base resins. *International Journal of Prosthodontics*, 1993 6, 495-501.
- [51] Pfeiffer, P; Rolleke, C; Sherif, L. Flexural strength and moduli of hypoallergenic denture base materials. *Journal of Prosthetic Dentistry*, 2005 93, 372-372.
- [52] Pfeiffer, P; Rosenbauer, EU. Residual methyl methacrylate monomer, water sorption, and water solubility of hypoallergenic denture base materials. *Journal of Prosthetic Dentistry*, 2004 92, 72-78.
- [53] Kedjarune, U; Charoenworarluk, N; Koontongkaew, S. Release of methyl methacrylate from heat-cured and autopolymerized resins: cytotoxicity testing related to residual monomer. *Australian Dental Journal*, 1999 44, 25-30.
- [54] Sheridan, PJ; Koka, S; Ewoldsen, NO; Lefebvre, CA; Lavin, MT. Cytotoxicity of denture base resins. *International Journal of Prosthodontics*, 1997 10, 73-77.
- [55] Bartoloni, JA; Murchison, DF; Wofford, DT; Sarkar, NK. Degree of conversion in denture base materials for varied polymerization techniques. *Journal of Oral Rehabilitation*, 2000 27, 788-793.
- [56] Cimpan, MR; Cressey, LI; Skaug, N; Halstensen, A; Lie, SA; Gjertsen, BT; Matre, R. Patterns of cell death induced by eluates from denture base acrylic resins in U-937 human monoblastoid cells. *European Journal of Oral Sciences*, 2000 108, 59-69.
- [57] Harrison, A; Huggett, R. Effect of the curing cycle on residual monomer levels of acrylic resin denture base polymers. *Journal of Dentistry*, 1992 20, 370-374.
- [58] Balgojevic, V; Murphy, VM. Microwave polymerization of denture base materials. A comparative study. *Journal of Oral Rehabilitation*, 1999 26, 804-808.
- [59] Patil, PS; Chowdhary, R; Mandokar, RB. Effect of microwave postpolymerization treatment on residual monomer content and the flexural strength of autopolymerizing reline resin. *Indian Journal of Dental Research*, 2009 20, 293-297.
- [60] Azzari, MJ; Cortizo, MS; Alessandrini, JL. Effect of the curing condition on the properties of an acrylic denture base resin microwave polymerized. *Journal of Dentistry*, 2003 31, 463-468.
- [61] Mendonça, MJ; Machado, AL; Giampaolo, ET; Pavarina, AC; Vergani, CE. Weight loss and surface roughness of hard chairside reline resins after toothbrushing: influence of postpolymerization treatments. *International Journal of Prosthodontics*, 2006 19, 281-287.
- [62] Urban, VM; Machado, AL; Vergani, CE; Giampaolo, ET; Pavarina, AC; de Almeida, FG; Cass, QB. Effect of water-bath post-polymerization on the mechanical properties, degree of conversion, and leaching of residual compounds of hard chairside reline resins. *Dental Materials*, 2009 25, 662-671.
- [63] Compagnoni, MA; Souza, RF; Marra, J; Pero, AC; Barbosa, DB. Relationship between *Candida* and nocturnal denture wear: quantitative study. *Journal of Oral Rehabilitation*, 2007 34, 600-605.
- [64] Casemiro, LA; Martins, CHG; Pires-de-Souza, FCP; Panzeri, H. Antimicrobial and mechanical properties of acrylic resins with incorporated silver-zinc zeolite - part I. *Gerodontology*, 2008 25, 187-194.
- [65] He, XY; Meurman, JH; Kari, K; Rautemaa, R; Samaranayake, LP. *In vitro* adhesion of *Candida* species to denture base materials. *Mycoses*, 2006 49, 80-84.

-
- [66] Sumi, Y; Kagami, H; Ohtsuka, Y; Kakinoki, Y; Haruguchi, Y; Miyamoto, H. High correlation between the bacterial species in denture plaque and pharyngeal microflora. *Gerontology*, 2003 20, 84-87.
- [67] Senpuku, H; Sogame, A; Inoshita, E; Tsuha, Y; Miyazaki, H; Hanada, N. Systemic diseases in association with microbial species in oral biofilm from elderly requiring cases. *Gerontology*, 2003 49, 301-309.
- [68] Verran, J; Maryan, JC. Retention of *Candida albicans* on acrylic resin and silicone of different surface topography. *Journal of Prosthetic Dentistry*, 1997 77, 142-143.
- [69] Paranhos, HF; Panzeri, H; Lara, EH; Candido, RC; Ito, IY. Capacity of denture plaque removal and antimicrobial action of a specific paste formulated for denture cleaning. *Brazilian Dental Journal*, 2000 11, 97-104.
- [70] Abelson, DC. Denture plaque and denture cleansers: review of the literature. *Gerodontology*, 1985 1, 202-206.
- [71] Dills, SS; Olshan, AM; Goldner, S; Brogdon, C. Comparison of the antimicrobial capability of an abrasive paste and chemical-soak denture cleaners. *Journal of Prosthetic Dentistry*, 1988 60, 467-470.
- [72] Shay, K. Dental Hygiene: A review and Update. *Journal of Contemporary Dental Practice*, 2000 1, 28-41.
- [73] Pavarina, AC; Pizzollitto, AC; Machado, AL; Vergani, CE; Giampaolo, ET. An Infection control protocol: effectiveness of immersion solutions to reduce the microbial growth on dental prostheses. *Journal of Oral Rehabilitation*, 2003 30, 532-536.
- [74] Hotta, M; Nakajima, H; Yahamoto, K; Aono, M. Antibacterial temporaty filling materials: the effect of adding various ratios of Ag-Zn-Zeolite. *Journal of Oral Rehabilitation*, 1998 25, 485-489.
- [75] Thom, DC; Davies, JE; Santerre, JP; Friedman, S. The hemolytic and cytotoxic properties of a zeolite-containing root filling material *in vivo*. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology Endodontics*, 2003 95, 101-108.
- [76] Thaw, M; Addy, M; Handley, R. The effects of drug and water incorporation upon some physical properties of cold cured acrylic. *Journal of Biomedical Materials Research*, 1981 15, 29-36.
- [77] Uchida, T; Maru, N; Furuhashi, M; Fujino, A; Muramoto, S; Ishibashi, A; Koshiba, K; Shiba, T; Kikuchi, T. Anti-bacterial zeolite balloon catheter and its potential for urinary tract infection control. *Hinyokika Kyo*, 1992 38, 973-978.
- [78] Williams, RL; Doherty, PJ; Vince, DG; Grashoff, GJ; Williams, DF. The biocompatibility of silver. *Critical Reviews of Biocompatibility*, 1989 5, 221-223.
- [79] Ueshige, M; Abe, Y; Sato, Y; Tsuga, K; Akagawa, Y; Ishii, M. Dynamic viscoelastic properties one antimicrobial tissue conditioners containing silver-zeolite. *Journal of Dentistry*, 1999 27, 517-522.
- [80] Abe, Y; Ueshige, M; Takeushi, M; Ishii, M; Akagawa, Y. Citotoxicity of antimicrobial tissue conditioners containing silver-zeolite. *International Journal of Prosthodontics*, 2003 16, 141-144.
- [81] Shibata, T; Hamada, N; Kimoto, K; Sawada, T; Sawada, T; Kumada, H; Umemoto, T; Toyoda, M. Antifungal effect of acrylic resin containing Apatite-coated TiO₂ photocatalyst. *Dental Materials Journal*, 2007 26, 437-444.

- [82] Cunha, TR; Regis, RR; Bonatti, MR; Souza, RF. Influence of incorporation of fluoroalkyl methacrylates on roughness and flexural strength of a denture base acrylic resin. *Journal of Applied Oral Science*, 2009 17, 103-107.
- [83] Pesci-Bardon, C; Fosse, T; Madinier, I; Serre D. *In vitro* new dialysis protocol to assay the antiseptic properties of a quaternary ammonium compound polymerized with denture acrylic resin. *Letters of Applied Microbiology*, 2004 39, 226-231.
- [84] Arai, T; Ueda, T; Sugiyama, T; Sakurai, K. Inhibiting microbial adhesion to denture base acrylic resin by titanium dioxide coating. *Journal of Oral Rehabilitation*, 2009 36, 902-908.
- [85] Paleari, AG; Marra, J; Pero, AC; Rodriguez, LS; Ruvolo-Filho, A; Compagnoni, MA. Effect of incorporation of 2-tert-butylaminoethyl methacrylate on flexural strength of a denture base acrylic resin. *Journal of Applied Oral Sciences*, 2011 19, 195-199.
- [86] Park, SE; Blissett, R; Susarla, SM; Weber, HP. *Candida albicans* adherence to surface-modified denture resin surfaces. *Journal of Prosthodontics*, 2008 17, 365-369.
- [87] El-Charkawi, H; el-Said, EA; Safouh, HM; el-Raghi, N. Effect of addition antimicrobial agents to denture reliners. *Egyptian Dental Journal*, 1994 40, 785-790.
- [88] Zamperini, CA; Machado, AL; Vergani, CE; Pavarina, AC; Giampaolo, ET; da Cruz, NC. Adherence in vitro of *Candida albicans* to plasma treated acrylic resin. Effect of plasma parameters, surface roughness and salivary pellicle. *Archives of Oral Biology*, 2010 55, 763-770.
- [89] Sesma, N; Lagana, DC; Morimoto, S; Gil, C. Effect of Denture Surface Glazing on Denture Plaque Formation. *Brazilian Dental Journal*, 2005 16, 129-134.

Chapter VIII

Alternatives to Removable Complete Dentures: Viability for Public Health Systems in Developing Countries

*Mara Rodrigues Martins³, Wellington Márcio dos Santos Rocha¹,
Débora Emrich Magalhães³,
Fernanda Piana Santos Lima de Oliveira³, Virgínia Sanábio Vilela³,
Marcos Azeredo Furquim Werneck²
and Efigênia Ferreira e Ferreira^{2*}*

¹Department of Restorative Dentistry,

²Department of Public Health,

Universidade Federal de Minas Gerais (UFMG), Dentistry School, Brazil

³Private clinics, Brazil

Abstract

The global population is ageing at an unprecedented rate. The result of this process has produced an important change in the healthcare system, with the cure of acute diseases giving way to the treatment of chronic illnesses. The impact of this demographic change is of considerable concern due to the increase in the cost of healthcare systems, although the current correlation between health costs and the elderly population is weak. In the last ten years, Brazil has made considerable progress in the prevention and control of caries among children. However, the situation among adults and the elderly is among the poorest in the world. However, the rehabilitation of these individuals is hindered by a number of factors. The difficulty of retaining and adapting lower dentures is common and constitutes an obstacle to the complete success of prosthesis use, as reported in a number of studies. Moreover, the lack of information on the part of patients regarding the limitations of conventional complete dentures contributes toward dissatisfaction and the

* E-mail address: efigeniaf@gmail.com.

abandonment of complete lower denture use. The aim of this chapter is to carry out a systematic review of methods for the rehabilitation of individuals with an edentulous lower jaw in the search for an alternative solution that is biological, functional and financially viable. This review will consist of the treatment and options for edentulismo including the bone loss risk, the patient satisfaction, the masticatory efficiency, the number of implants and the economic cost/analysis. At the final presents a discussion on clinical decisions in the edentulous rehabilitation.

Introduction

The demographic shift that is taking place around the world, as a result of an aging population, has been attributed to increased life expectancy, birth control, and improvements made in the social and health areas (Lebrão, 2007). As a result of this phenomenon, there has also been a major change in the health care system that, today, confronts a growing ability to cure acute diseases and treat chronic illnesses (Bury, 2001; Turkyilmaz et al., 2010). Although the correlation between current spending on health care and the elderly population is weak (Niessen, 2000; Anderson and Hussey, 2000), the impact of this shift has caused great concern because of the possibility of rising costs in the health care systems.

Furthermore, the search for a better quality of life for this population group needs to focus on healthy aging that includes a low risk of contracting diseases and their resulting functional disabilities, a high degree of mental and physical functioning, and an active involvement with life (Cupertino et al., 2007). In this context, health is considered not as a life goal, but as a resource for life itself (WHO, 1986).

Oral health encompasses a range of biological and psychological conditions, which enables the human being to carry out functions such as chewing, swallowing, and talking. It also comprises an aesthetic dimension, which plays a role in self-esteem and social relationships, without inhibition or embarrassment. It is a health field that, when problems arise, inevitably affects the quality of a person's life. However, the current state of oral health, especially in adults and the elderly in developing countries, is still a long way from facilitating healthy aging. In fact, the model of care focused on healing and disabling practices was one of the determinants of this panorama (Moreira et al., 2005; Vargas and Paixão, 2005). A high degree of edentulism has a negative impact on the quality of life, particularly in terms of the psychological distress caused by the difficulty of being accepted by others and a sense of humiliation that a toothless person often experiences (Silva et al., 2010/b). The complete loss of teeth results in negative consequences such as shame, difficulty in eating, impairment of social relationships, and a sense of incompleteness (Silva et al., 2010/a).

Using Brazil as an example, in recent decades, there has been a major advance in the prevention and control of dental caries in children and adolescents. However, the situation in adults and the elderly is among the worst in the world. According to the National Survey of Oral Health sponsored by the Ministry of Health, three million people between the ages of 65 and 74 require complete dentures in both arches (15%) and 4 million are in need of a complete denture in one of the arches (23%) (Roncalli, 2011). Edentation has become a public health problem, engendering a growing demand in the population and need for

prosthetic treatment (Colussi and Freitas, 2002, Moreira et al., 2005; Vargas and Paixão, 2005; Campostrini et al., 2007; Ferreira et al., 2007).

Facing the possibility of the prosthetic replacement of teeth triggers anxiety in many people, but the thought of returning to a generally accepted social standard and regaining their self image motivates them (Silva et al., 2010/a). The benefit of using a prosthesis is undermined when the person fails to achieve sufficient stability for chewing, especially in the mandibles. The resulting dissatisfaction often leads the user to abandon the prosthesis, as has been frequently observed in the case of removable dentures (De Souza e Silva et al., 2009).

Consequently, the search for alternatives, which could be implemented with safe protocols, especially in public services, is being driven by need and urgency.

Treatment Options for Edentulism

Edentulism associated with poor access to treatment and its limitations produces functional and psychosocial disorders that tend to segregate the individual and burden the health care system.

The successful rehabilitation of this patient by means of prosthesis should result in an improved functional and aesthetic gain in the quality of life. The prosthetic options for this purpose include: conventional dentures (CD), the fixed implant-supported prosthesis (FISP), and the implant-retained overdenture (IRO).

Conventional Denture

A conventional denture (CD) consists of a removable denture, made of acrylic resin and on the mucosa. It is the most widely used therapeutic modality for the rehabilitation of edentulous patients, having been accepted until recently as the prosthesis of choice for this purpose. With its low cost and technical simplicity, it was an unprecedented success, rehabilitating millions of people around the world (Raghoobar, et al., 2003; Turkyilmaz et al., 2010).

The fact of relying on the mucosa for support causes problems for the patient, such as increased bone resorption, adjustment difficulties, pain, discomfort, low stability and retention of the prosthesis, and struggling while eating, especially as related to the lower dentures (Raghoobar, et al., 2003; Allen and McMillan, 2003). In addition, the reduced salivary flow, motor abnormalities of the tongue, reduction in bite force, and decreased sensory function contribute to its functional loss (Wolff et al., 2003; Allen and McMillan, 2003; Ikebe et al., 2006).

The weakened act of chewing can lead to the development of disease and lead to a reduced expectation and quality of life (Slagter et al., 1993; Slagter et al., 1992; Shinkai, et al., 2002; Emami, et al., 2009). Patient disinformation with regard to the limitations of this type of prosthesis also contributes to the increased rate of dissatisfaction and the abandonment of the use of the lower denture (Vervoorn, et al., 1988; Raghoobar et al., 2003; Silva et al., 2010/a).

Patient satisfaction is the yardstick in the professional's clinical planning. The satisfied patient does not come looking for treatment often or return to have the existing prosthesis replaced. However, the dissatisfied patient, whether due to the inefficiency of their current prosthesis or emotional problems, needs alternatives that facilitate his/her oral rehabilitation.

Observation and confirmation of the osseointegration phenomenon by Brånemark in the mid-1960s provided a new rehabilitative dimension for these edentulous individuals. The use of osseointegrated implants in their different ways suggests and encourages the conclusion that the patient's quality of life is significantly enhanced by prosthesis over implant (Heydenrijk, et al., 2002; Zarb and Bolender, 2006).

Fixed Implant-Supported Prosthesis

The fixed implant-supported prosthesis (FISP) originated with the Brånemark protocol, which was the first use of implants in the treatment of the edentulous mandible and consisted of a fixed denture on implants in 4 to 6 regions between the mental foramen. It is characterized by a metal infrastructure, in resin, associated with the teeth and gums, and a bilateral distal cantilever that should not exceed two teeth (15mm) screwed into the implants. The surgery was performed in two stages (installation and re-opening of the implants) with a spacing of 3 to 4 months between the stages, at which time the prosthetic (loading) phase was begun (Brånemark et al., 1969 and 1977).

The success of these prostheses in the rehabilitation of edentulous mandibles was confirmed by longitudinal studies (Brånemark et al., 1977; Adell, et al., 1981; Albrektsson et al., 1987).

Subsequently, changes took place in the Brånemark protocol due to patient-related factors. These changes deal with the number and location of implants, type of prosthesis fixation (cemented and screwed), and load (conventional, early, and immediate).

The impossibility of placing the implants in suitable number and arrangement, with or without economic reasons, led to the resurgence of overdentures (removable dentures), which are now on implants and no longer only by the residual roots (Naert et al., 1997; Awad et al., 2003).

Implant Retained Overdenture

The implant-retained overdenture (IRO) is an acrylic resin removable denture, retained by implants, and may or may not be on mucosa. The number and location of the implants, along with the various types of attachments, determine the variations in this type of prosthesis. According to Fitzpatrick (2006), this is a less invasive, cheaper, and equally effective treatment option when compared to the fixed implant-supported prosthesis.

The choice between an implant-retained overdenture and a fixed implant-supported prosthesis as an alternative to conventional dentures in edentulous mandibles suggests some questions, such as implant survival, bone loss, patient satisfaction and quality of life as related oral health, chewing efficiency, the number of implants, and cost, all of which will guide the decision (see also below).

Implant Survival and Bone Loss

Implant Survival

Bryant et al., (2007), suggest, in a systematic review that included meta-analysis, that the survival of implants in the fixed mandibular implant-supported prostheses is 6.6% higher than in fixed maxillary implant-supported prostheses. They conclude that the implant's survival and success are not affected by prosthesis type.

Bone Loss

With regard to bone loss, the first aspect to be discussed is the relationship between peri-implant bone loss and the different types of loading and prostheses. Sunyoung and Payne (2010) cite three types of loading for implant-retained overdentures on two implants: the conventional load, in which the prosthesis is attached after a second surgical procedure following a healing period ranging from three to four months, early loading, in which the prosthesis, along with the fastening system, is installed 48 hours after the implant, in up to 3 months, and immediate loading, in which the prosthesis, along with the fastening system, is installed up to 48 hours of the implant installation. This study suggests that, in the short term, there are no harmful effects on the marginal bone level using the immediate or early loading protocols, and the long-term results still need to be obtained. They conclude that recommending immediate loading protocols in mandibular overdentures on two implants may still be premature.

Jokstad and Carr (2007) analyzed the influence of loading time on the different types of prostheses on implants and concluded that there is no evidence that early or immediate loading is not a safe procedure.

The success rate of immediate loading (up to 1 week), early (from 1 week to 2 months), and conventional (after 2 months) showed that it is possible to succeed with the different load types, with the degree of the implant's primary stability appearing to be a prerequisite for success with immediate and early loading (Esposito et al., 2007).

Alsabeeha et al., (2010) also suggest that both immediate and early loading protocols using 2, 3, and 4 implants supporting or retaining mandibular overdentures may achieve comparable success to that obtained with the conventional load for more than two years. They suggest, however, that long-term studies are needed to validate the early and immediate loading protocol. The second aspect to be considered in terms of bone loss is the relationship between resorption of the posterior mandibular residual ridge and the different types of prostheses.

Jacobs et al., (1992) related the subsequent bone resorption to the different types of prostheses. In fixed implant-supported prosthesis for 4 to 6 implants, there was minimum resorption, while in implant-retained overdentures on 2 implants, it was found to be from 2 to 3 times higher than in conventional dentures in a six-month period of post-extraction remodeling. In patients who have been edentulous for more than 10 years, the difference in absorption rates for the latter two groups disappeared. The use of implant-retained

overdentures in younger patients should be carefully evaluated from a long-term perspective, although it is an option that offers advantages from a financial standpoint.

Kordatzis et al., (2003) compared conventional dentures and implant-retained overdentures on two implants and found 1.62-mm bone loss in dentures, when compared with 0.69 mm in implant-retained overdentures over a five-year period in patients who had been edentulous for more than twenty-two years.

The degree of residual ridge resorption differed when the number of implants in implant-retained overdentures ranged from 2 to 4, which is 4% lower when a larger number of implants were used in the ten-year control study (Meijer et al., 2010).

Patient Satisfaction

Numerous studies have focused on understanding the impact that different types of treatment for mandibular edentulism have on the patient's life. Despite the conventional denture's known and prevailing limitations, such as instability, poor mastication, decreased self-confidence, quality of life, satisfaction, and social relations, and the positive impact of implant therapy, the satisfaction-based results need to be better understood. Edentulous patients have highly variable, individual functional requirements that are influenced by social, cultural, and financial factors, along with their capacity to adapt (Fitzpatrick, 2006).

Thomason et al., (2007) found strong evidence that the quality of life could be improved through the use of implant-retained overdentures; however, there was no perceivable difference with regard to patient satisfaction. Emami et al., (2009) found evidence showing that overall satisfaction and quality of life associated with oral health are higher in users of implant-retained overdentures when compared with conventional dentures, although the magnitude of these effects remains uncertain. This study suggests that an implant-retained overdenture on two implants should be the minimum offered to mandibular edentulous patients as a first treatment option. Evaluations of satisfaction, stability, comfort, and masticatory efficiency were significantly better in patients treated with implant-retained overdentures on two and four implants when compared with conventional denture users. The authors concluded that both the implant-retained overdentures and the fixed implant-supported prostheses significantly improve the quality of life of edentulous patients when compared with conventional dentures (Turkilmaz et al., 2010).

Masticatory Efficiency

Users' masticatory efficiency with CD in the upper and lower arches is less than 20% when compared with that of the natural tooth (Kapur and Soman, 1964; Heath, 1982). Functional loss is linked to low stability and support, but may also be affected by reduced salivary flow, motor control of the tongue, reduced bite force, and decreased sensory function (Koshino et al., 1997; Wolff et al., 2003; Allen and McMillan, 2003; Ikebe et al., 2006). Many studies relate the reduction in bite force to masticatory muscle atrophy, reduced masticatory ability, and nutritional status (Kapur and Soman, 1964; Wayler and Chauncey, 1983; Slagter et al., 1992; Slagter et al., 1993; Shinkai et al., 2002).

Dentists and patients should be aware of the functional limitations and benefits of implant-retained overdentures and implant-supported bridges when making a well-informed treatment choice (Fueki et al., 2007).

The combination of an implant-retained overdenture or fixed mandibular implant-supported prosthesis with a conventional denture provides significant improvement in masticatory efficiency when compared to the use of dentures in both arches. This applies to conventional mandibular denture users with functional problems attributable to persistent severe resorption (Fueki et al., 2007).

Van der Bilt et al., (2010) evaluated the long-term effects of treatment with implant-retained mandibular overdentures on two implants on oral function. The authors quantified the maximum bite force and masticatory performance 10 years after implant treatment. The study concludes that treatment with implant-retained mandibular overdentures significantly improves oral function and that after 10 years of practice, the maximum bite force and masticatory performance remained unchanged.

The various types of attachments in implant-retained mandibular overdentures have limited impact on masticatory performance (Çehreli et al., 2010).

Number of Implants

In a longitudinal study, Branemark et al., (1995) concluded that the use of four or six 10-mm long implants in the area between the mental foramen is highly likely to support a fixed implant-supported prosthesis. A minimum of four 7-mm long implants is suggested by Lekholm (1998) for the same reason. However, the author recommends that, whenever possible, five implants, preferably longer, be used. The same number of implants has also been suggested for a fixed implant-supported prosthesis (Zarb and Bolender, 2006). To support an implant-retained overdenture, two implants, whether splinted or not, are a generally accepted treatment option (Naert et al., 2004; Carlsson et al., 2004; Meijer et al., 2010). The use of a single implant to support this type of prosthesis has also been employed (Krennmair and Ulm, 2001; Walton et al., 2009; Kronström et al., 2010).

Klemetti (2008) carried out a systematic review to evaluate whether there is an optimal number of implants for retaining implant-retained mandibular overdentures. The results showed that patient satisfaction and prosthesis performance do not seem to be dependent on the number of implants or type of attachment. However, an implant-retained overdenture with two implants and a bar appears to have fewer complications.

Cost and Economic Analysis

The cost greatly influences the choice of prosthesis, especially when it comes to public services. The financial issue is the most frequent reason for the choice of implant-retained overdentures rather than a fixed implant-supported prosthesis. In a number of countries, the amounts charged for fixed implant-supported prostheses seem to be higher than necessary when analyzing the clinical, laboratory, and material costs. Palmqvist et al., (2004) report that the implant-supported fixed prosthesis can be provided at the same cost as an implant-

retained overdenture when carried out in the All -in -One concept and that the choice between a fixed implant-supported prosthesis and implant-retained overdenture need not be an economic issue any longer.

In the long term, little is known about the cost-effectiveness of conventional dentures, implant-retained overdentures, and fixed implant-supported prostheses in edentulous patients (Heydecker et al., 2005). In most studies, the estimated costs and/or effects have been restricted to monitoring during the trial period (Jönsson and Karlsson, 1990; Tang et al., 1997; Palmqvist et al., 2004; Takanashi et al., 2004). However, most of the costs for prosthetic restorations typically accumulate during the first year, while their benefits are usually seen throughout the prosthetic's functional period (Attard and Zarb, 2004).

A study by Zitzmann et al., (2006) analyzed the cost-effectiveness of three different treatments: conventional dentures, implant-retained overdentures on two implants, and implant-retained overdentures on four implants on an edentulous mandible. They prepared a cost projection over a ten-year period using a mathematical model and came to the conclusion that an implant-retained overdenture on two implants is the most cost-effective treatment option when compared to conventional dentures and the implant-retained overdenture on four implants. The cost of treatment rose substantially with the increasing number of implants, becoming 3 to 6 times higher when the number of implants jumps from 2 to 4 in implant-retained overdentures, as compared with treatment using conventional dentures. The proportion of maintenance costs in relation to initial costs decreased with the increased number of implants used, i.e., 46% of the initial costs for conventional dentures, 28% for implant-retained overdentures on two implants, and 13% in implant-retained overdentures on four implants.

Attard et al., (2005) reported that the cost of initial treatment and maintenance costs during the study's observation period were higher for fixed implant-supported prostheses when compared with implant-retained overdentures. The fixed implant-supported prosthesis, however, presented a greater reduction in maintenance costs with respect to the initial cost. Furthermore, maintenance costs indicate that implant-retained overdenture treatment was cheaper when compared with fixed implant-supported prostheses.

Emami et al., (2009) reported that additional meta-analyses are needed in well-controlled randomized clinical trials that include relevant economic assessments to provide policymakers and insurance companies with the help they need when making decisions about the adoption of implant therapies.

Discussion for Guidance in Clinical Decisions Regarding Edentulous Rehabilitation

Making a decision concerning the rehabilitation of edentulous patients initially involves identifying their profiles and getting answers related to their complaints and expectations. The denture's low retention and stability, especially in the mandible, result in functional deficiency, which in turn has a negative impact on the patient's satisfaction and quality of life. Prostheses supported or retained by an implant partially overcome these limitations and may even approach the functional performance of dentate subjects, thereby improving patient satisfaction and quality of life as related to oral health.

Table 1. Comparison among the decision factors regarding the choice of the type of total prosthesis

Factors	CD	IOD	FISP
Stability/ Retention	Lowest	Average	Greatest
Masticatory efficiency	Lowest	Average	Greatest
Patient satisfaction	Lowest	Improved	Improved
Quality of life	-	Improved	Improved
Number of implants	-	Fewer	Greater
Implant survival	-	Same	Same
Peri-implant resorption	-	Same	Same
Subsequent bone resorption*	Same	Same	Lowest
Soft tissue abrasion	More	Less	Absent
Initial cost	Lowest	Average	Highest
Maintenance /Initial cost	46%	28%	13%

(* = In patients who have been edentulous for more than 10 years).

The number of implants used for this purpose varies with the type of prosthetic option, being higher in fixed implant-supported prostheses than in implant-retained overdentures and inversely proportional to the rate of residual ridge resorption and soft tissue abrasion, especially when looking at conventional dentures. However, implant survival and peri-implant bone loss are not altered by the type of prosthesis. It should be noted that the implants under mandibular overdentures survive longer when compared with the maxillary. The cost aspect is treated last of all because it is currently the primary deciding factor in choosing among these rehabilitation options. The conventional denture has a low initial cost (clinical hours, laboratory, and material), followed by the implant-retained overdenture, and finally the implant-supported fixed prosthesis. However, maintenance costs decrease as the number of implants increases. Maintenance on a conventional removable prosthesis represents 46% of its initial cost, while maintenance on an implant-retained overdenture represents 28%, and on the fixed implant-supported prosthesis, 13%. Table 1 presents a summary of the deciding factors for choosing the type of prosthesis.

Longitudinal follow-up studies would facilitate a better evaluation of these aspects and the possible associations with individual conditions, such as length of time of edentulism, health conditions, and other pertinent factors.

Conclusion

In developing countries, edentulism is regarded as a public health problem. Among the therapeutic options available for the rehabilitation of these patients, the conventional denture is the most frequently used prosthesis.

However, difficulties in the retention of and adaptation to the lower prosthesis are common and represent a major obstacle to the successful use of conventional prostheses. Therapies using implants have increased the range of prosthetic options for treating these

patients. Within the scope of this review, there is no evidence of a single universal treatment modality for edentulous mandibles. Long-term studies are needed to further explore the differences in patient acceptance for each intervention.

References

- Adell, R; Lekholm, U; Rockler, B; Brånemark, P-I. A 15-year study of osseointegrated implants in the treatment of edentulous jaw. *International Journal of Oral Surgery*, 1981 10, 387-416.
- Albrektsson, T; Blomberg, S; Brånemark, P-I. Edentulousness: an oral handicap-patient reactions to treatment with jawbone-anchored prosthesis. *Journal of Oral Rehabilitation*, 1987 14, 503-511.
- Allen, PF. and McMillan, AS. A longitudinal study of quality of life outcomes in order adults requesting implant prostheses and complete removable dentures. *Clinical Oral Implants Research*, 2003 14, 173-179.
- Alsabeeha, N; Atieh, M; Payne, AGT. Loading Protocols for Mandibular Implant Overdentures: A Systematic Review with Meta-Analysis. *Clinical Implant Dentistry and Related Research*, 2010 12, 28-38.
- Anderson, GF. and Hussey, PS. Population aging: a comparison among industrialized countries. *Health Affairs*, 2000 19, 191-203.
- Attard, NJ. and Zarb, GA. Long-term treatment outcomes in edentulous patients with implant overdentures: the Toronto study. *International Journal of Prosthodontics*, 2004 17, 425-433.
- Attard, NJ; Zarb, GA; Laporte, A. Long-term treatment costs associated with implant-supported mandibular prostheses in edentulous patients. *International Journal of Prosthodontics*, 2005 18, 117-123.
- Awad, MA; Lund JP; Dufresne E; Feine JS. Comparing the efficacy of mandibular implant-retained overdenture and conventional dentures among middle-aged edentulous patients: Satisfaction and functional assessment. *International Journal of Prosthodontics*, 2003 16, 110-116.
- Brånemark, P-I; Breine, U; Adell, R; Hansson, BO; Lindström, J; Ohlsson, A. Intra-osseous anchorage of dental prostheses. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery*, 1969 3, 81-100.
- Brånemark, P-I; Hansson, BO; Adell, R; Breine, U; Lindström, J; Hallen, O; Öhman, A. Osseointegrated implants in treatment of the edentulous jaw. Experience from 10-year period. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery*, 1977 16, 1-132.
- Brånemark, P-I; Svensson, B; van Steenberghe D. Ten-year survival rates of fixed prostheses on four or six implants ad modum Branemark in full edentulism. *Clinical Oral Implants Research*, 1995 6, 227-231.
- Bryant, SR; McDonald-Jankowski, D; Kim, K. Does the Type of Implant Prosthesis Affect Outcomes for the Completely Edentulous Arch? *The International journal of oral maxillofacial implants*, 2007 22, 117-139.

- Bury, M. Illness narratives: fact or fiction? *Sociology of Health and Illness*, 2001 23, 263-285.
- Campostrini, EP; Ferreira, EF; Rocha, FL. Condições da saúde bucal do idoso brasileiro (Oral health conditions of Brazilian elderly). *Arquivos em Odontologia*, 2007 43, 48-56.
- Carlsson, GE; Kronström, M; de Baat, C; Cune, M; Davis, D; Garefis, P; Heo, SJ; Jokstad, A; Matsuura, M; Närhi, T; Ow, R; Pissiotis, A; Sato, H; Zarb, GA. A survey of the use of mandibular implant overdentures in 10 countries. *International Journal Prosthodontics*, 2004 17, 211-217.
- Çehreli, MCT; Karasoy, D; Kökat, AM; Akça, K; Eckert, SE. Systematic Review of Prosthetic Maintenance Requirements for Implant-Supported Overdentures. *The International Journal of Oral and Maxillofacial Implants*, 2010 25, 163-180.
- Colussi, CF. and Freitas, SFT. Aspectos epidemiológicos da saúde bucal do idoso no Brasil (Epidemiological aspects of oral health among the elderly in Brazil). *Cad. Saúde Pública*, 2002 18, 1313-1320.
- Cupertino, APFB; Rosa, FHM; Ribeiro, PCC. Definição de envelhecimento saudável na perspectiva de indivíduos idosos. (Definition of healthy aging from the perspective of the elderly). *Psicologia: Reflexão e Crítica*, 2007 20, 81-86.
- De Souza e Silva, ME; De Magalhães, CS; Ferreira, EF. Complete removable prostheses: from expectation to (dis)satisfaction. *Gerodontology*, 2009 26, 143-149.
- Emami, E; Heydecke, G; Rompre, PH; Grandmont, P de; Feine, JS. Impact of implant support for mandibular dentures on satisfaction, oral and general health-related quality of life: a meta-analysis of randomized-controlled trials. *Clinical Oral Implants Research*, 2009 20, 533-544.
- Esposito, M; Grusovin, MG; Willings, M; Coulthard, P; Worthington, HV. The Effectiveness of Immediate, Early, and Conventional Loading of Dental Implants: A Cochrane Systematic Review of Randomized Controlled Clinical Trials. *The International Journal of Oral and Maxillofacial Implants*, 2007 22, 893-904.
- Ferreira, RC; Magalhães, CS; Moreira, AN. Tooth loss, denture wearing and associated factors among an elderly institutionalized Brazilian population. *Gerodontology*, 2007 25, 168-178.
- Fitzpatrick, B. Standard of care for the edentulous mandible: A systematic review. *The Journal of Prosthetic Dentistry*, 2006 95, 71-78.
- Fueki, K; Kimoto, K; Ogawa, T; Garrett, NR. Effect of implant-supported or retained dentures on masticatory performance: A systematic review. *The Journal of Prosthetic Dentistry*, 2007 98, 470-477.
- Heath, MR. The effect of maximum biting force and bone loss upon masticatory function and dietary selection of the elderly. *Int Dent J*, 1982 32, 345-356.
- Heydecker, G; Penrod, JR; Takanashi, Y; Lund, JP; Feine, JS; Thomason, JM. Cost-effectiveness of mandibular two-implant overdentures and conventional dentures in the edentulous elderly. *J. Dent Res.*, 2005 84, 794-799.
- Heydenrijk, K; Raghoobar, GM; Meijer, HJA; van der Reijden, WA; van Winkelhoff, AJ; Stegenga, B. Two-stage IMZ implants and ITI implants inserted in a single-stage procedure. A prospective study. *Clin Oral Implants Res*, 2002 13, 371-380.
- Ikebe, K; Matsuda, K; Morii, K; Furuya-Yoshinaka, M; Nokubi, T; Renner, RP. Association of masticatory performance with age, posterior occlusal contacts, occlusal force, and salivary flow in older adults. *Int. J.. Prosthodont*, 2006 19, 475-481.

- Jacobs, R; Schotte, A; van Steenberghe, D; Quirynen, M; Naert, I. Posterior jaw bone resorption in osseointegrated implant - supported overdentures. *Clin Oral Implants Res*, 1992 3, 63-70.
- Jokstad, A. and Carr, AB. What Is the Effect on Outcomes of Time-to-Loading of a Fixed or Removable Prosthesis Placed on Implant(s)? *The International Journal of Oral and Maxillofacial Implants*, 2007 22, 19-61.
- Jönsson, B. and Karlsson, G. Cost-benefit evaluation of dental implants. *Int. J. Technol Assess Health Care*, 1990 6, 545-557.
- Kapur, KK. and Soman, SD. Masticatory Performance and Efficiency in Denture Wearers. *J Prosthetic Dent*, 1964 14, 687-694.
- Klemetti, E. Is there a certain number of implants needed to retain an overdenture? *Journal of Oral Rehabilitation*, 2008 35, 80-84.
- Kordatzis, K; Wright, PS; Meijer, HJA. Posterior Mandibular Residual Ridge Resorption in Patients with Conventional Dentures and Implant Overdentures. *The International Journal of Oral and Maxillofacial Implants*, 2003 18, 447-452.
- Koshino, H; Hirai, T; Ishijima, T; Ikeda, Y. Tongue motor skills and masticatory performance in adult dentates, elderly dentates, and complete denture wearers. *J. Prosthet Dent*, 1997 77, 147-152.
- Krennmair, G. and Ulm, C. The symphyseal single-tooth implant for anchorage of a mandibular complete denture in geriatric patients: a clinical report. *International Journal Maxillofacial Implants*, 2001 16, 98-104.
- Kronström, M; Davis, B; Loney, R; Gerrow, J; Hollender, L. A prospective randomized study on the immediate loading of mandibular overdentures supported by one or two implants : a 12- month follow-up report. *The International Journal Oral Maxillofacial Implants*, 2010 25, 181-188.
- Lebrão, ML. O envelhecimento no Brasil: Aspectos da transição demográfica e epidemiológica (Aging in Brazil: Aspects of demographic and epidemiological transition). *Saúde Coletiva*, 2007 4, 135-140.
- Lekholm, U. Surgical considerations and possible shortcomings of host sites. *J. Prosthet Dent*, 1998 79, 43-48.
- Meijer, HJA; Raghoobar, GM; Batemburg, RHK; Visser, A; Vissink, A. Mandibular overdentures supported by two or four endosseous implants: 10-year clinical trial. *Clinical Oral Implants Research*, 2010 20, 722-728.
- Moreira, RS; Nico, LS; Tomita, NE; Ruiz, T. Oral health of Brazilian elderly: a systematic review of epidemiologic status and dental care access. *Cad. Saúde Pública*, 2005 21, 1665-1675.
- Naert, I; Alsaadi, G; Quyrinen, M. Prosthetic aspects and patient satisfaction with two-implant-retained mandibular overdentures: a 10-year randomized clinical study. *International Journal of Prosthodontics*, 2004 17, 401-410.
- Naert, I; Hooghe, M; Quirynen, M; van Steenberghe, D. The reliability of implant-retained hinging overdentures for the fully edentulous mandible. An up to 9-year longitudinal study. *Clin Oral Investig*, 1997 1, 119-124.
- Niessen, LC. Geriatric dentistry in the next millennium: opportunities for leadership in oral health. *Gerodontology*, 2000 17, 3-7.

- Palmqvist, S; Owall, B; Schou S. A Prospective Randomized Clinical Study Comparing Implant-Supported Fixed Prostheses and Overdentures in the Edentulous Mandible: Prosthodontic Production Time and Costs. *Int. J. Prosthodont*, 2004 17, 231-235.
- Raghoebar, GM; Meijer, HJA; van't Hof, M; Stegenga, B; Vissink, A. A randomized prospective clinical trial on the effectiveness of three treatment modalities for patients with lower denture problems. *Int. J. Oral Maxillofac Surg.*, 2003 32, 498-503.
- Roncalli, AG. Projeto SB Brasil 2010 – Pesquisa Nacional de Saúde Bucal revela importante redução da cárie dentária no país (National Oral Health Research reveals a significant reduction of dental caries in the country). *Cad. Saúde Pública*, 2011 27, 4-5.
- Shinkai, RSA; Hatch, JP; Rugh, JD; Sakai, S; Mobley, CC; Saunders, MJ. Dietary intake in edentulous subjects with good and poor quality complete dentures. *J. Prosthet Dent*, 2002 87, 490-498.
- Silva, ME; Magalhães CS; Ferreira, EF. Dental loss and prosthetic replacement expectation: qualitative study. *Cien Saude Colet*, 2010/a 15, 813-820.
- Silva, ME; Villaça, EL; Magalhães, CS; Ferreira, EF. Impact of tooth loss in quality of life. *Cien Saude Colet*, 2010/b 15, 841-850.
- Slagter, AP; Bosman, F; van der Bilt, A. A human jaw-elevator muscle activity and food comminution in the dentate and edentulous state. *Arch. Oral. Biol.*, 1993 38, 195-205.
- Slagter, AP; Olthoff, LW; Steen, WH; Bosman, F. Communion of food by complete denture wearers. *J. Dent Res.*, 1992 72, 380-386.
- Sunyoung, M. and Payne, AGT. Marginal Bone Loss with Mandibular Two-Implant Overdentures Using Different Loading Protocols: A Systematic Literature Review. *Int. J. Prosthodont*, 2010 23, 117-126.
- Takanashi, Y; Penrod, JR; Lund, JP; Feine, JS. A cost comparison of mandibular two-implant overdenture and conventional denture treatment. *Int. J. Prosthodont*, 2004 17, 181-186.
- Tang, L; Lund, JP; Taché, R; Clokie, CML; Feine, JS. A within subject comparison of mandibular long-bar and hybrid implant-supported prostheses: psychometric evaluation and patient preference. *J. Dent Res.*, 1997 76, 1675-1683.
- Thomason, JM; Heydecke, G; Feine, JS; Ellis, JS. How do patients perceive the benefit of reconstructive dentistry with regard to oral health-related quality of life and patient satisfaction? A systematic review. *Clin Oral Implants Res*, 2007 18, 168-188.
- Turkyilmaz, I; Company, AM; McGlumphy, EA. Should edentulous patients be constrained to removable complete dentures? The use of dental implants to improve the quality of life for edentulous patients. *Gerodontology*, 2010 27, 3-10.
- Van der Bilt, A; Burgers, M; van Kampen, FM; Cune, MS. Mandibular implant-supported overdentures and oral function. *Clinical Oral Implants Research*, 2010 21, 1209-1213.
- Vargas, AMD. and Paixão, HH. The loss of teeth and its meaning in the quality of life of adults who use the municipal oral health services of the Boa Vista Health Center, in Belo Horizonte. *Cien Saude Colet*, 2005 10, 1015-1024.
- Vervoor, JM; Duinkerke, AS; Luteijn, F; van der Poel, AC. Assessment of denture satisfaction. *Community Dent Oral Epidemiol*, 1988 16, 364-367.
- Walton, JN; Glick, N; Macentee, MI. A randomized clinical trial comparing patient satisfaction and prosthetic outcomes with mandibular overdentures retained by one or two implants. *Int. J. Prosthodont*, 2009 22, 331-339.

- Wayler, AH. and Chauncey, HH. Impact of complete dentures and impaired natural dentition on mastigatory performance and food choice in healthy aging men. *J. Prosthet Dent*, 1983 49, 427-433.
- WHO- World Health Organization. The Ottawa Charter for Health Promotion. In *Health Promotion, World Health Organization*, Geneva, 1986 1.
- Wolff, A; Gadre, A; Begleiter, A; Moskona, D; Cardash, H. Correlation between patient satisfaction with complete dentures and denture quality, oral condition, and flow rate of submandibular/sublingual salivary glands. *Int. J. Prosthodont*, 2003 16, 45-48.
- Zarb, G. and Bolender, C. *Tratamento protético para os pacientes edêntulos* (Prosthetic treatment for edentulous). (12ª Ed). São Paulo: Santos; 2006.
- Zitzmann, NU; Marinello, CP; Sendi, P. A Cost-effectiveness Analysis of Implant Overdentures. *Journal Dentistry Research*, 2006 85, 717-721.

Chapter IX

Denture Adhesives – A Boon or a Bane

Puja Chowdhry Malhotra*

SGT Dental College, Gurgaon, India

Abstract

Denture Adhesives have always been surrounded by an infinite number of controversies. Prosthodontists and dentists in general have never really accepted the role of denture adhesives in improving denture retention, stability and function. Despite substantial and significant records encouraging the use of adhesives, the dental fraternity by large still considers adhesive use as a poor echo of their treatments and prosthetic proficiencies. However, in view of a longer-living population with an increasing rate of chronic residual ridge resorption and a consequent increase in dentures that are unsteady because of parameters beyond the control of the patient or dentist, denture adhesives deserve a place in removable prosthodontics as an effective and active aide in denture treatment and denture aftercare.

Complete dentures constitute one of the most important treatment options in prosthodontics, more so with an increase in average life expectancy of the individual. However, retention of complete dentures has always posed to be a problem for the dentist and the dental industry. Solutions to the problem, over the years have included overdentures, implants, and denture adhesives. From time immemorial adhesives have been used by denture wearers and advertised commercially, however these dental materials have not been given their due place in prosthetic dentistry. These materials are specially promoted in certain special conditions such as complicated prostheses—obturators, dry mouth, difficult and exacting patients (e.g. Parkinson's disease, Alzheimer's disease), compromised ridges, single complete denture or in public figures. Besides, adhesives are also normally used by denture wearers to psychologically support the patient so as to make the complete dentures more acceptable.

Conflicting views still persist. The ancient views regard these materials as a poor replacement for a proper fit and adjustment of dentures. However, existing literature supports the use of denture adhesives i.e. if properly used. Adhesives can be a valuable aid in management of challenging denture patients. Therefore, the following article aims

* E-mail address: pujchow@yahoo.com.

to solve this dilemma regarding the use denture adhesives, oust some of the myths related to adhesives in order to allow good use and avoid misuse of the same.

Introduction

One of the biggest challenges in complete denture Prosthodontics is undoubtedly the oral rehabilitation of a patient with resorbed ridges. One alternative which has been most commonly used by the patients is denture adhesives. In spite of this, little scientific information is available on them and most of the data that is available is based upon patients' verdict, opinion and experiences. Therefore, Dental clinicians have always been in a predicament regarding the use and abuse of denture adhesives, their use has been and still is paralleled to poorly fabricated dentures. Subsequently, the clinicians are still doubtful and not very inclined to prescribe adhesives to denture wearers as a means to boost retention, stability and comfort.

Dentistry has grown exponentially in the last few decades. With the development of implant dentistry, Conventional removable Prosthodontics has taken a back seat. Despite this complete dentures treatment is still considered to be an integral part of prosthodontics, especially with an increase in average life expectancy of the individual. The objectives of impression making (i.e. for retention, support and stability) are designed to accomplish a full coverage of denture bearing areas, a close tissue to denture contact, and an operative peripheral seal. However, in certain situations, the same may not be probable. In these instances patients frequently choose to use denture adhesives for e.g. in circumstances of immediate or single complete dentures, complicated maxillofacial prostheses (obturators, testing patients), hysterical patients, patients suffering from systemic diseases (like myasthenia gravis or parkinsonism etc.), in cases of compromised ridge structures, or public figures (such as attorneys, actors, and public speakers where denture wobble cannot be tolerated). Besides this the adhesives are also usually used as a support during the fabrication phase of a denture and as a means to aid in the application of drugs to the oral mucosa e.g. radiation carriers.

"Denture adhesive" [1] is a commercially available, non toxic, soluble material that when applied to the tissue surface of the denture enhance their retention, stability and performance. These are available as powder, paste, tape or cushion. Soluble adhesives [2] i.e., the powder and paste forms are universally accepted now because they do not damage the soft tissues.

Composition and Mechanism of Action

Stafford in 1970 [3] has given the basic composition of denture adhesives which has also been reported by the manufacturers and the American Dental Composition. The ingredients fall mainly into three categories:

1.: Those materials that swell gel and show increasing and varying degrees of viscosity in water (e.g. gum karaya, gelatin and pectin). These materials also suspend insoluble powders in aqueous preparations and act as emulsifying, dispersing and thickening agents (e.g. sodium carboxy methyl cellulose, hydroxy ethyl cellulose, and methyl cellulose).

2.: Those materials that act as antibacterial/ antiseptic agents (e.g. methyl salicylate, sodium borate, sodium tetraborate, hexachlorophene). These materials are added by some manufacturers but it is doubtful whether these materials serve any real purpose.

3.: Additional materials like sodium lauryl sulphate. This reduces the surface tension and acts as a wetting agent. Propyl hydroxy benzene (propyl paraben) acts as a preservative and is mildly active against *Candida albicans*. Magnesium oxide is a filler. Peppermint oil has flavoring properties with mild antiseptic actions. Petrolatum is used as an emollient.

These materials basically increase the forces of adhesion and cohesion between the intaglio surface of the denture and basal tissues allowing complete wetting and eliminating the voids between the two. Shay [4] described the mechanism of action of adhesives in 1991. These materials swell 50% to 150% by volume in the presence of water, filling the spaces between the prosthesis and the tissues. As water is absorbed by the adhesive agents, the resulting cations are attracted to anions in mucous membrane proteins, producing the stickiness. The properties of current adhesives depend upon the combination of both physical and chemical properties, saliva increases the viscosity of the adhesive thereby increasing the force required to separate the prosthesis from the tissue surface. Most adhesives use ingredients that provide bioadhesion via carboxyl groups. As the adhesive hydrates, free carboxyl groups form electrovalent bonds that produce stickiness. Polymethyl vinyl ether – maleic anhydride (PVM - MA) is a synthetic compound widely used in denture adhesives because of high levels of carboxyl groups. Sodium carboxymethyl cellulose has the advantage of being more soluble, beginning in the early 1970's manufacturers combined divalent salts of PVM-MA, specifically calcium salts with CMC to make denture adhesive. The divalent salt increased product performance by reducing the rate of dissolution and also increasing the cohesive strength of overall material.

In the late 1980s, companies introduced products that combined PVM-MA zinc and calcium salts with carboxy methyl cellulose. These materials provided an even greater cohesive strength for longer duration because of stronger covalent bond that develops via divalent zinc cation. Commercial denture adhesives that contain these materials have clinically proved to provide stronger, longer hold than products containing calcium salts of PVM-MA copolymer alone.

Denture Adhesives – Myth and Reality

It is now accepted that denture adhesives if used sensibly and under supervision can aid in retention of complete dentures. Even though, the ancient former views regard these materials as a poor substitute for a proper fit and adjustment of dentures and their use was equated to poor denture skills and lack of prosthetic expertise [5, 6]. In 1967, Kapur [7] conducted a study on 26 denture wearers and devised a method for scoring denture retention and stability as ill, fair and well fitting prosthesis. He concluded that denture adhesives unequivocally increased denture retention thereby increasing denture wearers' incisive ability.

Tarbet et al. [8, 9] addressed the role of denture adhesives in retention and stability by counting denture dislodgements in patients eating standardized portions of food with and without denture adhesive. Results showed a significant reduction in dislodgement when an adhesive was used. Researchers have devised new techniques such as the cineradiographic

technique and a hydraulic and electrical transducer to measure maximum retentive force and to compare the effects of materials during chewing. A decrease in dislodgement of dentures while chewing was observed [10, 11]. Radiotelemetry and gnathodynamometry have also been used in this field and results have shown that the use of denture adhesives, in patients with poor bone support, improve the retention and stability comparable to that in patients with good bone support [9].

Most recently, a multi-channel magnetometer tracking system has been used to study the effect of denture adhesives on mandibular movement during chewing. The subjects were made to chew dried apricots and fresh white bread. It was concluded from the study that the use of denture adhesives resulted in a faster and more natural rate of chewing [12].

In spite of all the studies done, negative attitude regarding denture adhesives still prevails and a lot of literature focuses on the harmful effects of these materials i.e.: *I.*: enhancing the prolonged use of ill fitting dentures thus promoting residual ridge resorption,; *II.*: interfering with the occlusion as a result of uneven and uncontrolled thickness of intervening adhesive; *III.*: act as allergens [13]. This is especially true for cushion type of denture adhesives which are insoluble adhesives and involve a greater risk of inducing alveolar ridge resorption. These “home reliners” are not accepted by ADA. The soluble adhesives cannot be abused to the extent of changing vertical dimension since they rapidly absorb water becoming gelatinous and spreading over the denture through chewing stress [2, 14, 15].

In 1980, Tarbet and Grossman [16] reported a six-month investigation on incidence and severity of mucosal irritation in a group of 111 denture wearers using adhesive materials. No increased incidence of mucosal irritation was observed, on the contrary the irritation present at the time of insertion of dentures showed a definite decline. However, at the same time denture adhesive should not be prescribed on a regular basis as they do increase the wearing time of ill-fitting dentures leading to deterioration of bony support [16].

Use or Abuse

Denture Adhesives act as a cushion, reduce the collection of food under the denture flanges, hinder the growth of *Candida*. They help to distribute the occlusal load during various centric and eccentric movements. Adisman [2] in 1989 said that dentists may prescribe and use denture adhesives as a legitimate, therapeutic and effective procedure in denture treatment and denture after care. The patients should be instructed in its proper use and cautioned against misuse. The use of denture adhesives in ill fitting dentures is contraindicated. All denture patients should be advised to consult their dentists for periodic examinations of their dentures and the health of the oral cavity.

Denture Adhesives are commonly available in paste and powder forms. The paste forms of denture adhesives are more retentive than powder forms which has been attributed to an increased viscosity of the pastes. This was demonstrated by Chew [17] in 1990 and was later confirmed by Ghani et al. [18] who demonstrated that the liquid / paste form of denture adhesive rendered the ill-fitting dentures almost as retentive as well fitting one.

Retention and stability of dentures has been a continuing problem for dentists in clinical practice. Retention of dentures in the oral cavity is a multifactorial phenomenon which relies upon a number of variables namely adhesion, cohesion, atmospheric pressure, surface tension

and viscosity, these are in turn determined by the quantity and quality of saliva interspersed between the mucosal surface and the hard acrylic of the corresponding surface of the dentures. Studies conducted in the past, have distinctly shown that the denture adhesives significantly augment the retentive ability of complete dentures. This rise could probably be explained by the consequent rise in viscosity of the intervening film of saliva existing between the acrylic dentures and the mucosal surface. Thus, denture adhesives can be used to improve denture retention, offering security, certainty, assurance, and function to the patient but at the same time it must be noted that denture adhesives improve denture retention only if the principal properties of the complete denture such as retention, stability and close adaptation of the denture base to the underlying tissues are satisfactory. An ill-fitting, poorly fabricated complete denture that does not satisfy the basic requirements of retention and stability will not become retentive with the mere use of adhesive materials. On the contrary, the unwise, imprudent use of such materials may lead to undesirable harm and deterioration of supporting structures, which may be permanent and irreversible.

Dentists should caution the patient against the misuse of such materials and insist upon the regular recall visits for evaluation of their dentures and health of the oral cavity. Woelfel et al. [19] in 1965 investigated the hazards of additives sold over the counter and they concluded that a longer than normal wearing period of complete dentures is encouraged by the sale of over-the-counter denture additives, adhesives, cushions, pads, reliners etc. A four-fold program was suggested in an effort to promote better denture health among denture wearers: *I.*: Educate the public to the full benefits of professional dental care, including essential periodic recalls while wearing complete dentures; *II.*: Warn patients against buying or using any do-it-yourself additives on their dentures; *III.*: Prohibit the promotion and public sale of all over the counter reliners, cushions and repair kits; *IV.*: Place the sale of denture adherent powders and pastes on a basis whereby a prescription from a dentist would be required to obtain them from a pharmacist. Most recently Tezvergil - Mutluay, have reported an increase in serum zinc levels and decrease in serum copper levels in denture patients overusing denture adhesives or rather misusing denture adhesives. This condition is known to cause bone marrow suppression and widespread sensory and motor neuropathies. Epidemiologic studies have revealed the source of excessive zinc from overuse of denture adhesives [20].

Conclusion

Denture adhesives are commonly used by denture wearers to increase the retention and stability of the dentures, to improve the chewing and masticatory efficiency, and to psychologically support the patient so as to make the complete dentures more acceptable. In 1965, Great Britain alone produced about 88 tons of denture adhesives, 12% of population included women and 10% men [3] and in the U.S more than 5 million denture wearers use denture adhesives [2]. This number is on a continuous rise because of the advances in the medical field which has led to an increase in average life expectancy of the individual. Literature review has revealed the use of mechanical devices like springs, suction cups, suction chambers, magnets, undercuts, overdentures and implants to alleviate retention problems in the difficult aforementioned conditions. Although most of premised devices may

increase the retention, but many of them either cause further damage to the foundation tissues, or are rather expensive (i.e. implants, magnets). Therefore, denture adhesives being commercially and readily available, has emerged as an acceptable solution to meet the challenges of retention in such patients. Numerous surveys have been conducted in the past taking into consideration the patient's opinion on the effectiveness of adhesive materials. In general, most of the patients responded positively indicating that the retention was little better or much better with the use of adhesive materials [21].

References

- [1] Zarb, GA; Bolender, Cl. *Boucher's Prosthodontic Treatment of the edentulous patient*, 11th edition, St Louis: Mosby; 1997; pp 442
- [2] Adisman, IK. The use of denture adhesives as an aid to denture treatment. *J. Prosthet Dent*, 1989 62, 711-715.
- [3] Stafford, GD. Denture Adhesives – A Review of their uses and compositions. *Dent Practit*, 1970 21, 17-19.
- [4] Shay, K. Denture Adhesives – Choosing the right powders and pastes. *J. Am. Dent. Assoc.*, 1991 122, 70-76.
- [5] Mc Kevitt, FV. The measured vertical dimension and denture adhesive powders. *J. Prosthet Dent*, 1951 1, 393-401.
- [6] Woelfel, NB; Curry, RL. Additives sold over the counter dangerously prolong wearing period of ill fitting dentures. *J. Am. Dent. Assoc.*, 1965 71, 603-613.
- [7] Kapur, KK. A clinical evaluation of denture adhesives. *J. Prosthet Dent*, 1967 18, 550-558.
- [8] Tarbet, WJ; Boone, M; Schmidt, NF. Effect of denture adhesive on complete denture dislodgement during mastication. *J. Prosthet Dent*, 1980 44, 374-378.
- [9] Tarbet, WJ; Boone, M; Schmidt, NF. Maximum incisal biting force in denture wearers as influenced by adequacy of denture-bearing tissues and the use of an adhesive. *J. Dent Res.*, 1981 60, 115-119.
- [10] Karlsson, S; Swartz, B. Effect of denture adhesive on mandibular denture dislodgement. *Quintessence Int*, 1990 21, 625-627.
- [11] Ow, RKK; Bern, EM. A method of studying the effect of adhesives on denture retention. *J. Prosthet Dent*, 1983 50, 332-337.
- [12] Rendell, JK; Gay, T; Grasso, JE; Baker, RA; Winston, JL. The effect of denture adhesive on mandibular movement during chewing. *J. Am. Dent. Assoc.*, 2000 131, 981-986.
- [13] Panagiotouni, E; Pissiotis, A; Kapari, D; Kaloyannides, NN. Retentive ability of various denture adhesive materials: an in vitro study. *J. Prosthet Dent*, 1995 73, 578-585.
- [14] Means, CR. A Report of a User of Home Reliner Materials. *J. Prosthet Dent*, 1964 14, 935-938.
- [15] Woelfel, JB; Kreider, JA; Berg, TJ. Deformed Lower Ridge Caused by Relining of a Denture by a Patient. *J. Am. Dent. Assoc.*, 1962 67, 763-769.

-
- [16] Tarbet, WJ; Grossman, E. Observations of denture supporting tissue during six-months of denture adhesive wearing". *J. Am. Dent. Assoc.*, 1980 101, 789-791.
 - [17] Chew, CL. Retention of denture adhesives – an in-vitro study. *J. Oral. Rehabil.*, 1990 17, 425-434.
 - [18] G'hani, F; Picton, DCA; Likeman, PR. Some factors affecting retention forces with the use of denture fixatives in in-vivo. *Brit. Dent J.*, 1991 171, 15-21.
 - [19] Woelfel, NB; Curry, RL. Additives sold over the counter dangerously prolong wearing period of ill fitting dentures. *J. Am. Dent. Assoc.*, 1965 71, 603-613.
 - [20] Tezvergil-Mutluay A; Carvalho RM ; Pashley DH. Hyperzincemia from ingestion of Denture Adhesives . *J. Prosthet Dent*, 2010 103 : 380-383.
 - [21] Kesley, CC; Lang, BR; Wang, RF. Examining patients' response about the effectiveness of five denture adhesive pastes. *J. Am. Dent. Assoc.*, 1997 128, 1532-1538.

Chapter X

Changes in the Mycological Ecology of Dentures

Carlos Madrid^{*1}, Jacques Bille² and Marcelo Abarca¹

¹University of Lausanne, School of Medicine, Lausanne, Switzerland,
Department of Oral Surgery, Oral Medicine and Hospital Dentistry, Switzerland
Department of Ambulatory Care and Community Medicine, Switzerland

²University of Lausanne, School of Medicine, Lausanne
Institute of Microbiology, Centre Hospitalier Universitaire Vaudois, Switzerland

Abstract

A wide range of medical devices used in elderly patients shown to support colonization and biofilm formation by *Candida spp.* Among them, more attention must be paid to denture plaque because similar to dental plaque it also serves as a reservoir of potentially infectious microorganisms. Continuous swallowing or aspiration of microorganisms from denture plaque exposes elderly patients to the risk of unexpected infections. The level of oral hygiene in elderly people, especially those living in long-term care institutions, has been reported to be poor, both for dentures and remaining teeth. The upper denture has been suggested to be the major source of infection in long term hospital care patients. It has also been reported that larger quantities of *Candida spp.* reside in the denture base and not in the palate.

The most frequent isolated yeast species in dentures is *C. albicans*. Although, recent reports show high proportions of *non-albicans Candida spp.*, thus showing a shift from *C. albicans* towards *non-albicans spp.* This shift has already been reported over the last years with *C. glabrata* emerging as the second most prevalent species found in dentures. Infections related to *C. glabrata* can lead to systemic infections with high mortality rates. The increased inadequate use of topical and systemic antifungals has been proposed to contribute to this fungal profile change.

Several studies show a correlation between poor denture hygiene and prevalence of *Candida* species and also a statistically significant relationship between denture related

* E-mail addresses: carlos.madrid@hospvd.ch.

stomatitis (DRS), presence of *Candida* and denture cleanliness. High frequency and quantity of *Candida spp.* detection is sometimes associated with few clinically detectable cases of DRS. Hyphae detection by microscopy is now the gold standard (with the exception of *C. glabrata*) for the diagnosis of oropharyngeal fungal infections (OFI). Presence of oral lesions associated with OFI has been reported only in one third of microscopically affected patients.

Frequent use of toothpaste to clean dentures has an influence on dentures surface roughness due to the presence of abrasives. Surface roughness being positively correlated with the rate of fungal colonization of biomaterials, a rougher surface may be a risk factor for microorganism adhesion and biofilm formation.

Conclusion: High levels of *Candida spp.* are reported on dentures of elderly patients. A high frequency of *non-albicans Candida spp.* (particularly *C. glabrata*) has been detected in recent studies. This seems to confirm a change in the mycological ecology involved in yeast denture colonization and DRS. Keywords: *Candida spp.*, dentures, elderly.

Introduction

Oropharyngeal candidosis and the particular form of denture related stomatitis (DRS) are a main concern among denture users and oral health practitioners. Poor fitting and/or prolonged use of the denture are the most common factors related to denture related stomatitis (Budtz-Jørgensen, 1981). The upper denture has been suggested to be the major source of infection in long term hospital care patients (Wilkie *et al.*, 1991). Denture plaque and poor oral hygiene are commonly associated with *Candida* infection (Coulthwaite and Verran, 2007). Moreover, dentures may serve as a reservoir of other potentially infectious pathogens (Sumi *et al.*, 2003). Continuous swallowing or aspiration of microorganisms from denture plaque for example, exposes elderly patients to the risk of unexpected infections (Nikawa *et al.*, 1998).

Among elderly patients, especially those in a frail situation, the lack of instructions concerning oral and denture hygiene is long past reported (Arendorf and Walker, 1987). The level of oral hygiene in elderly people living in long-term care institutions has been reported to be poor, both for dentures and remaining teeth (De Visschere *et al.*, 2006). Recently, Kossioni confirms a common belief among the patients that after provision of dentures, elderly patients assume that regular recall is not necessary anymore (Kossioni, 2010). Several studies show a correlation between poor denture hygiene and prevalence of *Candida spp.* They also demonstrate a statistically significant relationship between DRS, presence of *Candida* and denture cleanliness (Kanli *et al.*, 2005, Kulak-Ozkan *et al.*, 2002). Besides, the frequency of denture brushing does not mean effective cleansing. Cognitive deficits, loss of visual acuity and lack of manual dexterity are a common issue in older people, and this strongly influences the effectiveness of brushing (Nevalainen *et al.*, 1997, Darwazeh *et al.*, 2001). Dentures night – wearing is also a common habit within the elderly. In a group of 150 complete denture wearers, 64% (96 subjects) had the habit to sleep with their dentures (de Castellucci Barbosa *et al.*, 2008).

Gendreau and Loewy published a systematic review focused on epidemiology of DRS. Data from several countries around the world were collected and presented. The usual assessment method of DRS in the different studies was generally based on the scales

developed by *Newton* or *Budtz-Jorgensen and Bertman*, meaning that oral examinations (some of them standardized) were performed by a trained dentist or by an undefined examiner. The number of subjects wearing dentures in the different studies varies between 37 (Scottish study) and 3450 (a US - NHANES III study). DRS prevalence among the different studies varies between 14.7% (Slovenian study/ n=163) and 77.5% (Canadian study/ n=40), and for the larger studied population in Finland the prevalence of DRS was of 48%, distributed in 43.2% for males and 50.6% for females (*Gendreau and Loewy, 2011*).

***Candida* on Dentures and other Implantable Medical Devices**

Candida spp. may develop as biofilm on a wide range of artificial materials used in medicine. These biofilms protect them from antifungal and host defenses. As a matter of fact, shunts, prostheses (valve, knee, etc.), stents, implants, endotracheal tubes, pacemakers and catheters have been shown to support colonization by yeasts (*Ramage et al., 2006*).

Many cases of infections related to medical devices by *Candida* spp. have been described in the literature: infections of peritoneal dialysis catheters leading to *Candida* peritonitis (*Bibashi et al., 2003*); yeast infections in neurosurgical shunts, which have increased within the last few years (*Montero et al., 2000*); and, even if scarce, knee and arthroplasties infections with *Candida* (which are mainly related to *C. albicans*, *C. parapsilosis* and *C. glabrata*) have also been reported (*Phelan et al., 2002*).

Dentures and oral implants are also medical devices used in patients (mainly elderly). They too have shown to support colonization and biofilm formation by *Candida* spp. (*Douglas, 2002*). Denture wearing is therefore commonly associated with DRS, which is characterized primarily by an inflamed mucosa (*Coco et al., 2008*). Oropharyngeal candidosis have a high prevalence of 25% in partially or fully edentulous patients, with 68% of them suffering of DRS (*Madrid et al., 2008*).

Adhesion and Colonization of Dentures by *Candida* Spp.

Candida, as well as other yeasts, has the ability to adhere and grow to synthetic materials in a similar way as they do to oral tissues (*Blankenship and Mitchell, 2006*). Dentures are made principally of polymethylmetacrylate, sometimes associated to soft liners, tissue conditioners or dental adhesives, and all these materials are permeable and/or present depression. This means that yeasts may be easily retained and form complex biofilms (*Allison and Douglas, 1973*).

Several publications of *Sumi and coworkers* show that respiratory pathogens preferentially colonize dentures rather than soft tissues like the palate (*Sumi et al., 2002, Sumi et al., 2003, Sumi et al., 2007*).

Bacteria's proliferation has a close relationship with the development of *Candida* spp. in biofilm (*Hsu et al., 1990*). At least in vitro, the adhesion of *C. albicans* to the acrylic would

be in presence of streptococci (*S. sanguis* and *S. salivarius*) previously colonizing an acrylic surface like the dentures. Another in vitro study has shown that colonization of the acrylic by *C. albicans* is improved if this occurs simultaneously with *S. mutans* (Branting et al., 1989).

The different characteristics of free energy of a biomaterial surface, of surface roughness, and of hydrophobic modification influence both the quantity and quality of fungal adhesion (Ramage et al., 2006, Klotz et al., 1985, Park et al., 2003). Surface roughness has been positively correlated with the rate of fungal colonization of biomaterials, so that a rougher surface may be a risk factor for microorganism adhesion and biofilm formation (Pereira-Cenci et al., 2008, Quirynen et al., 1990). When the adhesion of *C. albicans* to a clear heat cured polymethylmetacrylate was compared with two different soft lining materials, each of them with varying degrees of surface roughness, it has been demonstrated that, of the 3 different tested materials, there was less adhesion of *Candida* to smooth surfaces than to rough surfaces. There was also less adhesion of *C. albicans* to acrylic surfaces than to the soft lining materials. Finally, adhesion of *Candida* was reduced by the presence of a salivary pellicle (Radford et al., 1998).

As mentioned above, the ability of *Candida* to develop themselves on oral implants, which are commonly used to support dentures (e.g. overdentures), needs also to be considered, as *C. albicans* has shown to adhere and grow over different implant surfaces such as machined titanium, sand-blasted titanium, sand blasted and acid etched titanium and zirconia (Bürgers et al., 2010).

Most patients suffering from chronic periodontitis wear partial or total dentures. Adhesion of *C. albicans* to epithelial cells (samples collected at subgingival sites) has been reported to be significantly higher in the periodontitis group when compared to a control group (periodontitis free). Even if the role of *Candida* spp. in periodontitis is still controversial and not yet entirely established, the periodontal pocket might serve as reservoir for regrowth of *Candida* in partial edentulous patients wearing dentures (Machado et al., 2011a).

Denture Related Stomatitis: Clinical Manifestation and Diagnosis

Candida spp. are present in the normal oral flora of an important proportion of healthy individual (Fotos et al., 1992). As commensal members of the oral microflora, they are present in 40% (range 20-60) of the human population (Webb et al., 1998). Certain specific conditions like local factors (e.g., xerostomia or hyposalivation, steroid inhaler use, trauma provoked by the denture) and/or systemic factors (e.g., immunodeficiencies like HIV, systemic corticosteroid use, broad spectrum antibiotic use, malnutrition) increase the risk of overgrowth in older persons (Gonsalves et al. 2008).

The clinical presentation of oral candidiasis ranges from no symptoms (for the majority of the cases) to a prominent burning sensation or an unpleasant salty or bitter taste. The most readily recognized clinical pattern, which is the acute pseudomembranous candidiasis (thrush), is characterized by adherent, curd-like plaques that can be removed with a tongue blade or gauze. Another clinical pattern is angular cheilitis, which is a manifestation of *Candida albicans* (but at times *Staphylococcus*) infection. (Gonsalves et al., 2008).

In denture wearers, oral candidiasis may lead to an erythematous lesion (DRS) (Coco *et al.*, 2008). According to 1962 Newton's classification, revised by Budtz-Jørgensen and Bertram in 1970, 3 types of DRS have been described, based on their clinical features: Type I (localized inflammation or hyperaemia), Type II (diffuse erythema), or Type III (papillary hyperplasia of the palate) (Newton 1962, Budtz-Jorgense and Bertram, 1970).

Microbiologic diagnosis uses different samples and techniques, which include: oral rinses, imprint cultures, and swabbing of specific oral tissue sites (e.g., palate) or swabbing of the denture itself (Bagg *et al.*, 2003). However, hyphae detection (an important factor related to the invasiveness, adhesive capacity and virulence) by means of microscopy is now the gold standard for the diagnosis of DRS fungal infections (Webb *et al.*, 1998).

With regard to basic morphologic microscopic features, all yeasts are Gram-positive. The shapes of blastospores can vary from an ovoid to an elongated or spherical form. Size for the *C. albicans* varies between 3-7 x 3-14 μm (Odds, 1988). Since *C. glabrata* cannot form pseudohyphae, this feature is one of the major identification differences with other *Candida* spp. Presence of germ tubes and chlamydospores are helpful in the identification of *C. albicans* (Warren *et al.*, 1991). Several phenotypic or molecular approaches have been developed for fast and accurate identification of *C. glabrata*, including gene sequence by PCR, restriction fragment length polymorphism (RFLP) or randomly amplified polymorphism DNA (RAPD) methods. However, molecular means are laborious, not very useful in large-scale epidemiological studies and they require expensive equipments (Valerio *et al.*, 2006, Li *et al.*, 2006).

Mycological Changes in Dentures

Until today, the most frequent isolated yeast species in dentures is still *C. albicans*, due to its adhesion and proliferation ability on soft and hard tissues (e.g., dentures, implants) (Ramage *et al.*, 2005). *C. albicans* has been isolated from the palate mucosa (89,2%) and/or maxillary denture (78,5%) in patients suffering from DRS (Gasparoto *et al.*, 2009).

Nevertheless, a high proportion of *non albicans Candida spp.* has been detected in several studies throughout the last years, showing a shift from *C. albicans* towards *non albicans Candida spp.*, with *C. tropicalis*, *C. parapsilosis*, *C. glabrata*, *C. krusei* and *C. dubliniensis* isolated from diseased tissues (Samaranayake and Samaranayake, 2001, Coleman *et al.*, 1997, Samaranayake and Samaranayake, 1994). This shift has been already confirmed within the last years with *C. glabrata* emerging as the second most prevalent specie (Coco *et al.*, 2008, Vanden Abeele *et al.*, 2008). Systemic infections of immunosuppressed patients related to *C. glabrata* can lead to high mortality rates (Krcmery, 1999). Bagg *et al.* have reported in cancer patients that 72% of strains of *C. glabrata* were resistant to fluconazole and itraconazole (Bagg *et al.*, 2003). The increased treatment of patients with some antifungal drugs has been proposed to contribute to this fungal profile change (Martinez *et al.*, 2002).

The presence of *C. dubliniensis* in dentures demands a special attention. *C. dubliniensis* was first obtained from oropharyngeal lesions of HIV positive patients (Sullivan and Coleman, 1998). But diabetic patients also exhibit a high prevalence of *C. dubliniensis* (Willis *et al.*, 2000). This non-albicans species is commonly associated with immunosuppressed

individuals (Davies *et al.*, 2002). Nevertheless, some non-immunosuppressed populations have shown the presence of *C. dubliniensis* (Madrid *et al.*, 2010). Thus, the close phenotypic similarities of *C. albicans* and *C. dubliniensis* could probably led to the misidentification of this species in the past. The presence of *C. dubliniensis* was recently demonstrated by PCR or culture in palate and maxillary denture samples from denture wearers with or without DRS. *C. dubliniensis* was isolated at low rates from both palate (5.3 % and 10.7 %) and maxillary denture (5.3 % and 8.9 %) samples from wearers, regardless of the presence of the disease. However, when *C. dubliniensis* was detected in individuals with DRS, it was always associated with *C. albicans* (Gasparoto *et al.*, 2009).

Systemic Conditions, Aging and DRS

The principal germ implicated in oropharyngeal candidiasis is *C. albicans*, but the non *Candida* spp. (e.g. *C. tropicalis*, *C. glabrata*) are responsible for 46% of the systemic infections caused by this group of yeasts (Samaranayake, 1992, Wingard, 1995).

At cellular level it is important to recognize that, because they are early recruited to sites of infections, neutrophils have an influence on the establishment of Oral Diseases like DRS (Lükac *et al.*, 2003). Neutrophils are able to destroy yeasts by phagocytosis and by production of reactive oxygen species (Shoham and Levitz, 2005). Impaired microbicidal capacity against *C. albicans* has been demonstrated in elderly patients (Schroeder and Rink, 2003). Gasparoto and co-workers reported that individuals presenting DRS show a lower number of salivary neutrophils and dysfunctions in the phagocytosis and in the killing of *C. albicans* (Gasparoto *et al.*, 2009).

People with HIV form a particular group of patients in DRS infections. The primary targets of virus HIV are CD4 + lymphocytes and cells of the monocyte/macrophage lineage. CD4+ lymphocytes are important in the defense against microorganisms such as *Candida* (Perezous *et al.*, 2005). Recurrent oropharyngeal candidiasis develops in the majority of HIV positive patients with CD4+ counts below 300 cells/mm³ (McCarthy *et al.*, 1991). The denture surface can serve as a reservoir not only for local infection like DRS but also for disseminated ones (e.g. esophageal candidiasis). Even after the cleaning of the denture, this surface may influence the quantity and quality of fungal adhesion because of its roughness (Radford *et al.*, 1998). Recent clinical data show that while *C. albicans* remains the most common etiological agent of oropharyngeal candidiasis in patients with advanced AIDS (54%), *C. dubliniensis* (17%) and *C. glabrata* (16%) are the second and third most frequently isolated species, which may exhibit decreased susceptibility or resistance to antifungal agents like fluconazole (Thompson *et al.*, 2011).

Malnutrition is frequent in frail older individuals. A poor oral status has been associated with malnutrition (Sheiham, 2001). The relationship between oral candidiasis and nutritional deficiencies has been reported.

A high prevalence of undernutrition (lower leg circumference and lower serum nutritional protein levels) was reported in a population with candidal infection when compared to control a group without infection (Paillaud *et al.*, 2004).

Finally, nicotine addiction must be considered as a systemic condition with local effects on the oral cavity. It has been reported that DRS was facilitated by the presence of

histopathological alterations of oral tissues more frequent in smokers older than 75 years or in patients consuming more than 15 cigarettes/day (MacEntee et al., 1998, Shulman et al., 2005).

Topical and Systemic Antifungal Therapy

Compared to the accessibility of antibiotics, few antifungal drugs are available. Because of increasing drug resistance, probably provoked by repeated or suppressive courses of low-dose of antifungal drugs, assessment of patient's medications must be performed before starting therapy, in order to modify the etiologic factor causing DRS, e.g. change of drugs provoking reduction of salivary flow (Williams et al., 2011, Perezous et al., 2005). It is also essential to treat treating both the patient and the denture (need of rebase of merely make new dentures) the latter often acting as a reservoir of *Candida* isolates (Tawara et al., 1996).

Topical alternatives to treat DRS are polyenes (nystatine, amphotericine B) or imidazoles (clotrimazole, miconazole or ketoconazole) for a period of 14 days. Topical administration is safe, with very few side effects (Perezous et al., 2005). However, it must be noted that within immunocompromized patients relapses are common using this kind of administration (Scully and McCarthy, 1992).

Because of its good systemically delivery and the good concentration in the saliva (similar to what is attained in the blood), fluconazole (100mg day/14 days) is the first choice to treat complex forms of DRS. Even if secondary effects of fluconazole are limited, important interactions do occur with coumarin anticoagulants and sulfonylurea antidiabetic agents (Williams et al., 2011).

As topical alternative, chlorhexidine (CHX) mouth rinses are regularly used as antiseptic in oral care. In concentration of 0.2% CHX has a wide range spectrum of action against oral bacterias and yeasts like *Candida* spp., causing coagulation of nucleoproteins with inhibition of budding and cell changes. Furthermore, CHX also inhibits candidal adhesion to surfaces like dentures (Ellepola and Samaranayake, 2001). One recent in vitro study shows that immersion of sterile dentures inoculated with an azole-resistant *C. albicans* strains in 2% CHX is effective as a disinfection method (Mima et al., 2011). Also a clinical pilot study demonstrates that CHX gel 0.2% reduces by 68% the *Candida* spp. counts after 21 days of use in immunocompromized patients (Machado et al., 2011). Essential oil mouthwashes have also anticandidal in vitro activity, but the clinical benefits of these agents against *Candida* spp. remain to be established in future studies (Williams et al., 2011, Filoche et al., 2005).

C. glabrata, as abovementioned, associated with systemic infections having a high mortality rate is especially difficult to treat.

A mixed infection with *C. albicans* can cause even more severe symptoms and is therefore still more difficult to treat. *C. glabrata* is naturally resistant to a wide variety of antifungal drugs (innate or acquired) It also hosts derived molecules (e.g. defensins, histatins) compared to other *Candida* spp. Moreover, *C. glabrata* exhibits higher adherence to denture surfaces than *C. albicans* (Luo and Samaranayake, 2002, Redding, 2001, Redding et al., 1999).

Disinfection and Maintenance of Dentures

The understanding of an effective antibiofilm approach is essential to achieve adequate cleaning of dentures. Tooth brushing is a basic method to combat biofilms if it is combined with non-abrasive products (e.g. toothpaste) (Williams *et al.*, 2011). Unfortunately, a low percentage of dentures worn by elderly people are properly cleaned and additionally older dentures tend to be dirtier than the newer ones (Kanli *et al.*, 2005). The institutionalized elderly patient often needs help with oral health care and in particular with the cleaning and disinfecting of dentures. Therefore, a weekly professional mechanical cleaning of the dentures and oral cavity can be an important strategy for the prevention of diseases like aspiration pneumonia (Ishikawa *et al.*, 2008).

Several disinfection methods have been proposed. However, in most of the cases the dentures underwent sanitization rather than sterilization. Many disinfection procedures have been proposed: mechanical cleansing, microwave with varying temperatures, sonicating dentures, immersing the dentures (e.g. in chlorhexidine), using a vacuum, photodynamic therapy etc. In general, all methods reported different degrees of reductions in the levels of microorganisms (Glass *et al.*, 2011, Machado *et al.*, 2011, Ribeiro *et al.*, 2011).

Dentures contaminated by *C. albicans* and irradiated using different microwave energy for different time intervals and power, exhibited reduction in number of counts of yeasts. As a result, surfaces with larger biofilm areas required longer irradiation exposure to be disinfected (Senna *et al.*, 2010). An increased roughness after several cycles of disinfection of the dentures has been reported, and till today there is no standardized method for this technique (Izumida *et al.*, 2011, Augusto Brondani *et al.*, 2010).

Uludamar and coworkers tested different denture disinfection methods in patients with DRS by spraying different mouthwashes on their dentures, like CloSYSII/Chlorine dioxide (Portola Plaza Dental Group, Mission Viejo, CA, USA), and Corsodyl/0.2% chlorhexidine gluconate (GlaxoSmithKline Consumer, Health Group, Oakville, Ontario, Canada) or by using effervescent tablets of different denture cleaners like Polident (GlaxoSmithKline Consumer Health Group, Oakville, Ontario, Canada), Efferdent (Pfizer Consumer Health Care, Scarborough, Ontario, Canada) and Fittydent (Mag Hoeveler Co., Geinberg, Germany) placed in sterile distilled water where the dentures were immersed. Mouthwashes and the effervescent tablets were tested for different periods of time (15, 30 and 60 minutes). All materials tested showed a reduction in the number of *C. albicans*, but mouthwash-sprays are the most easy-to-use and effective method (Uludamar *et al.*, 2010). In general, denture cleansers are effective against *C. albicans* isolates. However, due to biofilm retention, regrowth on denture surface has been observed. This means that a mechanical cleansing will always be required (Jose *et al.*, 2010).

The American College of Prosthodontists published recently evidence-guidelines for care and maintenance of dentures, to reduce potentially harmful bacteria and fungi. For denture cleaning they suggest: the daily cleaning with non abrasive denture cleanser; cleansers should be used only to clean dentures outside the mouth; dentures should be stored in water only after efficient cleansing in order to avoid warping; even if the evidence is weak, the dentures should be cleaned annually by a dentist or dental professional using ultrasonic cleansers (for extensive review see Felton *et al.*, 2011).

Conclusion

The world is facing a significant augmentation of the population older than 60 year-old. Even if there has been stabilization over the last years, an augmentation from 20% to 30% is to be expected in 2030 (*Bellomo et al., 2005*).

In 2050 there is an estimation of 2 billion older people, and 80% of them will be patients in non-industrialized countries (*Erik Petersen, 2003*). The increasing number of people who will be wearing dentures seems obvious then.

Edentulous patients represent a particular group. These patients report not to receive much instructions concerning oral and denture hygiene (*Arendorf and Walker, 1987*). As mentioned above, DRS is a common problem among denture users. Additional measures, such as regular adjustment (rebase) and professional cleaning (in a dental laboratory) by means of mechanic - manual cleaning, polishing and decontamination of the denture, might contribute efficiently to eradicate microorganisms (e.g. *Candida* spp.) deeply lodged into the removable dentures. This approach may be far more effective than the inadequate use of antifungal agents.

The quality of life can be largely influenced by oral conditions. *Gagliardi* and co-workers focused on the relationship between oral care and quality of life, showing a significant improvement of it after oral rehabilitation in a population of 119 subjects ranging from 75 to 97 years old (*Gagliardi et al., 2008*).

The prolonged use of dentures, and especially not removing them at night, associated with poor hygienic habits, a low daily cleansing frequency and inadequate techniques of cleansing (use of abrasive means) and disinfection methods are probably related to the frequent detection levels of *Candida* spp.

The high frequency of *non-albicans Candida* species (particularly *C. glabrata*) detected in different studies seems to confirm a change in the mycological ecology involved in yeast denture colonization.

References

- Allison RT, Douglas WH. Micro-colonization of the denture-fitting surface by *Candida albicans*. *J. Dent.* 1973 1, 198-201.
- Arendorf TM, Walker DM. Denture stomatitis: a review. *J. Oral. Rehabil.* 1987 14, 217-227.
- Augusto Brondani M, Samim F, Feng H. A conventional microwave oven for denture cleaning: a critical review. *Gerodontology*. (in press, accepted 2010 Nov 17).
- Bagg J, Sweeney MP, Lewis MA, Jackson MS, Coleman D, Al MA, Baxter W, McEndrick S, McHugh S. High prevalence of non-albicans yeasts and detection of anti-fungal resistance in the oral flora of patients with advanced cancer. *Palliat Med.* 2003 17, 477-81.
- Bellomo F, de Preux F, Chung JP, Julien N, Budtz-Jørgensen E, Müller F (2005). The advantages of occupational therapy in oral hygiene measures for institutionalized elderly adults. *Gerodontology*. 2005 22, 24-31.

- Bibashi E, Memmos D, Kokolina E, Tsakiris D, Sofianou D, Papadimitriou M. Fungal peritonitis complicating peritoneal dialysis during an 11-year period: report of 46 cases. *Clin. Infect Dis.* 2003 36, 927-931.
- Blankenship JR, Mitchell AP. How to build a biofilm: a fungal perspective. *Curr. Opin. Microbiol.* 2006 9, 588-594.
- Branting C, Sund ML, Linder LE. The influence of *Streptococcus mutans* on adhesion of *Candida albicans* to acrylic surfaces in vitro. *Arch. Oral. Biol.* 1989 34, 347-353.
- Budtz-Jorgensen E. Oral mucosal lesions associated with the wearing of removable dentures. *J. Oral. Pathol.* 1981 10, 65-80.
- Budtz-Jorgensen E, Bertram U. Denture stomatitis. I. The etiology in relation to trauma and infection. *Acta Odontol. Scand.* 1970 28, 71-92.
- Bürgers R, Hahnel S, Reichert TE, Rosentritt M, Behr M, Gerlach T, Handel G, Gosau M. Adhesion of *Candida albicans* to various dental implant surfaces and the influence of salivary pellicle proteins. *Acta Biomater.* 2010 6, 2307-2313.
- Coco BJ, Bagg J, Cross LJ et al. Mixed *Candida albicans* and *Candida glabrata* populations associated with the pathogenesis of denture stomatitis. *Oral Microbiol Immunol.* 2008 23, 377-383.
- Coleman DC, Sullivan DJ, Bennett DE, Moran GP, Barry HJ, Shanley DB. Candidiasis: the emergence of a novel species, *Candida dubliniensis*. *AIDS.* 1997 11, 557-567.
- Coulthwaite L, Verran J. Potential pathogenic aspects of denture plaque. *Br J Biomed Sci.* 2007 64, 180-189.
- Darwazeh AM, Al-Refai S, Al-Mojaiwel S. Isolation of *Candida* species from the oral cavity and fingertips of complete denture wearers. *J. Prosthet Dent.* 2001 86, 420-423.
- Davies AN, Brailsford S, Broadley K, Beighton D. Oral yeast carriage in patients with advanced cancer. *Oral Microbiol Immunol.* 2002 17, 79-84.
- de Castellucci Barbosa L, Ferreira MR, de Carvalho Calabrich CF et al. Edentulous patients' knowledge of dental hygiene and care of prostheses. *Gerodontology.* 2008 25, 99-106.
- De Visschere LM, Grooten L, Theuniers G, Vanobbergen JN. Oral hygiene of elderly people in long-term care institutions--a cross-sectional study. *Gerodontology.* 2006 23, 195-204.
- Douglas LJ. Medical importance of biofilms in *Candida* infections. *Rev. Iberoam Micol.* 2002 19, 139-143.
- Erik Petersen P.E. *Rapport sur la santé bucco-dentaire dans le monde 2003. Poursuivre l'amélioration de la santé bucco-dentaire au XXIème siècle – l'approche du Programme OMS de santé bucco-dentaire.* Programme santé bucco-dentaire Prévention des maladies non transmissibles et promotion de la santé Organisation mondiale de la Santé Genève, Suisse. 2003.
- Ellepola AN, Samaranayake LP. Adjunctive use of chlorhexidine in oral candidoses: a review. *Oral. Dis.* 2001 7, 11-17.
- Felton D, Cooper L, Duquum I, Minsley G, Guckes A, Haug S, Meredith P, Solie C, Avery D, Deal Chandler N; American College of Prosthodontists. Evidence-based guidelines for the care and maintenance of complete dentures: a publication of the American College of Prosthodontists. *J. Prosthodont.* (in press, accepted 2011 Feb. 20 Suppl 1:S1-S12).
- Filoché SK, Soma K, Sissons CH. Antimicrobial effects of essential oils in combination with chlorhexidine digluconate. *Oral. Microbiol. Immunol.* 2005 20, 221-225.
- Fotos PG, Vincent SD, Hellstein JW. Oral candidosis. Clinical, historical, and therapeutic features of 100 cases. *Oral. Surg. Oral. Med. Oral. Pathol.* 1992 74, 41-49.

- Gagliardi DI, Slade GD, Sanders AE. Impact of dental care on oral health-related quality of life and treatment goals among elderly adults. *Aust. Dent. J.* 2008 53, 26-33.
- Gasparoto TH, Dionisio TJ, de Oliveira CE, Porto VC, Gelani V, Santos CF, Campanelli AP, Lara VS. Isolation of *Candida dubliniensis* from denture wearers. *J Med Microbiol.* 2009 58, 959-962.
- Gendreau L, Loewy ZG. Epidemiology and etiology of denture stomatitis. *J. Prosthodont.* 2011 20, 251-60.
- Glass RT, Conrad RS, Bullard JW, Goodson LB, Mehta N, Lech SJ, Loewy ZG. Evaluation of cleansing methods for previously worn prostheses. *Compend Contin. Educ. Dent.* 2011 32, 68-73.
- Gonsalves WC, Wrightson AS, Henry RG. Common oral conditions in older persons. *Am Fam Physician.* 2001 78, 845-52.
- Hsu LY, Minah GE, Peterson DE, Wingard JR, Merz WG, Altomonte V, Tylanda CA. Coaggregation of oral *Candida* isolates with bacteria from bone marrow transplant recipients. *J. Clin. Microbiol.* 1990 28, 2621-2626.
- Izumida FE, Ribeiro RC, Giampaolo ET, Machado AL, Pavarina AC, Vergani CE. Effect of microwave disinfection on the surface roughness of three denture base resins after tooth brushing. *Gerodontology.* (in press, accepted: 2011 Feb 23.)
- Jose A, Coco BJ, Milligan S, Young B, Lappin DF, Bagg J, Murray C, Ramage G. Reducing the incidence of denture stomatitis: are denture cleansers sufficient? *J. Prosthodont.* 2010 19, 252-257.
- Ishikawa A, Yoneyama T, Hirota K, Miyake Y, Miyatake K. Professional oral health care reduces the number of oropharyngeal bacteria. *J. Dent Res.* 2008 87, 594-598.
- Kanli A, Demirel F, Sezgin Y. Oral candidosis, denture cleanliness and hygiene habits in an elderly population. *Aging Clin. Exp. Res.* 2005 17, 502-507.
- Klotz SA, Drutz DJ, Zajic JE. Factors governing adherence of *Candida* species to plastic surfaces. *Infect Immun.* 1985 50, 97-101.
- Kossioni AE. The prevalence of denture stomatitis and its predisposing conditions in an older Greek population. *Gerodontology.* (in press, accepted: 2010 Jan 14.)
- Krcmery Jr. VV, Krupova I, Mateicka F, Jurga L, Sulcova M, Spanik S, Kunová A, Novotny J. *Candida glabrata* fungemia in a tertiary cancer institution in Slovakia. *J Infect Chemother.* 1999 5, 163-167.
- Kulak-Ozkan Y, Kazazoglu E, Arikan A. Oral hygiene habits, denture cleanliness, presence of yeasts and stomatitis in elderly people. *J. Oral. Rehabil.* 2002 29, 300-304.
- Li L, Redding S, Dongari-Bagtzoglou A. *Candida glabrata*: an emerging oral opportunistic pathogen. *J. Dent Res.* 2007 86, 204-215.
- Lukac J, Mravak-Stipetić M, Knezević M, Vrcek J, Sistig S, Ledinsky M, Kusić Z. Phagocytic functions of salivary neutrophils in oral mucous membrane diseases. *J. Oral. Pathol. Med.* 2003 32, 271-274.
- Luo G, Samaranayake LP. *Candida glabrata*, an emerging fungal pathogen, exhibits superior relative cell surface hydrophobicity and adhesion to denture acrylic surfaces compared with *Candida albicans*. *APMIS.* 2002 110, 601-610.
- MacEntee MI, Glick N, Stolar E. Age, gender, dentures and oral mucosal disorders. *Oral. Dis.* 1998 4, 32-36.
- Machado FC, de Souza IP, Portela MB, de Araújo Soares RM, Freitas-Fernandes LB, Castro GF. Use of chlorhexidine gel (0.2%) to control gingivitis and *Candida* species

- colonization in human immunodeficiency virus-infected children: a pilot study. *Pediatr Dent*. 2011 33, 153-157.
- Machado AL, Giampaolo ET, Pavarina AC, Jorge JH, Vergani CE. Surface roughness of denture base and reline materials after disinfection by immersion in chlorhexidine or microwave irradiation. *Gerodontology*. (in press, accepted: 2011 May 27).
- Machado AG, Komiyama EY, Santos SS, Jorge AO, Brighenti FL, Koga-Ito CY. In vitro adherence of *Candida albicans* isolated from patients with chronic periodontitis. *J. Appl. Oral. Sci.* (in press, accepted: 2011 Jun 24.)
- Madrid C, Abarca M, Pop S et al. Dents et tissues de soutien - partie I: Une denture fonctionnelle se compose d'environ vingt dents. *Geriatric Pratique*. 2008 2, 32-36.
- Madrid C, Korsvold T, Durussel C, Bille J, Abarca M. Candidal colonization of dentures in elderly population (abstract and oral presentation). *88th General Session and Exhibition of the IADR*. 2010 Barcelona, Spain.
- Martinez M, López-Ribot JL, Kirkpatrick WR, Coco BJ, Bachmann SP, Patterson TF. Replacement of *Candida albicans* with *C. dubliniensis* in human immunodeficiency virus-infected patients with oropharyngeal candidiasis treated with fluconazole. *J. Clin. Microbiol*. 2002 40, 3135-3139.
- McCarthy GM, Mackie ID, Koval J, Sandhu HS, Daley TD. Factors associated with increased frequency of HIV-related oral candidiasis. *J. Oral. Pathol. Med*. 1991 20, 332-336.
- Mima EG, Pavarina AC, da Silveira Vargas F, Giampaolo ET, Machado AL, Vergani CE. Effectiveness of chlorhexidine on the disinfection of complete dentures colonized with fluconazole-resistant *Candida albicans*: in vitro study. *Mycoses*. (in press, accepted: 2011 May 23.)
- Montero A, Romero J, Vargas JA, Regueiro CA, Sánchez-Aloz G, De Prados F, De la Torre A, Aragón G. *Candida* infection of cerebrospinal fluid shunt devices: report of two cases and review of the literature. *Acta Neurochir (Wien)*. 2000 142, 67-74.
- Nevalainen MJ, Narhi TO, Ainamo A. Oral mucosal lesions and oral hygiene habits in the home-living elderly. *J. Oral. Rehabil*. 1997 24, 332-337.
- Newton AV. Denture sore mouth: a possible etiology. *Br Dent J*. 1962 112, 357-360.
- Nikawa H, Hamada T, Yamamoto T. Denture plaque--past and recent concerns. *J. Dent*. 1998 26, 299-304.
- Odds FC. *Candida and candidosis. A review and bibliography (2nd edition)*. London: Baillière Tindall, 1988: 42-59.
- Paillaud E, Merlier I, Dupeyron C, Scherman E, Poupon J, Bories PN. Oral candidiasis and nutritional deficiencies in elderly hospitalized patients. *Br. J. Nutr*. 2004 92, 861-867.
- Park SE, Periathamby AR, Loza JC. Effect of surface-charged poly(methylmethacrylate) on the adhesion of *Candida albicans*. *J. Prosthodont*. 2003 12, 249-254.
- Pereira-Cenci T, Del Bel Cury AA, Crielaard W et al. Development of *Candida*-associated denture stomatitis: new insights. *J. Appl. Oral. Sci*. 2008 162, 86-94.
- Perezous LF, Flaitz CM, Goldschmidt ME, Engelmeier RL. Colonization of *Candida* species in denture wearers with emphasis on HIV infection: a literature review. *J. Prosthet Dent*. 2005 93, 288-293.
- Phelan DM, Osmon DR, Keating MR, Hanssen AD. Delayed reimplantation arthroplasty for candidal prosthetic joint infection: a report of 4 cases and review of the literature. *Clin Infect Dis*. 2002 34, 930-938.

- Quirynen M, Marechal M, Busscher HJ et al. The influence of surface free energy and surface roughness on early plaque formation. An in vivo study in man. *J. Clin. Periodontol.* 1990 17, 138-144.
- Radford DR, Sweet SP, Challacombe SJ, Walter JD. Adherence of *Candida albicans* to denture-base materials with different surface finishes. *J. Dent.* 1998 26, 577-583.
- Ramage G, López-Ribot JL. Techniques for antifungal susceptibility testing of *Candida albicans* biofilms. *Methods Mol. Med.* 2005 118, 71-79.
- Ramage G, Martínez JP, López-Ribot JL. *Candida* biofilms on implanted biomaterials: a clinically significant problem. *FEMS Yeast Res.* 2006 6, 979-86.
- Redding SW. The role of yeasts other than *Candida albicans* in oropharyngeal candidiasis. *Curr. Opin. Infect Dis.* 2001 14, 673-677.
- Redding SW, Zellars RC, Kirkpatrick WR, McAtee RK, Caceres MA, Fothergill AW, Lopez-Ribot JL, Bailey CW, Rinaldi MG, Patterson TF. Epidemiology of oropharyngeal *Candida* colonization and infection in patients receiving radiation for head and neck cancer. *J Clin Microbiol.* 1999 37, 3896-3900.
- Ribeiro DG, Pavarina AC, Dovigo LN, de Oliveira Mima EG, Machado AL, Bagnato VS, Vergani CE. Photodynamic inactivation of microorganisms present on complete dentures. A clinical investigation : Photodynamic disinfection of complete dentures. *Lasers Med. Sci.* (in press, accepted: 2011 Apr 12.)
- Samaranayake LP. Oral mycoses in HIV infection. *Oral. Surg. Oral. Med. Oral. Pathol.* 1992 73, 171-80.
- Samaranayake YH, Samaranayake LP. *Candida krusei*: biology, epidemiology, pathogenicity and clinical manifestations of an emerging pathogen. *J. Med. Microbiol.* 1994 41, 295-310.
- Samaranayake YH, Samaranayake LP. Experimental oral candidiasis in animal models. *Clin Microbiol Rev.* 2001 14, 398-429.
- Schröder AK, Rink L. Neutrophil immunity of the elderly. *Mech. Ageing Dev.* 2003 124, 419-425.
- Scully C, McCarthy G. Management of oral health in persons with HIV infection. *Oral. Surg. Oral. Med. Oral. Pathol.* 1992 73, 215-225.
- Senna PM, Da Silva WJ, Del Bel Cury AA. Denture disinfection by microwave energy: influence of *Candida albicans* biofilm. *Gerodontology.* (in press, accepted: 2010 Nov 17.)
- Sheiham A, Steele JG, Marcenes W, Lowe C, Finch S, Bates CJ, Prentice A, Walls AW. The relationship among dental status, nutrient intake, and nutritional status in older people. *J. Dent Res.* 2001 80, 408-413.
- Shoham S, Levitz SM. The immune response to fungal infections. *Br. J. Haematol.* 2005 129, 569-582.
- Shulman JD, Rivera-Hidalgo F, Beach MM. Risk factors associated with denture stomatitis in the United States. *J. Oral. Pathol. Med.* 2005 34, 340-346.
- Sullivan D, Coleman D. *Candida dubliniensis*: characteristics and identification. *J. Clin. Microbiol.* 1998 36, 329-334.
- Sumi Y, Miura H, Sunakawa M, Michiwaki Y, Sakagami N. Colonization of denture plaque by respiratory pathogens in dependent elderly. *Gerodontology.* 2002 19, 25-29.
- Sumi Y, Kagami H, Ohtsuka Y. High correlation between the bacterial species in denture plaque and pharyngeal microflora. *Gerodontology.* 2003 20, 84-87.

- Sumi Y, Kagami H, Ohtsuka Y, Kakinoki Y, Haruguchi Y, Miyamoto H. High correlation between the bacterial species in denture plaque and pharyngeal microflora. *Gerodontology*. 2003 20, 84-87.
- Sumi Y, Miura H, Michiwaki Y, Nagaosa S, Nagaya M. Colonization of dental plaque by respiratory pathogens in dependent elderly. *Arch. Gerontol. Geriatr*. 2007 44, 119-124.
- Tawara Y, Honma K, Naito Y. Methicillin-resistant *Staphylococcus aureus* and *Candida albicans* on denture surfaces. *Bull Tokyo Dent Coll*. 1996 37, 119-128.
- Thompson GR 3rd, Patel PK, Kirkpatrick WR, Westbrook SD, Berg D, Erlandsen J, Redding SW, Patterson TF. Oropharyngeal candidiasis in the era of antiretroviral therapy. *Oral. Surg. Oral. Med. Oral. Pathol. Oral. Radiol. Endod*. 2010 109, 488-495.
- Uludamar A, Ozkan YK, Kadir T, Ceyhan I. In vivo efficacy of alkalineperoxide tablets and mouthwashes on *Candida albicans* in patients with denture stomatitis. *J. Appl. Oral. Sci*. 2010 18, 291-296.
- Valério HM, Weikert-Oliveira Rde C, Resende MA. Differentiation of *Candida* species obtained from nosocomial candidemia using RAPD-PCR technique. *Rev. Soc. Bras. Med. Trop*. 2006 39, 174-178.
- Vanden Abbeele A, de Meel H, Ahariz M, Perraudin JP, Beyer I, Courtois P. Denture contamination by yeasts in the elderly. *Gerodontology*. 2008 25, 222-228.
- Webb BC, Thomas CJ, Willcox MD, Harty DW, Knox KW. *Candida*-associated denture stomatitis. Aetiology and management: a review. Part 1. Factors influencing distribution of *Candida* species in the oral cavity. *Aust. Dent. J*. 1998 43, 45-50.
- Wilkieson C, Samaranayake LP, MacFarlane TW, Lamey PJ, MacKenzie D. Oral candidosis in the elderly in long term hospital care. *J. Oral. Pathol. Med*. 1991 20,13-16.
- Williams DW, Kuriyama T, Silva S, Malic S, Lewis MA. *Candida* biofilms and oral candidosis: treatment and prevention. *Periodontol*. 2000. 2011 55, 250-265.
- Willis AM, Coulter WA, Sullivan DJ, Coleman DC, Hayes JR, Bell PM, Lamey PJ. Isolation of *C. dubliniensis* from insulin-using diabetes mellitus patients. *J. Oral. Pathol. Med*. 2000 29, 86-90.
- Wingard JR. Importance of *Candida* species other than *C. albicans* as pathogens in oncology patients. *Clin. Infect. Dis*. 1995 20, 115-25.

Chapter XI

Methods for the Treatment of Denture Induced Psychogenic Symptoms

***Anita Beck^{*1,2}, Gábor Varga², Péter Hermann¹,
Gábor Fábián³, Pál Fejérdy¹ and Tibor Károly Fábián¹***

¹Clinic of Prosthetic Dentistry,

²Department of Oral Biology,

³Clinic of Pediatric Dentistry and Orthodontics

Semmelweis University Budapest, Faculty of Dentistry, Budapest, Hungary, EU

Abstract

Psychogenic denture intolerance (PDI) patients suffering from several denture induced psychogenic symptoms refuse to accept psychological background of their symptoms frequently. Instead of psychiatrists or psychotherapists, first they visit dentist and insist on the somatic origin of their symptoms. Since most of these patients refuse a referral to psychiatrist and/or psychotherapist, an initial psychosomatic therapy is needed, which is a scope of dental profession's duty. Initiation of a palliative dental care and gradual escalation of any kind of mind-body therapy (as "basic therapeutics" for psychosomatic disorders) are "cornerstones" of such dental psychosomatic interventions. Introduction of any method of complementary/alternative medicine may also be useful especially for the prevention of relapse. To carry out efficient palliative dental care useless repetition of previous dental treatment (or any further invasive dental treatment) should be avoided, certain psychotherapeutic approaches should be introduced and several other treatment modalities like physiotherapy, medication, medicinal herb therapy or acupuncture should also be administered. For mind-body therapies, relaxation, hypnosis, self-hypnosis, meditation, photo-acoustic stimulation or biofeedback may be used. For complementary and alternative medicine prayer, breathing exercises, Tai-Chi and Qigong, Yoga or several other therapies (i.e. sleep deprivation, vigil, fasting therapy etc.) may be considered.

* E-mail address: becka@fok.usn.hu.

Introduction

Psychogenic denture intolerance (PDI) patients suffering from several denture induced psychogenic symptoms refuse to accept psychological background of their symptoms frequently [Ross et al 1953, Fábíán et al. 2005, Schwichtenberg and Doering 2008, Fábíán and Fejérdy 2010]. Instead of psychiatrists or psychotherapists, PDI patients usually visit dentists and insist on the somatic origin of their symptoms [Ross et al. 1953, Fábíán et al. 2005, Fábíán and Fejérdy 2010]. Major reason behind premised behavior of patients may be that most PDI manifestations (notwithstanding that they are primarily of psychogenic origin) are coupled with somatic type reactions like several pain symptoms, neuromuscular symptoms, TMJ manifestations, occlusal discomfort, allergic/irritative reactions, salivation problems (etc.) [Fábíán and Fejérdy 2010].

Since most of these patients refuse a referral to psychiatrist and/or psychotherapist, an initial psychosomatic therapy is needed, which is a scope of dental profession's duty [Moulton et al. 1957, Pomp 1974, Fábíán and Fejérdy 2010]. Initiation of a palliative dental care and gradual escalation of any mind-body therapy are "cornerstones" of such dental psychosomatic interventions. Introduction of any method of complementary/alternative medicine may also be useful.

To carry out efficient palliative dental care useless repetition of previous dental treatment (or any further invasive dental treatment) should be avoided [Brodine and Hartshorn 2004, Sarlani et al. 2005/b, Reewes and Merrill 2007, Baad-Hansen 2008], certain psychotherapeutic approaches should be introduced [Fábíán and Fejérdy 2010] and several other treatment modalities like physiotherapy, medication, medicinal herb therapy or acupuncture should also be administered [Fábíán and Fejérdy 2010]. For making decision that which premised treatment modality would be the best choice to administer in a concrete case, patient's major somatic type complaint should be considered first; and a method found to be efficient in the scientific literature to improve such complaint should be chosen.

Besides palliative dental care, any mind-body therapy should also be introduced and gradually escalated in the course of treatment. Mind-body therapies are "basic therapeutics" for psychosomatic disorders [Iversen 1989, Binder and Bider 1989, Krause 1994, Fábíán et al. 2009, Fábíán in press]. Mind-body therapies usually decrease anxiety and level of perceived stress [Fábíán and Fejérdy 2010, Fábíán in press], and advantageously improve patients' mood, vitality, sleep quality as well as psychoemotional and psychosocial functions [Fábíán and Fejérdy 2010, Fábíán in press]. For this purpose relaxation, hypnosis, self-hypnosis, meditation, photo-acoustic stimulation or biofeedback may be used. Mind-body therapies advantageously improve psychogenic symptoms in most of the cases leading either to a recovery or to an improvement coupled with increased motivation of patients to take part in a definitive psychosomatic therapy including psychiatric/psychotherapeutic treatment.

As mentioned above, introduction of any method of complementary/alternative medicine may also be useful, especially for the prevention of relapse [Fábíán and Fejérdy 2010]. Complementary and alternative medicine (CAM) denotes a wide range of variable therapies, including also treatments with established benefits and few if any side effects. CAM therapies emphasize self-care, which can lead to advantageously decreased load of the much more expensive health care system. For complementary and alternative medicine prayer, breathing

exercises, Tai-Chi and Qigong, Yoga or other therapies (i.e. sleep deprivation, vigil, fasting therapy etc.) may be considered [Fábián and Fejérdy 2010].

Methods for Palliative Dental Care

Psychotherapeutic Approaches

To carry out efficient palliative dental care use of certain techniques of psychotherapy may be needed [Fábián and Fejérdy 2010]. Although in general there is no contraindication of psychotherapy, however level of qualification of the dentist should determine the depth of psychotherapeutic intervention used. Deeper psychotherapeutic interventions targeting psychological structural changes of the personality should be avoided in the absence of full psychotherapeutic qualification. Deeper psychotherapeutic interventions should also be avoided in the case of severe pathologies in the absence of psychiatric support and professional background. For dental psychosomatic therapies client centered approach may be offered as a basic approach, however, certain techniques of cognitive-behavioral therapy, and psychoanalytic therapy may also be used. Client-centered therapy emphasizes the autonomy of clients and the healing effect of the encounter during therapy [Rogers 1965]. It also emphasizes the attitudinal qualities of the therapist such as warmth, empathy and genuineness (congruence) [Rogers 1965, Smith 2000] as well as the ability of therapist to emphatically recognize and reflect ("mirror") the real sense behind the client's communication [Rogers 1965]. Autonomy of client together with the encounter and the attitudinal qualities of the therapist help patient (client) to remove inside obstacles of personality development, and to become their best possible self [Rogers 1965, Smith 2000].

Cognitive-behavioral therapy is highly advantageous combination of behavioral therapy and cognitive therapy. Behavioral therapy is an operationalized approach, which has its roots in classical learning theory and other findings of experimental psychology. It focuses on current determinants of behavior and draws on the principles of learning to develop individual treatment strategies [de Silva 2000, Smith 2000]. The focus in behavior therapy is the presenting problem itself; and it is not assumed that the presenting problem is a manifestation of an underlying primary problem [de Silva 2000, Smith 2000]. The aim of behavior therapy is to modify the problem behavior at a behavioral level [de Silva 2000, Smith 2000]. Cognitive therapy is based on the assumption that, the person's feelings and behavior are determined by the way in which their experiences are processed cognitively [Ellis 1962, Beck 1976]. In this approach, irrational determinants of thoughts (i.e. spontaneously appearing negative thoughts, errors of logical thoughts, and depressive cognitive schemes [Beck 1976]) are assumed as a major cause of several pathopsychological conditions [Ellis 1962, Beck 1976]. Therefore, cognitive therapy aims to decrease premised maladaptive irrational cognitions and to increase adaptive cognitions of patients.

Psychoanalytic therapies are based on various theoretical streams of psychoanalytic theory. Besides the fundamental libido-theory and ego-psychology, object-relationships theory and self-theory are the most important such theories. Major therapeutical interventions of psychoanalysis are analysis of free associations, analysis of resistance, analysis of

transference, analysis of counter-transference and the related interpretations. Although psychoanalysis is a depth-psychological theory, focusing on unconscious psychological processes [Freud 1900] there are several short-term forms of psychoanalytic therapy (referred to as brief dynamic psychotherapy) which are explicitly problem focused and focal, therefore may be used also for dental psychosomatic therapies.

Physiotherapies

Certain techniques of physiotherapy may also be used efficiently for palliative dental care [Fábián and Fejérdy 2010]. However, it should be considered that, there are some contraindications and precautions of physiotherapies. In general, physiotherapy is contraindicated in the case of patients with any severe systemic diseases and with strongly compromised health. In the case of such patients, use of physiotherapy should be avoided in the absence of a clear agreement of the medical professional(s) responsible for the treatment of the systemic disease(s) at issue (and/or responsible for the medical care of strongly compromised health). Similarly, physiotherapy is usually contraindicated in the cases of patients with any hemorrhagic diathesis, precancerosis, and malignancy as well as in the case of patients having any oncological treatment in the anamnesis (at least in the absence of a clear agreement of the oncologist or other responsible medical professional). Importantly, use of any physiotherapy should be strictly avoided near to a nevus. In the absence of clear evidences about the innocuity, several forms of physiotherapy may be contraindicated for patients with pacemaker (or with another implanted electric instruments) as well as in the case of patients with any implants including also dental implants. Similarly, physiotherapeutic treatment should be avoided near to the ear, eye, brain or spinal cord (i.e. temporal or paravertebral use) and in the condilar area during the growth period, in the absence of clear evidence about the innocuity of the chosen treatment modality. (Especially those modalities utilizing electric/electromagnetic impulses or irradiative energy with significant heat generating effect like laser or ultrasound should be considered for such contraindication.) The most frequently used physiotherapies for the treatment of denture induced psychogenic manifestations are massage combined with muscle exercises, occlusal splint therapy, transcutaneous electrical nerve stimulation (TENS), pulsed electromagnetic field therapy (PEMF), and low level laser (soft laser) therapy.

Massage combined with muscle exercises are aiming to re-train the incoordinated masticatory musculature [Boos 1959, Laney 1983, Schulte 1988, Lund et al. 2006] and to improve the metabolic processes of the muscles and other tissues. Massage is a therapeutic friction, stroking, kneading or shaking of a part of the body. Massage aid the return of venous blood, lymph, and catabolites into the main circulation and therefore may reduce muscle pain or other symptoms [Laskin and Block 1986]. Muscle exercises re-train the incoordinated masticatory musculature and also improve the metabolic processes of the muscles. For most cases, three patterns of sequential movements of the mandible may be used 4 times daily for periods of 2 to 3 minutes as follows [Boos 1959, Laney 1983]: (1) maximal opening position held for 30 seconds and followed by relaxation (without tooth contact); (2) jaw movements to the right (and subsequently to the left) in a slow continuous stretch held for 30 seconds and

followed by returning to the rest position; (3) the mandible is protruded held for 30 seconds and then retruded to the resting posture.

Ultrasound treatment produces vibrations within the tissue that cause particle collision and the release of energy resulting in the production of heat [Laskin and Block 1986]. Therefore, ultrasonic treatment induces a combined deep effect of vibration ("micromassage") and deep heat effect (to a dept of 4 - 5 cm) [Bell 1969/b, Laskin and Block 1986]. Ultrasound therapy may be used to reduce muscle tension and to improve myofascial pain symptoms. Ultrasound treatment also increases tissue elasticity and improve circulation. The sound head is used over the involved muscles (i.e. masseter or temporal muscles) and should be moved slowly to avoid excessive heat build up [Laskin and Block 1986]. Treatments last 10 to 15 minutes, and can be applied twice a day for 1 to 2 weeks [Laskin and Block 1986]. Ultrasound should not be applied over the eye, over the ear, over an area of acute inflammation [Bell 1969/b, Laskin and Block 1986] and over metallic implants [Laskin and Block 1986]; and great care should be taken to avoid heat build up when ultrasound is used near the brain and spinal cord (i.e. temporal or paravertebral use). Ultrasound therapy is also contraindicated in patients with cardiac pacemaker, vascular insufficiency and malignancy [Laskin and Block 1986]. Use of ultrasound should also be avoided in the condylar area during the growth period [Bell 1969/b]

Transcutaneous electrical nerve stimulation (TENS) refers to commercially available devices that apply electric impulses to the peripheral nerves via electrodes placed on the skin [Bishop 1986, Curcio et al. 1987]. Although electrodes of TENS devices were originally placed on the skin, in dentistry "intraoral TENS" (intraoral electrostimulation therapy) utilizing electrodes placed intraorally may also be used [Wilder-Smith and Zimmermann 1989, Wilder-Smith 1990]. Based on the gate-control theory [Melzack and Wall 1965] it is hypothesized that stimulation of the branches of the trigeminal nerve due to TENS creates an inhibitory effect on the trigeminal nucleus and thereby reduces awareness of pain and helps (indirectly) to induce muscle relaxation. It is also hypothesized that, TENS induces release of endorphins, norepinephrine and serotonin in the central nervous system, which may also be responsible for the clinical effects [Black 1986, Hochman 1988]. Further, TENS increases circulation in the areas where electrodes are placed and of those muscles, which belong to the stimulated nerve [Hochman 1988, Laskin and Block 1986]; and therefore reduces muscle pain and increases muscles' resistance to fatigue [Laskin and Block 1986]. There is also a rather significant placebo effect coupled with TENS therapy [Gold et al. 1983, Dahlström 1992, Green and Laskin 2000]. Patients are usually treated by placing the electrode directly over the area of most discomfort for 30 minutes daily [Laskin and Block 1986]. TENS is contraindicated for certain patients including pregnant patients and pacemaker patients as well as for patients with cerebral convulsive disorders (possibility of initiating seizure), with pathologic hypotension (TENS lowers blood pressure) and with cerebral vascular disorders (blood flow increases in the areas where electrodes are applied) [Hochman 1988].

Occlusal splints are also frequently used. Reflex splints (occlusal disengagement) trigger mouth opening reflex due to functioning as an "artificial early contact" and as such may be used for reducing stress induced muscle spasm and parafunction activity [Mikami 1977, Laskin and Block 1986]. Reflex splints should be used for short run (at maximum for 2 weeks [Freesmeyer 1995]), because if such a splint were worn too long, a "new series" of muscle

spasm and parafunction could be triggered [Freesmeyer 1995]. Further, there is a danger of elongation (supereruption) of teeth, remodeling of the temporomandibular joint [Griffin 1975], as well as other severe and irreversible complications [Widmalm 1999]. If reflex splints are not worn continuously, but only at night and for 5 to 6 hours during the day [Laskin and Block 1986], reflex splints may be used somewhat longer (at maximum for 4 to 6 weeks) under regular control [Bell 1969/b, Laskin and Block 1986]. Reflex splints may be equilibrated and transformed into equilibrated splints due to filling up the existing space between the molars [Green and Laskin 1972/a], followed by occlusal adjustment until simultaneous tooth contacts are achieved on the complete occlusal coverage of the arch. Such equilibrated splints may be used for harmonizing muscle function [McNamara 1976, Mikami 1977, Dahlström et al. 1982, 1985, Dahlström 1989]. Importantly, the efficiency of splint therapy is expected to be based (at least partially) on placebo effects [Green and Laskin 1972/a, 2000, Laskin and Block 1986, Molin 1999]. However certain kind of splints like repositioning appliances (due to positioning of condyles) clearly exerts certain specific therapeutic effects as well [Green and Laskin 1972/a, Dahlström 1992, Karppinen et al. 1999].

Pulsed electromagnetic field (PEMF) therapy seems to be a highly promising method for psychosomatic dentistry. PEMF induces local vasodilatation [Smith et al. 2004] and increases angiogenesis [Tepper et al. 2004]. PEMF also enhances healing of bone fractures [Gossling et al. 1992, Pienkowski et al. 1992, Inoue et al. 2002], decreases the rate of residual ridge resorption following tooth extraction [Ortman et al. 1992], increases the rate of orthodontic tooth movement and coupled bone deposition [Stark and Sinclair 1987, Darendeliler et al. 1995], and improves bone formation in bone defects and around dental implants (osteointegration of implants) [Matsumoto et al. 2000]. Osteoporotic symptoms and symptoms of osteoarthritis patients may also be improved using PEMF. Besides the important bone-related effects, PEMF also improves tendon inflammation and returns to histological normality as well [Binder et al. 1984, Lee et al. 1997]. PEMF also improves wound healing of the gastrointestinal mucosa [Mentes et al. 1996] and the skin [Patino et al. 1996, Scardino et al. 1998]. Similarly, ulcerations may also be improved due to PEMF therapy [Stiller et al. 1992]. PEMF also improves regeneration of injured nerve [Raji 1984, Zienowicz et al. 1991, Kanje et al. 1993], and improves clinical symptoms (i.e. pain, decrease of conductive function, decrease of reflex excitability) of several neuropathic conditions [Musaev et al. 2003, Weintraub and Cole 2004]. PEMF may also be used for acupuncture therapy. Patient are usually treated by placing the applicator directly over the area of most discomfort for 10-15 (up to 30-40) minutes at least once (or may be twice or three times) a week. There is no specific contraindication of PEMF therapy, but general contraindications should be considered also for PEMF therapy.

Low level laser therapy (LLLT; also referred to as soft laser therapy) is a treatment with athermic low power lasers. There are many different types of low level laser (LLL) may be used for LLLT including either pulsed or continuous LLL, and with wavelength in both visible and invisible range [Khullar et al. 1995, Kimura et al. 2000]. In general, increase of cell metabolism, collagen synthesis and activity of immune cells are the most important pathways behind the efficiency of low level laser therapy [Qadri et al. 2005]. LLLT also inducing long-lasting expression of HSP70/HSPA type stress proteins [Souil et al. 2001]. There is also a prominent placebo effect coupled with low level laser therapy [Kimura et al.

2000, Payer et al. 2005]. LLLT may be used advantageously to control periodontal inflammation [Qadri et al. 2005, Schwarz et al. 2008] and more severe inflammatory and/or necrotizing bone pathologies [Vescovi et al. 2008]. LLLT may also be used to improve implant-tissue interaction in the bone and at the implant-soft tissue interface [Khadra et al. 2005/a,b, Schwarz et al. 2005]. LLLT may also be used to facilitate orthodontic tooth movement and related alveolar bone remodeling as well [Kawasaki and Shimizu 2000, Fujita et al. 2008]. LLLT may also facilitate bone metabolism during healing of bone defects. Besides its advantageous effect in the bone, LLLT improves myofascial pain [de Medeiros et al. 2005, Shirani et al. in press] and temporomandibular joint pain [Dahlström 1992, Mazzetto et al. 2007]. LLLT may also be used to improve functional recovery of injured peripheral nerves [Khullar et al. 1995, Ozen et al. 2006]. LLLT also accelerates oral wound healing [Amorim et al. 2006, Ozcelik et al. 2008] as well as healing of oral ulcerations [Navarro et al. 2007]. Low level lasers may also be used for acupuncture therapy.

Medicamentous Therapies

Medicamentous therapies are frequently used for dental psychosomatic therapies. The most frequently used drugs are pain killers and non-steroidal anti-inflammatory drugs, muscle relaxants, anxiolytics, and antidepressants. There are various specific contraindications and possible side effects of medicamentous therapies, which should be considered very carefully [Bell 1969/b, Laskin and Block 1986, Friedlander et al. 2004]. Utilizing pharmacological databases and an up-to-date pharmacological knowledge, dentists should be familiar with the potential hazards before prescribing a medication [Bell 1969/b, Laskin and Block 1986, Friedlander et al. 2004]. Considering premised precautions, short run (1 to 2 weeks, or maximum 3 to 4 weeks) administration of several medicaments can be carried out by the dentist in the frame of initial psychosomatic therapy [Bell 1969/b, Miyamoto and Ziccardi 1998, Laskin and Block 1986, Toyofuku and Kikuta 2006]. However, long-run medicamentous treatment of long-standing symptoms is clearly a scope of medical professionals such as psychiatrists, neurologists, rheumatologists, internists, family doctors (etc.). Therefore, if long-run medication is needed because of any reasons, it is necessary to place the patient under physicians' care [Laskin and Block 1986].

Pain killers and non-steroidal anti-inflammatory drugs (NSAIDs) are used primarily for the initial therapy of not yet chronic cases, because there is frequently no significant improvement of chronic orofacial psychosomatic symptoms after treatment with pain killers and non-steroidal anti-inflammatory drugs (NSAIDs) [Gellrich et al. 2002]. Since most of the orofacial pain symptoms are also (at least partially) of inflammatory origin, in most cases NSAIDs (instead of exclusive pain killers) are used for such purposes. Importantly, NSAIDs can also be applied locally [King 1988]. There are numerous NSAID gels and plasters available for local application, which may be used advantageously when applied onto the skin surface over the painful area. Local application may prevent drug tolerance and dependence which may appear following per os application of these drugs [Israel and Scrivani 2000].

Muscle relaxants may be used, because spasm produces pain that in turn produces more spasm in many cases [Laskin and Block 1986]. Thus, muscle relaxants may be used to

interrupt this vicious cycle, especially when used in conjunction with a pain-relieving medication [Laskin and Block 1986]. Importantly, certain anxiolytics (especially benzodiazepines) also exert significant myorelaxant properties [Chaco, 1973, Laskin and Block 1986, Dahlström 1989, 1992]. Since tranquilizing (anxiolytic) properties can be beneficial in relieving muscle tension too [Laskin and Block 1986], premised anxiolytics (benzodiazepines) are the drugs most often prescribed as muscle relaxants [Laskin and Block 1986, Dahlström 1992]. Again, benzodiazepines should also be taken in conjunction with an analgesic for the best effect [Laskin and Block 1986]. Interestingly, drugs with only muscle relaxant properties (i.e. without any anxiolytic effects) seem to be not as successful as premised anxiolytics (benzodiazepines) for the treatment of myofascial pain [Laskin and Block 1986].

Anxiolytics (especially benzodiazepines) may be used as an adjunct for the initial treatment of patients with high background anxiety level [Bell 1969/b]. Because of their significant antidepressive properties [Köhler 2005] benzodiazepines may also be used advantageously for patients with depression [Bell 1969/a], although cases with severe depression can not be managed with the use of benzodiazepines by itself [Köhler 2005]. Based on their myorelaxant effect, benzodiazepines may also be used advantageously as an adjunct to the initial therapy of patients with myofascial pain [Bell 1969/b, Green and Laskin 1972/b, Ryan et al. 1985, Laskin and Block 1986] and denture related (denture induced neuromuscular dysfunction related) functional TMD problems [Bell 1969/b, Chaco 1973, Dahlström 1989]. Benzodiazepines may also be used for the treatment of burning mouth syndrome [Sarhani et al. 2005/b].

Antidepressants can be used for pain reduction even if there are no current symptoms of depression [Maina et al. 2003, Derra and Egle 2003]; because they also exert a significant direct analgesic effect which is likely to be independent from the antidepressive effect [Derra and Egle 2003]. In contrast to antidepressive effect (appearing earliest after 2-4 weeks at best [Köhler 2005]) premised analgesic effect appears after 3-7 days [Miyamoto and Ziccardi 1998, Derra and Egle 2003]; and importantly, low doses are also sufficient to reach the premised analgesic effect [Miyamoto and Ziccardi 1998, Derra and Egle 2003]. For such low dosage administration, especially serotonin-norepinephrine reuptake inhibitors may be useful [Miyamoto and Ziccardi 1998, Derra and Egle 2003], because their direct analgesic effect seems to be stronger comparing to other groups of antidepressants [Miyamoto and Ziccardi 1998, Derra and Egle 2003]. Similarly, serotonin-norepinephrine reuptake inhibitors exert a direct effect on the somatosensory system as well; which is also likely to be independent from the antidepressive effect [Toyofuku and Kikuta 2006] and can be utilized efficiently for treating occlusal dysesthesia [Toyofuku and Kikuta 2006]. Similarly to pain treatment, low doses are likely to be sufficient to reach this kind of somatosensory effect too [Toyofuku and Kikuta 2006], however, this effect likely appears somewhat later (after 2-4 weeks) [Toyofuku and Kikuta 2006]. Although a definitive medicamentous treatment of affective disorders (including depression) is clearly a scope of psychiatrist; short run (1 to 2 weeks, or maximum 3 to 4 weeks) administration of low doses antidepressants (see above) can be carried out also by the dentist in the frame of initial psychosomatic therapy [Miyamoto and Ziccardi 1998, Laskin and Block 1986, Toyofuku and Kikuta 2006]. Especially pain symptoms including

myofascial pain, atypical facial pain, atypical odontalgia, phantom tooth pain, burning mouth syndrome as well as occlusal dysesthesia may be improved in this way.

Medicinal Herb Therapy

Medicinal herb therapy is an important supplemental modality of psychosomatic dental therapy. Proper use of medicinal herbs may be utilized for such therapeutic purposes efficiently. However, potential health hazards should be avoided. Therefore, great care should be taken especially in the case of patients with severe systemic diseases as well as during pregnancy and lactation. Similarly, great care should also be taken, when long run administration of medicinal herbs is needed. In such cases consultation with (or referral to) an internist and/or family doctor is highly recommended before starting medicinal herb therapy [Fábián and Fejérdy 2010]. It should be also considered that, medicinal herb therapy is frequently used for health enhancement by patients without any professional control; and unfortunately, they also use medicinal herbs during pregnancy and lactation or under severe systemic diseases, which may cause significant health hazards [Fábián and Fejérdy 2010]. Concurrent use of medicinal herbs and pharmaceutical drugs without any control of medical or dental professionals is relatively common and the majority of herbal medicine users are not aware of potential adverse effects [Zhang et al. 2008]. Therefore, dentist should carefully collect information about the patients' medicinal herb related usage. Besides specific contraindications, potential health hazards including possible interactions with medications [Segelman et al. 1976], risk of abuse of medicinal herbs [Siegel 1979] and risk of herbal intoxication [Siegel 1976] should also be considered very carefully before starting medicinal herb therapy.

A principled multi-target strategy of medicinal herb therapy may be advantageously used [Ginsburg et al. 2008]; however the use of an unprincipled combination of medicinal herbs should be strictly avoided. To avoid any potential adverse effects, dentist should administer medicinal herb therapy for a limited time span (few months) only. Following this period the therapeutic administration (at least two cups per day and/or use of medicinal herb extracts) of medicinal herbs should be either terminated or surrendered to the family doctor (or other professionals like internist, psychiatrist etc.). In the case of pregnancy, lactation, and any severe compromised health conditions, a consultation with the family doctor (or any other professionals responsible for the treatment of the condition at issue) is highly recommended before starting any medicinal herb therapy. For the treatment of psychogenic symptoms administration of Black Cohosh, Chastetree, Gingko, Ginseng and St John's wort as well as Cocoa, Coffee and Tea may be considered primarily.

Black cohosh (*Cimicifuga racemosa*) constituents likely act on serotonin receptors and may relieve hot flashes of women and improve mood through a serotonergic effect [Burdette et al. 2003, Geller and Studee 2005]. Improvement of sleeping, fatigue and abnormal sweating [Pockaj et al. 2004] could also be based on premised serotonergic (antidepressant) effect. Black cohosh may also exert bone-protective effects. The combination of black cohosh with St John's wort (see below) was found efficient also in psychovegetative disorders, likely via a synergistic effect [Liske 1998]. Use of Black cohosh is contraindicated in pregnancy, lactation, failed liver function and may be breast cancer.

Chastetree (*Vitex agnus-castus*) is a progesterone-like substance containing [Brown 1997, Geller and Studee 2005] plant approved for the treatment of premenstrual syndrome (PMS), breast tenderness and irregularities of menstrual cycle [Geller and Studee 2005]. Similarly, chastetree may improve peri- and postmenopausal symptoms including emotional problems and hot flashes [Lucks 2003]. The use of this plant should be avoided during pregnancy and lactation [Daniele et al. 2005] and interference of chastetree with dopaminergic antagonists may also be expected [Daniele et al. 2005].

Ginkgo (*Ginkgo biloba*) increases blood flow especially to the brain, increases uptake of glucose by brain cells and improve transmission of nerve signals [Geller and Studee 2005]. Ginkgo is approved for use of cerebral insufficiency, vertigo, tinnitus and also for peripheral vascular diseases like Raynaud's syndrome [Kleinen and Knipschild 1992, Geller and Studee 2005]. Ginkgo may also be used to improve memory function and dementia.

Ginseng (*Panax ginseng*) is used as a tonic for invigoration and fortification in times of fatigue, for improvement of declining capacity for work or concentration, and for coping with psychosocial stress [Geller and Studee 2005]. It may also be used to improve depression, insomnia, well being [Tode et al. 1999, Wiklund et al. 1999] and unexplained chronic fatigue [Bentler et al. 2005]. Ginseng also enhances the anti-allergic effect of green tea [Maeda-Yamamoto et al. 2007]. However, in the case of malignancies use of ginseng could be contraindicated [Geller and Studee 2005].

St John's wort (*Hypericum perforatum*) seems to be effective to improve mild to moderate depressive symptoms (but not major depression) [Linde et al. 1996, 2005, Volz 2005], as well as fatigue [Volz 2005], atypical depression [Volz 2005], several somatoform disorders [Volz 2005] and likely also anxiety [Volz 2005]. A synergistic effect of St John's wort with black cohosh (see above) in the case of psychovegetative disorders is also very likely [Liske 1998, Geller and Studee 2005]. Climacteric symptoms including several psychological and psychosomatic symptoms and sexual well-being may also be improved efficiently with the use of St John's wort [Grube et al. 1999]. However, great care should be taken with herb-drug interactions because St John's wort is a potent ligand for pregnane X receptor [Moore et al. 2000, Carlson et al. 2008] and consequently increases cytochrome P450 dependent hepatic drug metabolism. Therefore St John's wort may decrease blood concentration of several drugs including anticoagulants, oral contraceptives, cyclosporin, digoxin etc.

Cocoa contains a variety of active substances which act at the same site in the brain like cannabis [DiTomaso et al. 1996] and amphetamine [Benton and Donohoe 1999]. Cocoa also contains relatively high amount of stimulants like theobromine and caffeine as well as magnesium [Bruinsma and Taren 1999] which improves mood and may improve coping ability with stress. Certain cocoa flavonols also exert anti-inflammatory properties.

Coffee constituents exert several physiological effects including central nervous system stimulation, acute elevation of blood pressure, increase of metabolic rate and diuresis. Other constituents exert rather significant antioxidant activity [Higdon and Frei 2006]. Moderate coffee consumption (up to 3-4 cups in a day) is associated with a reduced risk of depression [Smith 2009], improved cognitive functioning [Smith 2009], improved mood [Robelin and Rogers 1998], improved psychomotor performance [Robelin and Rogers 1998] and decreased fatigue [Smith 2002] all of which are rather advantageous also for psychosomatic therapies.

Tea (*Camellia sinensis*) is a native plant of South Asia, which may be (black tea) or may not be (green tea) “fermented” prior drying. Black tea is rich in caffeine but, the total content of flavonoids is usually lower. Consumption of black tea is associated with a reduced risk of depression [Smith 2009], improved cognitive functioning [Smith 2009], improved mood [Robelin and Rogers 1998], improved psychomotor performance [Robelin and Rogers 1998] and decreased fatigue [Smith 2002]. Green tea is rich in flavonoids, but the caffeine content is much lower than in black tea. Tea flavonoids exert significant anxiolytic, sedative and psychological stress reducing properties [Adachi et al. 2006, Vignes et al. 2006]. Tea flavonoids also induce immunosuppressive alterations on human dendritic cells, decrease the adhesiveness and migration of mast cells and are likely to inhibit T-cell activation. They also inhibit activity and expression of matrix metalloproteinases (agents causing alveolar bone loss) and inhibits osteoclast formation [Yun et al. 2004]. Since tea flavonoids inhibit hepatic cytochrome P450-dependent enzymes [Yang et al. 2000, McKay et al. 2002], possible herb-drug interactions should also be considered especially in the case of green tea.

Acupuncture

Acupuncture is a traditional Chinese practice that attempts to regulate and restore energy balance by stimulating specific acupoints [Whittaker 2004]. Acupuncture therapy may be based on the trigger point method, in which painful trigger points are recognized and stimulated with acupuncture needles and/or with any other acupuncture modalities (i.e. electro acupuncture, laser- or PEMF acupuncture, acupressure, moxibustion etc.) [Baldry 2005, Fábíán and Fejérdy 2010]. Similarly, acupuncture treatment of intraoral painful points (i.e. oral acupuncture) may also be used [Schmid-Schwap et al. 2006]. In this case most painful points are selected by the “very-point method” [Gleditsch 1980] after palpation of the retromolar areas of maxilla and mandible and the vestibulum [Gleditsch 1980, Schmid-Schwap et al. 2006].

Besides trigger point (and/or intraoral “very-point”) based treatment, acupuncture therapy based on the traditional Chinese conception of vital energy (called “Chi” or “Qi”) may also be used [Baldry 2005, Fábíán and Fejérdy 2010]. Traditional acupoint Hegu (LI-4) may be used for the treatment of oral and/or dental pain as well as for the treatment of TMD disorder. Yifeng (SJ-17) may also be used for the treatment of oral and/or dental pain. Chengjiang (REN-24) may be used for control gag reflex and also for the treatment of xerostomia. Jiache (ST-6) and Xiaguan (ST-7) may be used for the treatment of xerostomia and also for the treatment of oral and/or dental pain. Daying (ST-5) and Tinggong (SI-19) may also be used to treat xerostomia. Serious adverse effects following acupuncture are rare [Vachiramon and Wang 2005]. However acupuncture may be contraindicated in the case of patients with certain psychiatric conditions like paranoid psychosis, hypochondria (etc.). Great care should be taken for infection control. Injuries of blood vessels nerves and any other sensitive tissues should also be strictly avoided when needle type acupuncture is used. Specific cautions should also be considered when laser-, electric-, or electromagnetic (PEMF) stimulation are used.

Mind-Body Therapies

Relaxation

Relaxation belongs to the natural behavior repertoire of human, but can also be induced by several methods following a learning process [Tóth and Fábíán 2006, Fábíán and Fejérdy 2010, Fábíán in press]. The most frequently used basic methods are simple eye-closure, standard exercises of Autogenic Training [Schultz 1932], progressive muscle relaxation [Jacobson 1938], and biofeedback assisted relaxation [Ray et al 1979]. Relaxation induces a relaxing feeling, reduced arousal level, and reduced stress. In relation with oral psychosomatic symptoms, relaxation may be used efficiently for the therapy of bruxism, myofascial pain and TMD symptoms of muscular dysfunction origin [Biondi and Picardi 1993, Sarlani et al. 2005/a]. Somatoform chronic pain may also be treated efficiently using progressive muscle relaxation [Kröner-Herwig et al. 1998, Derra 2003, Rehfish and Basler 2004]. Dental fear and dental phobia can also be treated with relaxation based methods [Hammarstrand 1995, Moore et al. 1996]. Relaxation may also be combined advantageously with numerous psychotherapeutic approaches. Accordingly, combination of progressive muscle relaxation with cognitive behavioral therapy and abdominal/diaphragmatic breathing was highly efficient to reduce pain-activity interference, pain intensity, depression, maladaptive pain beliefs, catastrophizing and to improve masticatory jaw function in a randomized controlled trial [Turner et al. 2006].

Hypnosis

Hypnosis may be defined as an altered state of consciousness (ASC) achieved by suggestions. Numerous important theories were developed to explain hypnosis [Orne 1959; Shor 1962, Bányai and Hilgard 1976, Hilgard 1991, Bányai 1991, Crawford and Gruzelier 1992]. However, there is currently consensus only on that the real essence of hypnosis lies in the experienced subjective alteration of consciousness (ASC) [Varga et al. 2001]. In this relation, altered state of consciousness (ASC) can be defined as sudden and transient subjective experience significantly different from those of common everyday experiences. ASC may induce significant changes of most psychological functions including attention, perceptions, sense of time, body image, self image, imagination, fantasy, cognition, emotions, arousal, memory, self-control, suggestibility, identity etc. [Kihlstrom 1984, Pekala and Kumar 1989, Farthing 1992, Halsband et al. 2009]. Importantly, significant pain relieving effect also occur following induction of hypnotic state; which can be influenced (i.e. increased or decreased) significantly using specific suggestions [Meier et al. 1993; Rainville et al. 1997, 1999; Fábíán and Fábíán 1998; Gáspár et al. 2003]. Data indicate that, changes of attentional function during hypnosis may induce dissociation of several psychological functions [Hilgard 1991, 1994] including dissociation of sensory and affective components of pain [Meier et al. 1993] as well as other pain related functions including generation of pain at the periphery, sensitization of secondary sensory neurons, modulation of endocrine-, immune- or autonomic responses and modulation of psychological functions [Miltner et al. 1992,

Larbig 2004, Carli 2009, Vanhaudenhuyse et al. 2009]. Besides pain perception, other physiological functions such as autonomic and hormonal responses, neuromuscular function, immune surveillance and immune/inflammatory reactions may also be advantageously influenced using hypnosis [Gruzelier et al. 2001, Gruzelier 2002, Mawdsley et al. 2008]. Hypnosis may be used for the treatment of bruxism and other neuromuscular symptoms, myofascial pain, atypical facial pain, oroposyrosis and other psychogenic pain [Staats and Krause 1995, Chaves 1997, Schmierer 1997,2004, Fábíán and Fábíán 1998] Hypnosis may be used highly efficiently also in the treatment of psychogenic denture intolerance [Ament and Ament 1970, Barsby 1997, Eli and Kleinhaus 1997, Fábíán and Fábíán 2000, Fábíán et al. 2007].

Self-Hypnosis

Self-hypnosis may be defined as an altered state of consciousness (ASC) achieved by self-suggestions. During practicing self-hypnosis, a deep hypnoid trance state (altered state of consciousness) develops similarly to those of hypnosis (see above). Autosuggestive techniques have been widely used from the earliest time of history [Heinze 1993, Hoppál 1993/a,b] but self-suggestions stood in the limelight of scientific interest in the last century only. There are several methods may be classified as self-hypnosis including (self)imagination techniques, active-graded-(self)hypnosis [Kretschmer 1946, Langen 1969], and biofeedback coupled techniques. Photo-acoustic stimulation (see below) may also be used to induce peculiar self-hypnotic states. The prototype of self-hypnotic methods is Autogenic Training [Schultz 1932], which is based on giving suggestions toward phenomena spontaneously occurring under relaxation to amplify and control them [Schultz 1932, Krause 1994]. Self-hypnosis practice can be resulted in prolonged increase of overall mental health, including decrease of depression scores, (trait)anxiety level, perceived stress scores, increase of self-observed level of mental energy and improved sleep quality under stress conditions. Sleeplessness (light sleep, disturbed sleep, insomnia) of patients can also be improved significantly using several self-hypnotic methods. Self-hypnosis may also be used efficiently for most forms of somatoform disorders; including somatoform chronic pain, pseudoneurological symptoms, autonomic dysfunction, pruritus (itching), several motor symptoms, fatigue, and allergic symptoms [Leuner and Schroeter 1997, Bongartz and Bongartz 2000, Langewitz et al. 2005].

Meditation

Although there are several forms of meditation, and no two meditation practices are alike in all features [Ospina et al. 2007, Halsband et al. 2009] meditation can be defined as volitional self-induced altered state of consciousness (ASC), established by mental faculties, without dominant contribution of other persons, highly intense body exercises or use of drugs [Fábíán in press]. Practices of meditation may be divided into two subgroups such as concentrative type and nonconcentrative type methods [Orne and Whitehouse 2000, Fábíán in press]. Although concentrative and nonconcentrative approaches differ from each other, the

subjective experience of deeper stages seems to be rather similar phenomenologically. Finally both group of meditation techniques, lead to a state of "Being", a state of "Knowing" or "Experiencing" without objectification and discursive thinking [Fábián in press]. In religious form of meditation, premised deep meditative states and experiences have certain religious overtones [Fábián in press]. There is an overall improvement of psychological abilities of meditators leading to more efficient social functioning and improved stress tolerance coupled with decreased level of neuroticism, depression, (trait)anxiety, trait negative affect, stress induced emotional irritability and stress induced anger. Long-run training of meditation also shortens habituation of stressful stimuli, and decreases the level of perceived stress [Fábián and Fejérdy 2010]. Meditative states also possess highly efficient anti-nociceptive effects. Sustained increase of pain tolerance [Kingston et al. 2007], reduction of pain induced distress [Mills and Farrow 1981] and reduced reactivity to pain in the brain [Kakigi et al. 2005, Orme-Johnson et al. 2006] were also reported following meditation training.

Photo-Acoustic Stimulation

Early studies demonstrated that turning the light or sound on and off induce alpha desynchronization on EEG [Berger 1930, Walter et al. 1946], leading to powerful stimulating effect on the central nervous system for a short time. In contrast, long lasting stimulation with flash light and tone signals (5-10 Hz frequency) leads to drowsiness and mixed alpha-theta activity [Williams and West 1975] coupled with body relaxation [Brauchli 1993] and appearance of significantly altered state of consciousness [Fábián et al. 2002]. These data indicate that, photo-acoustic stimulation whereas helps to keep the body relaxed, activates psychophysiological functions conducive to meditation (ASCs) coupled with somewhat increased arousal level [Fábián et al. 2005, 2009]. Photo-acoustic stimulation also exerts anti-depressive properties [Fábián et al. 2005, 2009].

Besides premised effects, flash light stimuli interact with the visual imagination leading to spontaneous appearance of various aspecific colored simple forms (i.e.: line, curve, web, lattice, spiral, cloud, tunnel etc.) [Fábián et al. 2005, 2009] similar to those induced by several hallucinogens in the phase of non-complex images [Siegel and Jarvik 1975, Siegel 1977]. (Using lower light intensities real visual imaginations may also appear [Fábián et al. 2005, 2009].) Since such visual imaginations appear spontaneously, a delightful experience of altered state of consciousness can be achieved even in case of patients with low motivation such as psychogenic denture intolerance (PDI) patients [Fábián et al. 2005, 2006, 2009].

Photo-acoustic stimulation improves oral immune surveillance. Both secretory immunoglobulin A [Brauchli 1993] and salivary chaperokine HSP70/HSPA [Fábián et al. 2004/b] (a salivary defense protein, molecular chaperone and cytokine) were increased significantly under photo-acoustic stimulation induced meditative state. Similarly, other oral defense proteins such as salivary amylase [Fábián et al. 2002, 2004/b] proposed to inhibit microbial growth [Shugars and Wahl 1998] and salivary lysozyme [authors' unpublished data] being bacteriolytic for gram-positive bacteria were also increased under premised meditative state. All premised changes induced by photo-acoustic stimulation advantageously improve defense of the oral cavity leading to clinical improvement of several oral mucosal symptoms especially of psychosomatic origin [Fábián and Fábián 2000, Fábián et al. 2005; 2007, 2009].

Somatoform orofacial chronic pain can also be treated efficiently using photo-acoustic stimulation [Fábián et al. 2002, 2005, 2006, 2009, Bálint et al. 2003]. Improvement of salivary gland function including significant increase of both flow rate and protein concentration following photo-acoustic stimulation treatment was also reported [Kaán et al. 2003, Fábián et al. 2009]. Premised improvement of salivary gland function is likely because of a prolonged decrease of sympathetic activity following treatment [Kaán et al. 2003]; notwithstanding that, there is a prompt, but short-term sympathetic activation under photo-acoustic stimulation. Tinnitus was also improved using photo-acoustic stimulation [Tönnies 2006]. Photo-acoustic stimulation was also used efficiently in the treatment of PDI cases with various somatic type manifestations [Fábián et al. 2004/a, 2006, 2009]. Photo-acoustic stimulation may be used highly efficiently also as a group-therapy [Fábián et al. 2006, 2009].

Biofeedback Methods

Biofeedback methods alter patient's physiological processes using devices that amplify signs of physiological processes that are ordinarily difficult to perceive without some type of amplification [Forgione and Mehler 2006]. Patients use the provided feedback signals as a guide. Measurable parameters of body functions coupled with relaxation such as muscle tension (EMG), skin conductance level (GSR or SCL), heart rate, heart rate variability (HRV) and skin temperature may be used efficiently for biofeedback treatment of orofacial psychosomatic patients [Forgione and Mehler 2006]. Feedback of breathing with light and sound signals called respiratory feedback [Leuner 1984, 1997, Barolin 2001] may also be used efficiently, because advantageous psychophysiological effects of breathing exercises and photo-acoustic stimuli also succeed in this case. Spontaneous appearance of altered state of consciousness was reported in significant proportion of subjects learning relaxation with the assistance of biofeedback equipment [McKee 1980, Leuner 2001] indicating that, therapeutic suggestions given under biofeedback may also be used highly efficiently [Leuner 2001]. Somatoform chronic pain may be treated efficiently using several biofeedback methods [Derra 2003, Kröner-Herwig 2004]. Other somatoform symptoms including pseudoneurological symptoms, autonomic dysfunction, pruritus (itching), several motor symptoms, fatigue (etc.) may also be treated efficiently with several biofeedback methods [Winkler and Krause 1989, Martin and Rief 2006]. Sleeplessness (light sleep, disturbed sleep, insomnia) of patients can also be improved with biofeedback [Krause 1983, Niepoth and Korn 2006].

Contraindication of Mind-Body Therapies

Contraindication of mind-body therapies includes several prepsychotic- and psychotic conditions, dementia and other deficiency of intelligence and also narcolepsy. Mind-body therapies may also be contraindicated under acute psychic trauma (crisis) because of increased liability toward regression of the patient. Treatment of borderline and narcissistic patients, strongly depressive or hysteric patients, introverted patients and also hypochondria patients may also be contraindicated [Fábián et al 2009, Fábián and Fejérdy 2010, Fábián in

press]. Great care should be taken with somatic diseases of unknown origin, because palliation of symptoms via mind-body techniques may impede definitive diagnosis of dangerous diseases (if any) [Leuner and Schroeter 1997]. A possible disadvantageous parasympathetic “rebound” effect (parasympathetic “overshoot”) induced by generalized relaxation of asthmatic patients was also expected in a study [Lehrer et al. 1997]. For the sake of completeness it can be noted that, increased epilepsy risk of meditation was also expected, but these expectations are likely exaggerated [Barnes 2005, Orme-Johnson 2005]. Photo-acoustic stimulation induced ASCs should be avoided in case of epileptic patients (danger of visually induced seizure), in several eye disorders (especially glaucoma) and also in case of blepharospasm [Fábián et al. 2005, 2009]. Photo-acoustic stimulation should also be used carefully in case of pregnancy, cardiac problems, pacemaker patients and other strongly compromised health conditions because of its strong influence on the central and autonomic nervous system [Fábián et al. 2005, 2009].

Complementary and Alternative Medicine

Prayer

Prayer is an active process of religious communicating with and/or appealing to God or other supernatural being belonging to God. The importance of religious methods is derived from that, they possess spiritual (sacral, transcendent) surplus, which improves efficiency of these kinds of methods [Fábián and Müller 2008, Fábián and Fejérdy 2010]. Evidently, religious techniques also improve religiosity, which is known to be associated with health promoting attitude and decreased frequency of medical symptoms and diseases. For psychosomatic therapy of religious patients at least prayer (the most frequently used religious CAM practice [Krause 2004, Tindle et al. 2005, Wahner-Roedler et al. 2005]) should be considered and utilized. Besides intercessory prayer (petitions on behalf of others) there are three major forms of prayer [Jantos and Kiat 2007] including conversational prayer (informal conversation with God), ritual prayer (reciting or reading well known prayers) and meditative prayer (i.e. repetition of short formulas, contemplation, imagery of religious themes etc.). A prolonged increase of overall mental health frequently occurs following regular practice of prayer [Meisenhelder and Chandler 2000] including positive mood [Wachholtz and Pargament 2005], decreased level of illness-related depression [Coleman et al. 2006], decreased anxiety [Harris et al. 2005, Wachholtz and Pargament 2005], higher level of perceived control over emotional reactions [Harris et al. 2005] and existential well being [Wachholtz and Pargament 2005]. Long-run practice of prayer also decreases psychophysiological stress reactions including salivary cortisol stress response [Tartaro et al. 2005]. It is also likely that, prayer improves immunological functions, and defense against infectious disorders [Fitzpatrick et al. 2007]. In general there is no contraindication of prayer, however practice of meditative forms may have contraindications similar to those of mind-body methods (see above).

Breathing Exercises

Breathing exercises can be classified primarily by frequency (low $\approx 6/\text{min}$; high $\approx 60\text{--}140/\text{min}$ and very low frequency $\leq 3\text{--}4/\text{min}$) [Fábián in press] but also by tidal volume (shallow or deep breathing), by location (abdominal, thoracic) and by specifically modified phases during breathing (shortening or lengthening inhalation, retention, exhalation and apnea) [Fábián in press]. Breathing exercises are primarily used to preserve healthiness, however they may also be used efficiently for therapeutic purposes. Breathing techniques can be utilized for mind-body therapies as a self-focus skill or anchor [Ospina et al. 2007, Fábián in press]. Similarly, combination of abdominal/diaphragmatic breathing with cognitive-behavioral therapy and progressive muscle relaxation was highly efficient to reduce pain-activity interference, pain intensity, depression, maladaptive pain beliefs, and catastrophizing as well as to improve masticatory jaw function in a randomized controlled trial [Turner et al. 2006]. Breathing exercises may be contraindicated in case of pregnancy, cardiac problems, pacemaker patients and other conditions with strongly compromised health. Breathing coupled meditation may have contraindications similar to those of mind-body methods (see above).

Tai-Chi and Qigong

Tai-Chi (Tai Chi Chuan) and Qigong (Chi Kung) are traditional Chinese practices of meditation designed to control expected vital energy (called "Chi" or "Qi") of the body to promote health and spiritual development. Thai-Chi (but not Qigong) may also be considered as a Chinese martial art based on suppleness and evasion [Cheng 2007] coupled with control of premised expected vital energy ("Chi"/"Qi"). Besides some static body postures and voice training, there are three major components of Tai-Chi and Qigong training including slow moving exercises, breathing exercises and meditation [Lewis 2000; Jones 2001; McCaffre and Flower 2003].

Movement in Qigong usually involve meditative visualizing of internal consequences of flow of the expected vital energy, although in some cases vital energy is expected to be transferred into other persons (patients) on purpose to cure. Movement in Tai-Chi (although usually practiced slowly) may be sped up and might involve meditative visualizing the external consequences of a motion as well (i.e. to provide a self-defense) which is not the case in Qigong. Tai-Chi and Qigong practice can be advantageously utilized for the treatment of psychosomatic patients especially when practiced repeatedly for longer-run. Prolonged increase of overall mental health can be achieved with long-run practicing regularly, including decrease of perceived stress and (trait)anxiety; as well as improve of mood, vitality, sleep quality and general social functioning [Fábián and Fejérdy 2010]. Tai-Chi and Qigong practices may be contraindicated in case of pregnancy, cardiac problems, pacemaker patients and other conditions with strongly compromised health [Vigh 1972, Fábián in press]. Meditative forms may have contraindications similar to those of mind-body methods (see above).

Yoga

Yoga is a collective noun of traditional practices developed in India to improve health, and to prepare body and mind for meditation and spiritual development. Classical yoga incorporates moral and ethical observances (yama, niyama), body postures and exercises (i.e. hatha yoga), breathing techniques (pranayama) and meditative techniques (i.e. dhyana, samadhi) [Joshi 1967; Ospina et al. 2007, Fábíán in press]. There are several major subdivisions of yogic tradition such as hatha yoga, pranajama, kundalini yoga, and raja yoga [Fábíán in press]. There are also some modern schools of yoga such as Iyengar yoga and Vivekananda yoga [Fábíán in press]. Hatha yoga includes some dynamic body exercises, but most typically several breathing exercises (pranayama, see below) and body postures like sittings, standing poses, inverted poses, muscle and spine stretching poses, trunk rotating poses, symbolic hand/body gestures, and relaxing poses. Pranayama is a collective noun of all breathing exercises of yoga tradition including breathing exercises of hatha yoga and kundalini yoga. Kundalini yoga is a yogic practice aimed at control of expected vital-energy of the body (called "prana") via meditation coupled with certain body postures and breathing exercises. Raja yoga is a pure meditative form of yogic exercises aimed at self-absorption experience (called "samadhi") [Fábíán in press]. Iyengar yoga is based on hatha yoga traditions (including pranayama and meditation) but employs props that allow to practice body postures despite limited experience and flexibility. Vivekananda yoga is also based on hatha yoga traditions (including pranayama and meditation) but aimed at therapeutic utilization of yoga tradition. Long-run practice of yoga induce prolonged increase of overall mental health including increased level of optimism and decreased level of perceived stress, neuroticism, depression, (trait)anxiety and anger under chronic illness-stress [Shapiro et al. 2007, Smith et al. 2007, Kjellgren et al. 2007]. Yoga was found highly efficient also in the treatment of unexplained chronic fatigue as well [Bentler et al. 2005]. Yoga may be contraindicated in case of pregnancy, cardiac problems, pacemaker patients and other conditions with strongly compromised health [Vigh 1972, Fábíán in press]. Meditative forms may have contraindications similar to those of mind-body methods (see above).

Other Therapies

Some other forms of CAM therapies like sleep deprivation, keeping fast and light therapy may also be used. Sleep deprivation (vigil) positively influences mood having prompt and significant antidepressive effect [Benedetti et al. 2003/a,b, Wirz-Justice et al. 2005]. Vigil decreases anterior cingulate cortex metabolism likely due to an increase in the activity of brain serotonergic [Benedetti et al. 2003/a] and dopaminergic [Benedetti et al. 2003/b] pathways as well. Sleep deprivation coupled increase of the serotonergic neural activity in dorsal raphe nucleus was also demonstrated in an animal study [Gardner et al. 1997]. Keeping fast is frequently used as a practice to preserve healthiness. Fasting therapy may also improve chronic pain [Michalsen et al. 2002] as well as syndromes of stress and exhaustion [Michalsen et al. 2002]. Fasting activates hypothalamic-pituitary-adrenocortical axis (HPA axis) leading to increased serum cortisol level and sympathetic activity [Fichter and Pirke 1986]. Altered function of hypothalamic and hippocampal septal systems was also assumed

[Winkelman 1986; Heinze 1993]. Fasting also activates the immune system (NK cell activity) and alleviates fatigue [Masuda et al. 2001], however a complete food restriction for two-three days (or more) may lead to immune suppression [Chiapelli and Hodgson 2000]. Light therapy (taking delight in the sunrise, sunset, candlelight, glittering surfaces of water and snow as well as sun-bathing) is frequently practiced by many people. Besides the rewarding and symbolic meaning of such practices, also neurophysiologic effects of the light should be counted in these cases. Light stimuli influences the function of pineal gland, epithalamus and supraoptic region of hypothalamic medial zone [Clark et al 2006]. Further, significant effect of light exposure on mood (i.e. antidepressive effect [Benedetti et al 2003/a, Wirz-Justice et al. 2005]) likely based on brain serotonergic pathways and phase changes of biological rhythms [Benedetti et al 2003/a] was also reported.

Conclusion

Even though prosthesis is fabricated conscientiously and properly, there is no assurance that the patient will be comfortable while wearing it or satisfied with the therapy [Mazurat and Mazurat 2003]. A normative evaluation by a dentist and a subjective evaluation by the patient related to the denture or to the dental treatment may be rather different [Lechner and Roessler 2001]. There are various psychogenic symptoms, which may occur in relation with making denture [Fábián and Fábián 2000, Fábián et al. 2006, 2007]. Although these symptoms are primarily of psychogenic origin, their pathomechanisms may include both somatic and psychogenic mechanisms. Therefore, both somatic and psychogenic pathways of pathomechanism should be considered carefully for a proper choice of methods used for palliative dental care, mind-body therapy as well as prevention of relapse using certain CAM therapies. Since lacking a religious sensitivity, medical science will always be woefully incomplete, no matter how great its discoveries [Pollack 1999] the importance of religious methods of complementary and alternative medicine should also not be underestimated.

References

- Adachi, N; Tomonaga, S; Tachibana, T; Denbow, DM; Furuse, M. (-)-Epigallocatechin gallate attenuates acute stress response through GABAergic system in the brain. *Eur. J. Pharmacol.*, 2006 531, 1717-175.
- Ament, P. and Ament, A. Body image in dentistry. *J. Prosthet Dent*, 1970 24, 362-366.
- Amorim, JC; de Sousa, GR; de Barros Silveira, L; Prates, RA, Pinotti, M; Ribeiro, MS. Clinical study of the gingiva healing after gingivectomy and low-level laser therapy. *Photomed Laser Surg*, 2006 24, 588-594.
- Baad-Hansen, L. Atypical odontalgia - pathophysiology and clinical management. *J. Oral Rehabil.*, 2008 35, 1-11.
- Baldry, PE. *Acupuncture, trigger points and musculoskeletal pain*. Third edition. Edinburgh-London-New York-Philadelphia-San Francisco-Toronto: Elsevier Churchill Livingstone; 2005. 3-13; 29-36.

- Bálint, M; Krause, M; Krause W-R; Kaán, B; Fejérdy, L; Gáspár, J; Fábíán, TK. Modification of the photo-acoustic stimulation in the psychotherapy of oral psychosomatic patients. Preliminary experiences. *Fogorv Szle*, 2003 96, 171-174.
- Bányai, ÉI. and Hilgard ER. A comparison of active-alert hypnotic induction with traditional relaxation induction. *J. Abnorm. Psychol.*, 1976 85, 218-224.
- Bányai, ÉI. Toward a social-psychobiological model of hypnosis. In: Lynn SJ, Rhue JW editors. *Theories of hypnosis: Current models and perspectives*. New York - London: Quilford Press; 1991. 564-598
- Barnes, VA. EEG, hypometabolism, and ketosis during transcendental meditation indicate it does not increase epilepsy risk. *Med Hypothesis*, 2005 65, 202-203.
- Barolin, GS. Das Respiratorische Feedback (RBF) - Basis und Praxis. In: Barolin GS, editor. *Das Respiratorische Feedback nach Leuner*. Berlin: VWB Verlag für Wissenschaft und Bildung; 2001. 9-32.
- Barsby, M. Hypnosis in the management of denture intolerance. *Hypn Int. Monographs*, 1997 3, 71-78.
- Beck, AT. *Cognitive theory and emotional disorders*. New York: International University Press; 1976.
- Bell, WE. Clinical diagnosis of the pain-dysfunction syndrome. *JADA*, 1969/a 79, 154-160.
- Bell, WH. Nonsurgical management of the pain-dysfunction syndrome. *JADA*, 1969/b 79, 161-170.
- Benedetti, F; Colombo, C; Serretti, A; Lorenzi, C; Pontiggia, A; Barbini, B; Smeraldi, E. Antidepressant effects of light therapy combined with sleep deprivation are influenced by a functional polymorphism within the promoter of the serotonin transporter gene. *Biol. Psychiatry*, 2003/a 54, 687-692.
- Benedetti, F; Serretti, A; Colombo, C; Lilli, R; Lorenzi, C; Smeraldi, E. Dopamine receptor D2 and D3 gene variants are not associated with the antidepressant effects of total sleep deprivation in bipolar depression. *Psychiatry Res.*, 2003/b 118, 241-247.
- Bentler, SE; Hartz, AJ; Kuhn, EM. Prospective observational study of treatments for unexplained chronic fatigue. *J. Clin. Psychiatry*, 2005 66, 625-631.
- Benton, D. and Donohoe, RT. The effects of nutrients on mood. *Public Health Nutr.*, 1999 2, 403-409.
- Berger, H. Über das Elektro-Enkephalogram des Menschen. II. *J. Psychol. Neurol. (Lpz)*, 1930 40; 160-169.
- Binder, A; Parr, G; Hazleman, B; Fitton-Jackson, S. Pulsed electromagnetic field therapy of persistent rotator cuff tendinitis. A double-blind controlled assessment. *Lancet*, 1984 1(8379), 695-698.
- Binder, H. and Binder, K. *Autogenes Training, Basispsychotherapeutikum*. Köln: Deutscher Ärzteverlag; 1989.
- Biondi, M. and Picardi, A. Temporomandibular joint pain-dysfunction syndrome and bruxism: etiopathogenesis and treatment from a psychosomatic integrative viewpoint. *Psychother Psychosom*, 1993 59, 84-98.
- Bishop, TS. High frequency neural modulation in dentistry. *JADA*, 1986 112, 176-177.
- Black, RR. Use of transcutaneous electrical nerve stimulation in dentistry. *JADA*, 1986 113, 649-652.
- Bongartz, W. and Bongartz, B. *Hypnosetherapie* 2-nd edition. Göttingen - Bern -Toronto - Seattle: Hogrefe; 2000; 252-254, 312.

- Boos, RH. Preparation and conditioning of patients for prosthetic treatment. *J. Prosthet Dent*, 1959 9, 4-10.
- Brauchli, P. Comparative study of the psychophysiologic relaxation effects of an optic-acoustic mind machine with relaxation music. *Z. Exp. Angew Psychol.*, 1993 40, 179-193.
- Brodine, AL. and Hartshorn, MA. Recognition and management of somatoform disorders. *J. Prosthet Dent*, 2004 91, 268-273.
- Brown, D. The use of Vitex agnus castus for hyperprolactinemia. *Quarterly Rev Nat Med*, 1997 (SPRING), 19-21.
- Bruinsma, K. and Taren, DL. Chocolate: Food or drug? *J. Am. Diet. Assoc.*, 1999 99, 1249-1256.
- Burdette, JE; Liu, J; Chen, SN; Fabricants, DS; Piersen, CE; Barker, EL; Pezzuto, JM; Mesecar, A; Van Breemen, RB; Farnsworth, NR; Bolton, JL. Black cohosh acts as a mixed competitive ligand and partial agonist of the serotonin receptor. *J. Agric. Food Chem.*, 2003 51, 5661-5670.
- Carli, G. An update on pain physiology: the relevance of Craig's and Jänning's hypotheses for hypnotic analgesia. *Contemp Hypnosis*, 2009 26, 4-14.
- Carlson, S; Peng, N; Prasain, JK; Wyss, JM; Effects of botanical dietary supplements on cardiovascular, cognitive, and metabolic function in males and females. *Gender Med*, 2008 5 (Suppl.A), S76-S90.
- Chaco, J. Electromyography of the masseter muscles in Costen's syndrome. *J. Oral Med.*, 1973 28, 45-46.
- Chaves, JF. Hypnosis in dentistry: Historical overview and current appraisal. *Hypn Int. Monographs*, 1997 3, 5-24
- Cheng, TO. Thai Chi: The Chinese ancient wisdom of an ideal exercise for cardiac patients. *Int. J. Cardiology*, 2007 117, 293-295.
- Chiapelli, F. and Hodgson, D. Immune suppression. In: Fink G, editor in chief. *Encyclopedia of stress*. (Vol. 2.). San Diego: Academic Press; 2000; 531-536.
- Clark, DL; Boutros, NN; Mendes, MF. *The brain and behavior. An introduction to behavioral neuroanatomy*. Cambridge: Cambridge University Press; 2006. 131, 133, 135.
- Coleman, CL; Holzemer, WL; Eller, LS; Corless, I; Reynolds, N; Nokes, KM; Kemppainen, JK; Dole, P; Kirksey, K; Sefcik, L; Nicholas, P; Hamilton, MJ. Gender differences in use of prayer as a self-care strategy for managing symptoms in African Americans living with HIV/AIDS. *J. Assoc. Nurses Aids Care*, 2006 17, 16-23.
- Crawford, HE. and Gruzelier, JH. A midstream view of the neuropsychology of hypnosis: Recent research and future directions. In: Fromm E, Nash MR editors. *Contemporary hypnosis research*. New York - London: Guilford Press, 1992, 227-266.
- Curcio, FB; Tackney, VM; Berweger, R. Transcutaneous electrical nerve stimulation in dentistry: A report of a double-blind study. *J. Prosthet Dent*, 1987 58, 379-383.
- Dahlström, L; Carlsson, GE; Carlsson, SG. Comparison of effects of electromyographic biofeedback and occlusal splint therapy on mandibular dysfunction. *Scand J. Dent Res.*, 1982 90, 151-156.
- Dahlström, L; Haraldson, T; Janson, ST. Comparative electromyographic study of bite plates and stabilization splints. *Scand J. Dent Res.*, 1985 93, 262-268.
- Dahlström, L. Electromyographic studies of craniomandibular disorders: a review of the literature. *J. Oral. Rehabil.*, 1989 16, 1-20.

- Dahlström, L. Conservative treatment methods in craniomandibular disorder. *Swed. Dent J.*, 1992 16, 217-230.
- Daniele, C; Coon, JT; Pittler, MH; Ernst, E. Vitex agnus castus: A systematic review of adverse events. *Drug Safety*, 2005 28, 319-332.
- Darendeliler, MA; Sinclair, PM; Kusy, RP. The effects of samarium-cobalt magnets and pulsed electromagnetic fields on tooth movement. *Am. J. Orthod. Dentofacial Orthop.*, 1995 107, 578-588.
- de Medeiros, JS; Vieira, GF; Nishimura, PY. Laser application effects on the bite strength of the masseter muscle, as an orofacial pain treatment. *Photomed Laser Surg*, 2005 23, 373-376.
- de Silva, P. Behavior therapy. In: Fink G, editor in chief. *Encyclopedia of stress*. (Vol. 1.). San Diego: Academic Press; 2000. 300-304.
- Derra, C. Entspannungsverfahren und Hypnose. In: Egle UT, Hoffmann SO, Lehmann KA, Nix WA editors. *Handbuch chronischer Schmerz. Grundlagen, Pathogenese, Klinik und Therapie aus bio-psycho-socialer Sicht*. Stuttgart-New York: Schattauer GmbH; 2003. 392-403.
- Derra C. and Egle, UT. Psychopharmaka in der Schmerztherapie. In: Egle UT, Hoffmann SO, Lehmann KA, Nix WA editors. *Handbuch Chronischer Schmerz. Grundlagen, Pathogenese, Klinik und Therapie aus bio-psycho-sozialer Sicht*. Stuttgart - New York: Schattauer GmbH; 2003; 352-359.
- DiTomaso, E; Beltramo, M; Piomelli, D. Brain cannabinoids in chocolate. *Nature*, 1996 382, 677-678.
- Eli, I. and Kleinhauz, M. Hypnosis and dentistry. *Hypn Int. Monographs*, 1997 3, 59-70.
- Ellis, AT. *Reason and emotion in psychotherapy*. New York: Lyle Stuart; 1962.
- Fábián, TK. *Mind-body connections. Pathways of psychosomatic coupling under meditation and other altered states of consciousness*. New York: Nova Science Publishers; in press.
- Fábián, TK. and Fábián, G. Stress of life, stress of death: Anxiety in dentistry from the viewpoint of hypnotherapy. *Ann. NY Acad. Sci.*, 1998 851, 495-500.
- Fábián, TK. and Fábián, G. Dental stress. In: Fink G, editor in chief. *Encyclopedia of stress*. (Vol. 1.). San Diego: Academic Press; 2000; 567-659.
- Fábián, TK and Fejérdy P. *Psychogenic denture intolerance. Theoretical background, prevention and treatment possibilities*. New York: Nova Science Publishers; 2010.
- Fábián, TK. and Müller, O. Prayer, meditation and healing [Ima és meditáció a gyógyításban]. In: Vértés G, Fábián TK editors. *Religion, faith and healing [Vallás és hit a gyógyításban]*. Budapest: Medicina; 2008. 118-125.
- Fábián, TK; Vértés, G; Szabó, A; Varga, K. Photo-acoustic stimulation and hypnotherapy. An effective combination for treatment of oral psychosomatic disorders. *Hypn Int. Monographs*, 2002 6, 199-207.
- Fábián, TK; Kaán, B; Fejérdy, L; Tóth, Zs; Fejérdy, P. Effectiveness of psychotherapy in the treatment of denture intolerance. Evaluation of 25 cases. *Fogorv Szle*, 2004/a 97, 163-168.
- Fábián, TK; Tóth, Zs; Fejérdy, L; Kaán, B; Csermely, P; Fejérdy, P. Photo-acoustic stimulation increases the amount of 70 kDa heat shock protein (Hsp70) in human whole saliva. A pilot study. *Int. J. Psychophysiol.*, 2004/b 52, 211-216.

- Fábián, TK; Krause, WR; Krause, M; Fejérdy, P. Photo-acoustic stimulation and hypnotherapy in the treatment of oral psychosomatic disorders. *Hypnos*, 2005 32, 198-202.
- Fábián, TK; Mierzwińska-Nastalska, E; Fejérdy P. Photo-acoustic stimulation. A suitable method in the treatment of psychogenic denture intolerance. *Protet Stomatol*, 2006 56, 335-340.
- Fábián, TK; Fábián, G; Fejérdy, P. Dental Stress. In: Fink G, editor in chef. *Encyclopedia of Stress*. 2-nd enlarged edition, (Vol. 1.) Oxford: Academic Press; 2007; 733-736.
- Fábián, TK; Kovács, KJ; Gótai, L; Beck, A; Krause, WR; Fejérdy, P. Photo-acoustic stimulation: Theoretical background and ten years of clinical experience. *Contemp Hypn*, 2009 26, 225-233.
- Farthing, GW. *The psychology of consciousness*. Prentice Hall New Jersey: Englewood Cliffs; 1992; 202-219, 478-497.
- Fichter, MM. and Pirke, KM. Effect of experimental and pathological weight loss upon the hypothalamo-pituitary-adrenal axis. *Psychoneuroendocrinology*, 1986 11, 295-305.
- Fitzpatrick, AL; Standish, LJ; Berger, J; Kim, JG; Calabrese, C; Polissar, N. Survival in HIV-1-positive adults practicing psychological or spiritual activities for one year. *Altern Ther. Health Med.*, 2007 13, 18-24.
- Forgione, AG. and Mehler, BL. Biofeedback in the treatment of myofascial pain disorder and temporomandibular joint pain. In: Mostofsky DI, Forgione AG, Giddon DB editors. *Behavioral dentistry*. Ames (Iowa): Blackwell Munksgaard; 2006; 51-64.
- Freesmeyer, WB. Okklusionsschienen. In: Koeck B, editor. *Funktionsstörungen des Kauorgans*. München - Wien - Baltimore: Urban and Schwarzenberg; 1995; 215-241.
- Freud, S. Die Traumdeutung (1900) In: Strachey J, editor. *The standard edition of the complete psychological works of Sigmund Freud*. (Vol. IV-V.) London: Imago; 1953-1966.
- Friedlander, AH; Marder, SR; Sung, EC. Panic disorder. Psychopathology, medical management and dental implications. *JADA*, 2004 135, 771-778.
- Fujita, S; Yamaguchi, M; Utsunomiya, T; Yamamoto, H; Kasai, K. Low-energy laser stimulates tooth movement velocity via expression of RANK and RANKL. *Orthod Craniofac Res.*, 2008 11, 143-155.
- Gardner, JP; Fornal, CA; Jacobs, BL. Effect of sleep deprivation on serotonergic neuronal activity in the dorsal nucleus of the freely moving cat. *Neuropsychopharmacology*, 1997 17, 72-81.
- Gáspár, J; Linninger, M; Kaán, M; Bálint, M; Fejérdy, L; Fábián, TK. The effectivity of standardized direct suggestions under dental hypnotic conditions. *Fogorv Szle*, 2003 96, 205-210.
- Geller, SE. and Studee, L. Botanical and dietary supplements for menopausal symptoms: What works, what doesn't. *J. Womens Health (Larchmt)*, 2005 14, 634-649.
- Gellrich, NC; Schramm, A; Böckmann, R; Kugler, J. Follow-up in patients with oral cancer. *J. Oral Maxillofac Surg*, 2002 60, 380-386.
- Ginsburg, I; Vennos, C; Koren, E. Inflammaging - Altern als Konsequenz chronischer Entzündungen: Das Beispiel Padma 28. *Schweiz Zschr GanzheitsMedizin*, 2008 20, 412-417.
- Gleditsch, JM. Punktversuche und Ermittlung von Reaktionsebenen mit Hilfe der Very-Point-Technik. *Akupunktur Theor Prax*, 1980 8, 58-61.

- Gold, N; Greene, CS; Laskin, DM. TENS therapy for treatment of MPD (abstract). *IADR Progr and Abstracts*. 1983, 244.
- Gossling, HR; Bernstein, RA; Abbott, J. Treatment of ununited tibial fractures: a comparison of surgery and pulsed electromagnetic fields (PEMF). *Orthopedics*, 1992 15, 711-719.
- Green, CS. and Laskin, DM. Splint therapy for the myofascial pain-dysfunction (MPD) syndrome: a comparative study. *JADA*, 1972/a 84, 624-628.
- Green, CS. and Laskin, DM. Therapeutic effects of diazepam (valium) and sodium salicylate in myofascial pain-dysfunction (MPD) patients (abstract). *IADR Progr and Abstr*, 1972/b, 96.
- Green, CS. and Laskin, DM. Temporomandibular disorders: Moving from a dentally based to a medically based model. *J. Dent Res.*, 2000 79, 1736-1739.
- Griffin, CJ: The treatment of the temporomandibular joint syndrome. In: Griffin CJ, Harris R editors. *The temporomandibular joint syndrome. The masticatory apparatus of man in normal and abnormal function*. (Monographs in oral science, Vol. 4.) Basel - München - Paris - London - New York - Sydney: Karger; 1975; 170-187.
- Grube, B; Walper, A; Wheatley, D. St John's wort extract: efficacy for menopausal symptoms of psychological origin. *Adv. Ther.*, 1999 16; 177-186.
- Gruzelier, JH. A review of the impact of hypnosis, relaxation, guided imagery and individual differences on aspects of immunity and health. *Stress*, 2002 5, 147-163.
- Gruzelier, J; Smith, F; Nagy, A; Henderson, D. Cellular and humoral immunity, mood and exam stress: the influences of self-hypnosis and personality predictors. *Int. J. Psychophysiol*, 2001 42, 55-71.
- Halsband, U; Mueller, S; Hinterberger, T; Strickner, S. Plasticity changes in the brain in hypnosis and meditation. *Contemp Hypn*, 2009 26, 194-215.
- Hammarstrand, G; Berggren, U; Hakeberg, M. Psychophysiological therapy vs. hypnotherapy in the treatment of patients with dental phobia. *Eur. J. Oral Sci*, 1995 103, 399-404.
- Harris, JI; Schoneman, SW; Carrera, SR. Preferred prayer styles and anxiety control. *J. Religion Health*, 2005 44, 403-412.
- Heinze, R-I. Shamanistic states of consciousness: Access to different realities. In: Hoppál M, Howard KD editors. *Shamans and cultures* (ISTOR books 5). Budapest - Los Angeles: Akadémiai Kiadó and International Society of Trans-Oceanic Research; 1993; 169-178.
- Higdon, JV. and Frei, B. Coffee and health: A review of recent human research. *Crit. Rev. Food Sci Res*, 2006 46, 101-123.
- Hilgard, ER. A neodissociation interpretation of hypnosis. In: Lynn SJ, Rhue JW editors. *Theories of hypnosis. Current models and perspectives*. New York - London: Guilford Press; 1991. 83-104.
- Hilgard, ER. Neodissociation theory. In: Lynn SJ, Rhue JW editors. *Dissociation: Clinical and theoretical perspectives*. New York: Guilford Press; 1994. 32-51.
- Hochman, R. Neurotransmitter modulator (TENS) for control of dental operative pain. *JADA*, 1988 116, 208-212.
- Hoppál, M. Shamanism: Universal structures and regional symbols. In: Hoppál M, Howard KD editors. *Shamans and cultures*. (ISTOR books 5). Budapest - Los Angeles: Akadémiai Kiadó and International Society of Trans-Oceanic Research; 1993/a. 181-192.
- Hoppál, M. Studies on Eurasian shamanism. In: Hoppál M, Howard KD editors. *Shamans and cultures*. (ISTOR books 5). Budapest - Los Angeles: Akadémiai Kiadó and International Society of Trans-Oceanic Research; 1993/b. 258-288.

- Inoue, N; Ohnishi, I; Chen, D; Deitz, LW; Schwaradt, JD; Chao, EY. Effect of pulsed electromagnetic fields (PEMF) on late-phase osteotomy gap healing in a canine tibial model. *J. Orthop. Res.*, 2002 20, 1106-1114.
- Israel, HA. and Scrivani, SJ. The interdisciplinary approach to oral, facial and head pain. *JADA*, 2000 131, 919-926.
- Iversen, G. Geleitwort. In: Binder, H; Binder, K. *Autogenes Training, Basispsychotherapeutikum*. Köln: Deutscher Ärzteverlag; 1989.
- Jantos, M. and Kiat, H. Prayer as medicine: how much have we learned? *Med. J. Australia*, 2007 186, S51-S53.
- Jones, BM. Change in cytokine production in healthy subjects practicing Guolin Qigong: a pilot study. *BMC Compl Altern Med*, 2001 1, 8.
- Joshi, KS. *Yoga and personality*. Allahabad: Udayana Publications, 1967. 16, 98, 127.
- Kaán, B; Krause, W-R; Krause, M; Fejérdy, L; Gáspár, J; Bálint, M; Fábíán, TK. Effects of photo-acoustic stimulation combined with hypnotherapy on saliva secretion. A pilot study. *Fogorv Szle*, 2003 96, 217-221.
- Kakigi, R; Nakata, H; Inui, K; Hiroe, N; Nagata, O; Honda, M; Tanaka, S; Sadato, N; Kawakami, M. Intracerebral pain processing in a yoga master who claims not to feel pain during meditation. *Eur. J. Pain*, 2005 9, 581-589.
- Kanje, M; Rusovan, A; Siskin, B; Lundborg, G. Pretreatment of rats with pulsed electromagnetic fields enhances regeneration of the sciatic nerve. *Bioelectromagnetics*, 1993 14, 353-359.
- Karppinen, K; Eklund, S; Suoninen, E; Eskelin, M; Kirveskari, P. Adjustment of dental occlusion in treatment of chronic cervicobrachial pain and headache. *J. Oral. Rehabil.*, 1999 26, 715-721.
- Kawasaki, K. and Shimizu, N. Effects of low-energy laser irradiation on bone remodeling during experimental tooth movement in rats. *Lasers Surg. Med.*, 2000 26, 282-291.
- Khadra, M; Kasem, N; Lyngstadaas, SP; Haanæs HR; Mustafa, K. Laser therapy accelerates initial attachment and subsequent behavior of human fibroblasts cultured on titanium implant material. *Clin. Oral Impl. Res.*, 2005/a 16, 168-175.
- Khadra, M; Lyngstadaas, SP; Hans, R; Haanæs, Mustafa, K. Effect of laser therapy on attachment, proliferation and differentiation of human osteoblast-like cells cultures on titanium implant material. *Biomaterials*, 2005/b 26, 3503-3509.
- Khullar, SM; Brodin, P; Messelt, EB; Haanæs, HR. The effects of low level laser treatment on recovery of nerve conduction and motor function after compression injury in the rat sciatic nerve. *Eur. J. Oral. Sci.*, 1995 103, 299-305.
- Kihlstrom, JF. Conscious, subconscious, unconscious: A cognitive perspective. In: Bowers KS, Meichenbaum D editors. *The unconscious reconsidered*. New York: John Wiley and Sons; 1984; 149-211.
- Kimura, Y; Wilder-Smith, P; Yonaga, K; Matsumoto, K. Treatment of dentine hypersensitivity by lasers: a review. *J. Clin. Periodontol.*, 2000 27, 715-721.
- King, RB. Concerning the management of pain associated with herpes zoster and of postherpetic neuralgia. *Pain*, 1988 33, 73-78.
- Kjellgren, A; Bood, SÅ; Axelsson, K; Norlander, T; Saatcioglu, F. Wellness through a comprehensive yogic breathing program - A controlled pilot trial. *BMC Compl Altern Med*, 2007 7, 43.
- Kleinen, J. and Knipschild, P. Ginkgo biloba. *Lancet*, 1992 340, 1136-1139.

- Köhler, T. Fortbildung: Psychopharmakotherapie (2. Teil). *Verhaltenstherapie and Verhaltensmedizin*, 2005 26, 141-168.
- Krause, WR. Grundlagen der Biofeedback-Therapie und ihre therapeutischen Anwendungsmöglichkeiten. *Z ärztl Fortbild*, 1983 77, 493-494.
- Krause, WR. Hypnose und Autogenes Training (Selbsthypnose) in der Rehabilitation. Ergänzung in: Schultz, JH. *Hypnose-Technik. Praktische Anleitung zum Hypnotisieren für Ärzte*. 9. Auflage - bearbeitet und ergänzt von G. Iversen und W.-R. Krause. Stuttgart - Jena - New York: Gustav Fischer Verlag; 1994. 71-79.
- Krause, WR. Hypnose und Autogenes Training (Selbsthypnose) in der Rehabilitation. Ergänzung in: Schultz, JH. *Hypnose-Technik. Praktische Anleitung zum Hypnotisieren für Ärzte*. 9. Auflage - bearbeitet und ergänzt von G. Iversen und W.-R. Krause. Stuttgart - Jena - New York: Gustav Fischer Verlag; 1994. 71-79.
- Krause, N. Religion, aging and health: Exploring new frontiers in medical care. *South Med. J.*, 2004 97, 1215-1222.
- Kretschmer, E. Über gestufte aktive Hypnoseübungen und den Umbau der Hypnosetechnik. *Dtsch Med. Wschr*, 1946 71, 281-283.
- Kröner-Herwig, B. Biofeedback. In: Basler H-D, Franz K, Kröner-Herwig B, Rehfisch H-P editors. *Psychologische Schmerztherapie*. Berlin-Heidelberg-New York: Springer; 2004; 551-565.
- Kröner-Herwig, B; Mohn, U; Pothmann, R. Comparison of biofeedback and relaxation in the treatment of pediatric headache and the influence of parent involvement on outcome. *Appl Psychophysiol Biofeedback*, 1998 23, 143-157.
- Laney, WR. Medical preparations of the mouth for prostheses. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment of prosthodontics*. Philadelphia: Lea and Febiger; 1983; 182-193.
- Langen, D. *Die gestufte Aktivhypnose*, 3. Aufl. Stuttgart: Huber; 1969.
- Langewitz, W; Izakovic, J; Wyler, J; Schindler, C; Kiss, A; Bircher, AJ. Effect of self-hypnosis on hay fever symptoms - a randomized controlled intervention study. *Psychoter Psychosom*, 2005 74, 165-172.
- Larbig W. Hirnphysiologische Korrelate der Hypnoanalgesie. *HyKog*, 2004 21, 39-59.
- Laskin, DM. and Block, S. Diagnosis and treatment of myofacial pain-dysfunction (MPD) syndrome. *J. Prosthet Dent*, 1986 56, 75-84.
- Lechner, SK. and Roessler, D. Strategies for complete denture success: beyond technical excellence. *Compend Contin Educ. Dent.*, 2001 22, 553-559.
- Lee, EW; Maffulli, N; Li, CK; Chan, KM. Pulsed magnetic and electromagnetic fields in experimental achilles tendonitis in the rat: a prospective randomized study. *Arch. Phys. Med. Rehabil*, 1997 78, 399-404.
- Lehrer, PM; Hochron, SM; Mayne, TM; Isenberg, S; Lasoski, AM; Carlson, V; Gilchrist, J; Porges, S. Relationship between changes in EMG and respiratory sinus arrhythmia in a study of relaxation therapy for asthma. *Appl. Psychophysiol Biofeedback*, 1997 22, 183-191.
- Leuner, H. Zur Indikation und wissenschaftlichen Fundierung des Respiratorisches Feedbacks. *Allgemeinarzt*, 1984 6, 334-354.
- Leuner, H. Respiratorische Feedback (RFB) als atembezogene Selbsthypnose. (Anhang). In: Leuner H, Schroeter E editors. *Indikationen und spezifische Anwendungen der*

- Hypnosebehandlung. Ein Überblick.* (2. erweiterte Auflage). Bern-Göttingen-Toronto-Seattle: Hans Huber Verlag; 1997; 204-209.
- Leuner, H. Manuskriptfragmente. In: Barolin GS, editor. *Das Respiratorische Feedback nach Leuner*. Berlin: VWB Verlag für Wissenschaft und Bildung; 2001; 39-105.
- Leuner, H. and Schroeter, E. *Indikationen und spezifische Anwendungen der Hypnosebehandlung*. Bern-Göttingen-Toronto-Seattle: Hans Huber Verlag; 1997; 24, 52, 66, 152-155, 158-160, 173-175.
- Lewis, DE. T'ai chi ch'uan. *Comp. Ther. Nurs. Midwifery*, 2000 6, 204-206.
- Linde, K; Ramirez, G; Mulrow, CD; Pauls, A; Weidenhammer, W; Melchart, D. St John's wort for depression -- an overview and meta-analysis of randomized clinical trials. *Brit. Med. J.*, 1996 313, 253-258.
- Linde, K; Ramirez, G; Mulrow, CD; Pauls, A; Weidenhammer, W; Melchart, D. St John's wort for depression: meta-analysis of randomized controlled trials. *Brit. J. Psychiatry*, 2005 186, 99-107.
- Liske, E. Therapeutic efficacy and safety of Cimifuga racemosa for gynecologic disorders. *Adv Ther*, 1998 15, 45-53.
- Lucks, BC. Vitex agnus castus essential oil and menopausal balance: a research update. *Compl Ther. Nurs. Midwifery*, 2003 9, 157-160.
- Lund, I; Lundeberg, T; Carleson, J; Sönnerrfors, H; Uhrlin, B; Svensson, E. Corticotropin releasing factor in urine - A possible biochemical marker of fibromyalgia responses to massage and guided relaxation. *Neurosci. Letters*, 2006 403, 166-171.
- Maeda-Yamamoto, M; Ema, K; Shibuchi, I. In vitro and in vivo anti-allergic effects of "benifuuki" green tea containing O-methylated catechin and ginger extract. *Cytotechnology*, 2007 55, 135-142.
- Maina, G; Vitalucci, A; Gandolfo, S; Bogetto, F. Comparative efficacy of SSRIs and amisulpride in burning mouth syndrome: a single-blind study. *J. Clin. Psychiatry*, 2003 64, 336-337.
- Martin, A; Rief, W. Somatoforme Störungen. In: Rief W, Birbaumer N editors. *Biofeedback. Grundlagen, Indikationen, Kommunikation, praktisches Vorgehen in der Therapie*. Stuttgart-New York: Schattauer Verlag; 2006; 73-96.
- Masuda, A; Nakayama, T; Yamanaka, T; Hatsutanmaru, K; Tei, C. Cognitive behavioral therapy and fasting therapy for a patient with chronic fatigue syndrome. *Internal Medicine*, 2001 40, 1158-1161.
- Matsumoto, H; Ochi, M; Abiko, Y; Hirose, Y; Kaku, T; Sakaguchi, K. Pulsed electromagnetic fields promote bone formation around dental implants inserted into the femur of rabbits. *Clin. Oral. Implant Res.*, 2000 11, 354-360.
- Mawdsley, JE; Jenkins, DG; Macey, MG; Langmead, L; Rampton, DS. The effect of hypnosis on systemic and rectal mucosal measures of inflammation in ulcerative colitis. *Am. J. Gastroenterol.*, 2008 103, 1460-1469.
- Mazurat, NM. and Mazurat, RD. Discuss before fabricating: Communicating the realities of partial denture therapy. Part II: Clinical outcomes. *J. Can. Dent. Assoc.* 2003 69, 90-94.
- Mazzetto, MO; Carrasco, TG; Bidinelo, EF; de Andrade Pizzo, RC; Mazzetto, RG. Low intensity laser application in temporomandibular disorders: a phase I double- blind study. *Cranio*, 2007 25, 186-192.
- McCaffrey, R. and Flower, NL. Qigong practice. A pathway to health and healing. *Holist Nurs. Pract.*, 2003 17, 110-116.

- McKay, DL. and Blumberg, JB. The role of tea in human health: An update. *J. Am. Coll Nutr.*, 2002 21, 1-13.
- McKee, MG. Spontaneous hypnotic states induced during biofeedback training. In: Pajntar M, Roškar E, Lavrič M editors. *Hypnosis in psychotherapy and psychosomatic medicine*. Kranj: Slovenian Society for Clinical and Experimental Hypnosis; 1980; 347-351.
- McNamara, DC. Electrodiagnosis at median occlusal position for human subjects with mandibular joint syndrome. *Arch. Oral. Biol.*, 1976 21, 325-328.
- Meier, W; Klucken, M; Soyka, D; Bromm, B. Hypnotic hypo- and hyperalgesia: divergent effects on pain ratings and pain-related cerebral potentials. *Pain*, 1993 53, 175-181.
- Meisenhelder, JB. and Chandler, EN. Prayer and health outcomes in church members. *Altern Ther Health Med*, 2000 6, 56-60.
- Melzack, R. and Wall, P. Pain mechanisms: a new theory. *Science*, 1965 150, 971-979.
- Mentes, BB; Tascilar, O; Tatlicioglu, E; Bor, MV; Isman, F; Turkozkan, N; Celebi, M. Influence of pulsed electromagnetic fields on healing of experimental colonic anastomosis. *Dis. Colon Rectum.*, 1996 39, 1031-1038.
- Michalsen, A; Weidenhammer, W; Melchart, D; Langhorst, J; Saha, J; Dobos, G. Kurzzeitiges therapeutisches Fasten in der Behandlung von chronischen Schmerz- und Erschöpfungssyndromen. Verträglichkeit und Nebenwirkungen mit und ohne begleitende Mineralstoffergänzung. *Forsch Komplementarmed Klass Naturheilk*, 2002 9, 221-227.
- Mikami, DB. A review of psychogenic aspects and treatment of bruxism. *J. Prosthet Dent*, 1977 37, 411-419.
- Mills, WW. and Farrow, JT. The transcendental meditation technique and acute experimental pain. *Psychosom Med*, 1981 43, 157-164.
- Miltner, W; Braun, C; Revenstorf, D. Nozizeption ist nicht gleich Schmerz. Eine Studie über schmerzreizkorrelierte hirnelektrische Potentiale unter Hypnose. *HyKog*, 1992 10, 22-34.
- Miyamoto, SA. and Ziccardi, VB. Burning mouth syndrome. *Mt Sinai J. Med.*, 1998 65, 343-347.
- Molin, C. From bite to mind: TMD -- a personal and literature review. *Int. J. Prosthodont*, 1999 12, 279-288.
- Moore, R; Abrahamsen, R; Brødsgaard, I. Hypnosis compared with group therapy and individual desensitization for dental anxiety. *Eur. J. Oral. Sci.*, 1996 104, 612-618.
- Moore, LB; Goodwin, B; Jones, SA; Wisely, GB; Serabjit-Singh, CJ; Willson, TM; Collins, JL; Kliever, SA. St John's wort induced hepatic drug metabolism through activation of the pregnane X receptor. *Proc. Natl. Acad. Sci. USA*, 2000 97, 7500-7502.
- Moulton, R; Ewen, S; Thieman, W. Emotional factors in periodontal disease. *Oral. Surg. Oral. Med. Oral. Pathol.*, 1957 5, 833-860.
- Musaev, AV; Guseinova, SG; Imamverdieva, SS. The use of pulsed electromagnethic fields with complex modulation in the treatment of patients with diabetic polyneuropathy. *Neurosci. Behav. Physiol.*, 2003 33, 745-752.
- Navarro, R; Marquezan, M; Cerqueira, DF; Silveira, BL; Correa, MS. Low-level-laser therapy as an alternative treatment for primary herpes simplex infection: a case report. *J. Clin. Pediatr. Dent.*, 2007 31, 225-228.
- Niepoth, L. and Korn, HJ. Schlafstörungen. In: Rief W, Birbaumer N editors. *Biofeedback. Grundlagen, Indikationen, Kommunikation, praktisches Vorgehen in der Therapie*. Stuttgart-New York: Schattauer Verlag; 2006; 250-269.

- Orme-Johnson, D. Transcendental meditation does not predispose to epilepsy. *Med. Hypotheses*, 2005 65, 201-202.
- Orme-Johnson, DW; Schneider, RH; Son, YD; Nidich, S; Cho, ZH. Neuroimaging of meditation's effect on brain reactivity to pain. *Neuroreport*, 2006 17, 1359-1363.
- Orne, MT. The nature of hypnosis: Artifact and essence. *J. Abnorm. Social Psychol.*, 1959 58, 277-299.
- Orne, MT. and Whitehouse, WG. Relaxation techniques. In: Fink G, editor in chief. *Encyclopedia of Stress*. (Vol. III.) San Diego: Academic Press; 2000; 341-348.
- Ortman, LF; Casey, DM; Deers, M. Bioelectric stimulation and residual ridge resorption. *J. Prosthet. Dent*, 1992 67, 67-71.
- Ospina, MB; Bond, TK; Karkhaneh, M; Tjosvold, L; Vandermeer, B; Liang, Y; Bialy, L; Hooton, N; Buscemi, N; Dryden, DM; Klassen, TP. *Meditation practices for health: State of research. Evidence Report/Technology Assessment No. 155*. (AHRQ Publication No. 07-E010). Rockville MD: Agency for Healthcare Research and Quality; 2007. 27-53.
- Ozcelik, O; Haytac, MC; Kunin, A; Seydaoglu, G. Improved wound healing by low-level laser irradiation after gingivectomy operations: a controlled clinical pilot study. *J. Clin. Periodontol.*, 2008 35, 250-254.
- Ozen, T; Orhan, K; Gorur, I; Ozturk, A. Efficacy of low level laser therapy on neurosensory recovery after injury to the inferior alveolar nerve. *BMC Head Face Med.*, 2006 2, 3.
- Patino, O; Grana, D; Bolgiani, A; Prezzavento, G; Mino, J; Merlo, A; Benaim, F. Pulsed electromagnetic fields in experimental cutaneous wound healing in rats. *J. Burn Care Rehabil.*, 1996 17, 528-531.
- Payer, M; Jakse, N; Pertl, C; Truschnegg, A; Lechner, E; Eskici, A. The clinical effect of LLLT in endodontic surgery: A prospective study on 72 cases. *Oral. Surg. Oral. Med. Oral. Pathol. Oral. Radiol. Endod.*, 2005 100, 375-379.
- Pekala, RJ. and Kumar, VK. Phenomenological patterns of consciousness during hypnosis: Relevance to cognition and individual differences. *Aust. J. Clin. Exp. Hypn*, 1989 17, 1-20.
- Pienkowski, D; Pollack, SR; Brighton, CT; Griffith, NJ. Comparison of asymmetrical and symmetrical pulse waveforms in electromagnetic stimulation. *J. Orthop. Res.*, 1992 10, 247-255.
- Pockaj, BA; Loprinzi, CL; Sloan, JA; Novotny, PJ; Barton, DL; Hagenmaier, A; Zhang, H; Lambert, GH; Reeser, KA; Wisbey, JA. Pilot evaluation of black cohosh for the treatment of hot flashes in women. *Cancer Invest*, 2004 22, 515-521.
- Pollack, R. Wisdom versus knowledge: an agenda for a more humane medical science. *FASEB J.*, 1999 13, 1477-1480.
- Pomp, AM. Psychotherapy for the myofascial pain-dysfunction syndrome: a study of factors coinciding with symptom remission. *JADA*, 1974 89, 629-632.
- Qadri, T; Miranda, L; Tunér, J; Gustafsson, A. The sort-term effects of low-level lasers as adjunct therapy in the treatment of periodontal inflammation. *J. Clin. Periodontol.*, 2005 32, 714-719.
- Rainville, P; Duncan, GH; Price, DD; Carrier, B; Bushnell, MC. Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science*, 1997 277, 968-971.
- Rainville, P; Hofbauer, RK; Paus, T; Duncan, GH; Bushnell, MC; Price, DD. Cerebral mechanisms of hypnotic induction and suggestion. *J. Cogn. Neurosci.*, 1999 11, 110-125.

- Raji, AM. An experimental study of the effects of pulsed electromagnetic field (Diapulse) on nerve repair. *J. Hand Surg. Br.*, 1984 9, 105-112.
- Ray, WR; Raczyński, JN; Rogers, T; Kimball, WH. *Evaluation of clinical biofeedback*. New York: Plenum Press; 1979.
- Reeves 2nd, JL. and Merrill, RL. Diagnostic and treatment challenges in occlusal dysesthesia. *J. Calif. Dent. Assoc.*, 2007 35, 198-207.
- Rehfish, HP. and Basler, HD. Entspannung und Imagination. In: Basler HD, Franz K, Kröner-Herwig B, Rehfish HP editors. *Psychologische Schmerztherapie*. Berlin-Heidelberg-New York: Springer; 2004. 537-550.
- Robelin, M. and Rogers, PJ. Mood and psychomotor performance effects of the first, but not of subsequent, cup-of-coffee equivalent doses of caffeine consumed after overnight caffeine abstinence. *Behav. Pharmacol.*, 1998 9, 611-618.
- Rogers, CR. The therapeutic relationship: Recent therapy and research. *Australian J. Psychol.* 1965 13, 95-108.
- Ross, GL; Bentley, HJ; Greene Jr., GW. The psychosomatic concept in dentistry. *Psychosom Med*, 1953 15, 168-173.
- Ryan, W; Green, CS; Laskin, DM. Comparison of diazepam, chlorazepate, carisoprodol and placebo in the treatment of MPD syndrome (abstract). *J. Dent Res.*, 1985 64 (spec issue), 232.
- Sarlani, E; Balciunas, BA; Grace, EG. Orofacial pain - Part I. Assessment and management of musculoskeletal and neuropathic causes. *AACN Clin. Issues*, 2005/a 16, 333-346.
- Sarlani, E; Balciunas, BA; Grace, EG. Orofacial pain - Part II. Assessment and management of vascular, neurovascular, idiopathic, secondary, and psychogenic causes. *AACN Clin. Issues*, 2005/b 16, 347-358.
- Scardino, MS; Swaim, SF; Sartin, EA; Steiss, JE; Spano, JS; Hoffman, CE; Coolman, SL; Peppin, BL. Evaluation of treatment with a pulsed electromagnetic field on wound healing, clinicopathologic variables, and central nervous system activity of dogs. *Am. J. Vet. Res.*, 1998 59, 1177-1181.
- Schmid-Schwap, M, Simma-Kletschka, I, Stockner, A; Sengstbratl, M; Gleditsch, J; Kundi, M; Piehslinger, E. Oral acupuncture in the therapy of craniomandibular dysfunction syndrome - a randomized controlled trial (RCT). *Wien Klin Wochenschr*, 2006 118, 36-42.
- Schmierer, H. *Einführung in die zahnärztliche Hypnose*. Berlin: Quintessenz; 1997; 236-244, 245-250, 253-275, 283-286.
- Schmierer, H. Rückmeldung nach 12 Jahren. Zahnextraction bei einer Patientin mit Trigeminusneuralgie. In: Ebell H, Schukall H editors. *Warum therapeutische Hypnose? Fallgeschichten aus der Praxis von Ärzten und Psychotherapeuten*. München-Berlin-Düsseldorf-Heidelberg: Richard Pflaum Verlag; 2004; 165-168.
- Schulte, W. Conservative treatment of occlusal dysfunctions. *Int. Dent J.*, 1988 38, 28-39.
- Schultz, JH: *Das Autogene Training. (Konzentrierte Selbstentspannung)*. Leipzig: Georg Thieme Verlag; 1932.
- Schwarz, F; Aoki, A; Becker, J; Sculean, A. Laser application in non-surgical periodontal therapy: a systematic review. *J. Clin. Periodontol.*, 2008 35(8 Suppl), 29-44.
- Schwichtenberg, J. and Doering, S. Success of referral in a psychosomatic-psychotherapeutic outpatient unit of a dental school. *Z Psychosom Med. Psychother*, 2008 54, 285-292.

- Segelman, AB; Segelman, FP; Karliner, J; Sofia, D. Sassafras and herb tea. Potential health hazards. *JAMA*, 1976 236, 477.
- Shapiro, D; Cook, IA; Davydov, DM; Ottaviani, C; Leuchter, AF; Abrams, M. Yoga as a complementary treatment of depression: effects of traits and moods on treatment outcome. *eCAM*, 2007 4, 493-502.
- Shirani, AM; Gutknecht, N. Taghizadeh, M; Mir, M. Low-level laser therapy and myofacial pain dysfunction syndrome: a randomized controlled clinical trial. *Lasers Med Sci*, in press. DOI: 10.1007/s10103-008-0624-5.
- Shor, RE. Three dimensions of hypnotic depth. *Int. J. Clin. Exp. Hypn*, 1962 10, 23-38.
- Shugars, DC. and Wahl, SM. The role of the oral environment in HIV-1 transmission. *JADA*, 1998 129, 851 - 858.
- Siegel, RK. Herbal intoxication. Psychoactive effects from herbal cigarettes, tea, and capsules. *JAMA*, 1976 236, 473-476.
- Siegel, RK. Hallucinations. *Scientific American*, 1977 237, 132-140
- Siegel, RK. Ginseng abuse syndrome. Problems with the panacea. *JAMA*, 1979 241, 1614-1615.
- Siegel, RK. and Jarvik, ME. Drug induced hallucinations in animals and man. In: Siegel RK, West LJ editors. *Hallucinations: Behavior, experience, and theory*. New York: Wiley; 1975; 81-161.
- Smith, GC. Psychotherapy. In: Fink G, editor in chief. *Encyclopedia of stress*. (Vol. 3.). San Diego: Academic Press; 2000; 310-315.
- Smith, A. Effects of caffeine on human behavior. *Food Chem. Toxicol.*, 2002 40, 1243-1255.
- Smith, AP. Caffeine, cognitive failures and health in a non-working community sample. *Hum Psychopharmacol Clin. Exp.*, 2009 24, 29-34.
- Smith, TL; Wong-Gibbons, D; Maultsby, J. Microcirculatory effects of pulsed electromagnetic fields. *J. Orthop. Res.*, 2004 22, 80-84.
- Smith, C; Hancock, H; Blake-Mortimer, J; Eckert, K. A randomized comparative trial of yoga and relaxation to reduce stress and anxiety. *Compl. Ther. Med.*, 2007 15, 77-83.
- Souil, E; Capon, A; Mordon, S; Dinh-Xuan, AT; Polla, BS. Bachelet, M. Treatment with 815-nm diode laser induces long-lasting expression of 72-kDa heat shock protein in normal rat skin. *Brit. J. Dermatol.*, 2001 144, 260-266.
- Staats, J. and Krause, WR. *Hypnotherapie in der zahnärztlichen Praxis*. Heidelberg: Hüthig; 1995; 87-104.
- Stark, TM. and Sinclair, PM. Effect of pulsed electromagnetic fields on orthodontic tooth movement. *Am. J. Orthod. Dentofacial. Orthop.*, 1987 91, 91-104.
- Stiller, MJ; Pak, GH; Shupack, JL; Thaler, S; Kenny, C; Jondreau, L. A portable pulsed electromagnetic field (PEMF) device to enhance healing of recalcitrant venous ulcers: a double-blind, placebo-controlled clinical trial. *Br. J. Dermatol.*, 1992 127, 147-154.
- Tartaro, J; Luecken, LJ; Gunn, HE. Exploring heart and soul: Effects of religiosity/spirituality and gender on blood pressure and cortisol stress response. *J. Health Psychol.*, 2005 10, 753-766.
- Tepper, OM; Callaghan, MJ; Chang, EI; Galiano, RD; Bhatt, KA; Baharestani, S; Gan, J; Simon, B; Hopper, RA; Levine, JP; Gurtner, GC. Electromagnetic fields increase in vitro and in vivo angiogenesis through endothelial release of FGF-2. *FASEB J.*, 2004 18, 1231-1233.

- Tindle, HA; Wolsko, P; Davis, RB; Eisenberg, DM; Phillips, RS; McCarthy, EP. Factors associated with the use of mind body therapies among United States adults with musculoskeletal pain. *Compl. Ther. Med.*, 2005 13, 155-164.
- Tode, T; Kikuchi, Y; Hirata, J; Kita, T; Nakata, H; Nagata, I. Effect of Korean red ginseng on psychological functions in patients with severe climacteric syndromes. *Int. J. Gynecol. Obstet.*, 1999 67, 169-174.
- Tóth, Zs. and Fábíán, TK. Relaxation techniques in dentistry. *Fogorv Szle*, 2006 99, 15-20.
- Toyofuku, A. and Kikuta, T. Treatment of phantom bite syndrome with milnacipran - a case series. *Neuropsych. Dis. Treatment*, 2006 2, 387-390.
- Tönnies, S. Entspannung für Tinnitusbetroffene durch Photostimulation. *HNO*, 2006 54, 481-486.
- Turner, JA; Mancl, L; Aaron, LA. Short- and long-term efficacy of brief cognitive-behavioral therapy for patients with chronic temporomandibular disorder pain: A randomized, controlled trial. *Pain*, 2006 121, 181-194.
- Vachiramon, A. and Wang, WC. Acupuncture and acupressure techniques for reducing orthodontic post-adjustment pain. *J. Contemp. Dent. Pract.*, 2005 1, 163-167.
- Vanhaudenhuyse, A; Boly, M; Laureys, S; Faymonville, ME. Neurophysiological correlates of hypnotic analgesia. *Contemp Hypnosis*, 2009 26, 15-23.
- Varga, K; Józsa, E; Bányai, ÉI; Gösi-Greguss, AC. Phenomenological experiences associated with hypnotic susceptibility. *Int. J. Clin. Exp. Hypn*, 2001 49, 19-29.
- Vescovi, P; Merigo, E; Manfredi, M; Meleti, M; Fornaini, C; Bonanini, M; Rocca, JP; Nammour, S. Nd:YAG laser biostimulation in the treatment of bisphosphonate-associated osteonecrosis of the jaw: clinical experience in 28 cases. *Photomed Laser Surg.*, 2008 26, 37-46.
- Vigh B. *Yoga and science [Jóga és tudomány]*. Budapest: Gondolat; 1972. 447-449.
- Vignes, M; Maurice, T; Lante, F; Nedjar, M; Thethi, K; Guiramand, J; Recasens, M. Anxiolytic properties of green tea polyphenol (-)-epigallocatechin gallate (EGCG). *Brain Res.*, 2006 1110, 102-115.
- Volz, HP. Clinical efficacy of St. John's wort in psychiatric disorders other than depression. In: Müller WE, editor. *St. John's worth and its active principles in depression and anxiety*. Basel-Boston-Berlin: Birkhäuser Verlag; 2005; 133-143.
- Wachholtz, AB. and Pargament, KI. Is spirituality a critical ingredient of meditation? Comparing the effects of spiritual meditation, secular meditation and relaxation on spiritual, psychological cardiac and pain outcomes. *J. Behav. Med.*, 2005 28, 369-384.
- Wahner-Roedler, DL; Elkin, PL; Vincent, A; Thompson, JM; Oh, TH; Loehrer, LL; Mandrekar, JN; Bauer, BA. Use of complementary and alternative medical therapies by patients referred to a fibromyalgia treatment program at a tertiary care center. *Mayo Clin. Proc.*, 2005 80, 55-60.
- Walter, WG; Dovey, VJ; Shipton, H. Analysis of the electrical responses of human cortex to photic stimulation. *Nature*, 1946 158, 540-541.
- Weintraub, MI. and Cole, SP. Pulsed magnetic field therapy in refractory neuropathic pain secondary to peripheral neuropathy: Electrodiagnostic parameters - Pilot study. *Neurorehab Neural Repair*, 2004 18, 42-46.
- Whittaker, P. Laser acupuncture: past, present, and future. *Lasers Med Sci*, 2004 19, 69-80.
- Widmalm, SE. Use and abuse of bite splints. *Compendium of Continuing Education in Dentistry*, 1999 20, 249-254.

- Wiklund, IK; Mattsson, LA; Lindgren, R; Limoni, C; Effects of a standardized ginseng extract on quality of life and physiological parameters in symptomatic postmenopausal women: a double-blind, placebo-controlled trial. (Swedish Alternative Medicine Group). *Int. J. Clin. Pharmacol. Res.*, 1999 19, 89-99.
- Wilder-Smith, P. and Zimmermann, M. Analgesie mittels Transkutaner Elektrischer Nervenstimulation (TENS). *Schweiz Montsschr Zahnmed.*, 1989 99, 653-657.
- Wilder-Smith, P. Untersuchungen zur Schmerzunterdrückung mittels transkutaner elektrischer Nervenstimulation (TENS). *DZZ*, 1990 45, 356-359.
- Williams, P. and West, M. EEG response to photic stimulation in persons experienced at meditation. *Electroencephal Clin. Neurophysiol.*, 1975 39, 519-522.
- Winkelman, M. Trance states: A theoretical model and cross cultural analysis. *Ethos*, 1986 14, 174-203.
- Winkler, G. and Krause, WR. Autogenes Training und thermales Feedback unter Kurbedingungen. *Z ärztl Fortbild*, 1989 83, 1187-1188.
- Wirz-Justice, A; Benedetti, F; Berger, M; Lam, RW; Martiny, K; Terman, M; Wu, JC. Chronotherapeutics (light and wake therapy) in affective disorders. *Psychol. Med.*, 2005 35, 939-944.
- Yang, C; Chung, J; Yang, G; Chhabra, S; Lee, M. Tea and tea polyphenols in cancer prevention. *J. Nutr.*, 2000 130, 472S-478S.
- Yun, JH; Pang, EK; Kim, CS; Yoo, Yj; Cho, KS; Chai, JK; Kim, CK; Choi, SH. Inhibitory effects of green tea polyphenol (-)-epigallocatechin gallate on the expression of matrix metalloproteinase-9 and on the formation of osteoclasts. *J. Periodontal. Res.*, 2004 39, 300-307.
- Zhang, AL; Story, DF; Lin, V; Vitetta, L; Xue, CC. A population survey on the use of 24 common medicinal herbs in Australia. *Pharmacoepidemiol. Drug Saf.*, 2008 17, 1006-1013.
- Zienowicz, RJ; Thomas, BA; Kurtz, WH; Orgel, MG. A multivariate approach to the treatment of peripheral nerve transection injury: the role of electromagnetic field therapy. *Plast Reconstr Surg.*, 1991 87, 122-129.

Chapter XII

Psychogenic Complications of Making Dentures. Theoretical Background, Prevention and Treatment Possibilities

***Tibor Károly Fábián^{*1}, Anita Beck^{1,2}, Laura Gótai¹,
Péter Hermann¹ and Pál Fejérdy***

¹Clinic of Prosthetic Dentistry

²Department of Oral Biology

Semmelweis University Budapest, Faculty of Dentistry, Budapest, Hungary, EU

Abstract

Psychogenic complications of making dentures as well as consequently appearing psychogenic denture intolerance is a complex and rising problem of dentistry and presents many intricate problems, which are being tackled by various disciplines of both basic and clinical research. Estimations based on the available data and clinical experience indicate that, at least 3-4% of denture wearers suffer from psychogenic symptoms caused by the treatment procedure, insertion or wearing of fixed or removable dentures. No wonder that, there is a high amount of scientific information gathered so far, however data are rather divergent, sometimes even contradictory and there are numerous questions without any available data to answer. Present chapter is primarily dedicated to the clinical aspects of these phenomena, including clinical manifestations, diagnosis, prevention and treatment possibilities. Other relevant subject areas of this chapter include theoretical background and peculiarities of denture-related psychological and psychophysiological phenomena, background and pathomechanisms of denture induced psychosomatic manifestations, basic principles of communication and patient-nurse-dentist interrelationships. This collection of information helps the reader to be at home in scientific field of denture related psychogenic manifestations.

^{*} E-mail address: fab@fok.usn.hu.

Introduction

The problem of psychogenic denture intolerance (PDI) stood in the limelight of scientific interest in the second half of the last century. Although early complete denture related studies sometimes missed the point that psychogenic denture intolerance may also develop in relation to other kind of dentures; later on psychogenic problems related to other kinds of dentures (i.e. removable partial dentures and/or fixed dentures) were also analyzed, and lastly a "holistic view" of psychogenic denture intolerance emerged. In this approach, denture intolerance is recognized as a member of the "large family" of orofacial psychosomatic manifestations [Ross et al. 1953, Schweitzer 1964, Marxkors and Müller-Fahlbusch 1981, Müller-Fahlbusch and Marxkors 1981, Müller-Fahlbusch and Sone 1982, Newton 1984, Demmel and Lamprecht 1996, Wolowski 2000].

In general, roughly 10 percent of the dental patients may be counted as a risk patient for any psychosomatic manifestations [Doering and Wolowski 2007], from which more than a third (roughly 3-4 % of all patients with dentures) suffer from psychogenic denture intolerance [Fábíán and Fejérdy 2007, 2010]. This is a not too large but still significant proportion, and the incidence may increase [Fábíán and Fejérdy 2007, 2010, Sugawara et al. 1998]. It should be also considered that, even two or three of such patients may unbalance dentist's entire schedule [Schweitzer 1964] and can occupy a large percentage of the dentist's time [Swoope 1973]. Moreover, it also gives rise to legal proceedings in certain cases [Figgener 1996]. Thus, the importance of psychogenic denture intolerance should not be underestimated, but considered as a rather important and difficult problem of dentistry.

Concept of Psychogenic Denture Intolerance

Emergence of the Concept

The problem of psychogenic denture intolerance stood in the limelight of scientific interest because of comparatively frequent unsuccess during making complete dentures (full dentures). Although complete denture related studies sometimes missed the point that psychogenic denture intolerance may develop in relation to other kind of dentures as well (i.e. partial removable and/or fixed dentures); huge knowledge of the psychological and psychopathological mechanisms behind psychogenic denture intolerance were collected based on this approach. Importance of patient's personality structure [Seifert et al. 1962, Ismail et al. 1974] and affective state [Golebiewska et al. 1998] as well as patient's level of psychosocial stress [Swoope 1973] level of anxiety [Swoope 1973] and level of fear of death [Dolder 1956, Kranz 1956] were recognized. Similarly, the importance of various unconscious psychological processes [Dolder 1956, Kranz 1956] and several mental disorders [Levin and Landesman 1976, Drost 1978/a,b] were documented. Further, significance of dentist's personality [Langer et al. 1961, Seifert et al. 1962] and communication skills [Guckes et al. 1978, Reeve et al. 1984] and quality of patient management [Hirsch et al. 1972, Reeve et al. 1984] were acknowledged. The importance of several transference and counter-transference processes [Drost 1978/a,b] were also recognized based on such studies.

Similarly, patient's persistence and motivation [Michman and Langer 1975] as well as patient's expectations [Levin and Landesman 1976] and emotional involvement in the development of a symptom [Holland-Moritz 1980] were documented. The importance of socio-cultural factors [Lowental and Tau 1980], self-image [Silverman et al. 1976, Tau and Lowental 1980], body-image [Ament and Ament 1970, Silverman et al. 1976] and gender differences [Massler 1951, Kotkin 1985] were also recognized. The significant influence of menopausal processes [Niedermeier et al. 1979] and the importance of quantitative and qualitative parameters of saliva secretion [Niedermeier 1991] were also recognized. The significance of neuromuscular coordination and adaptation [Michman and Langer 1975, Landt 1978], multifold sensory functions of mouth and teeth [Balters 1956/a,b], several masticatory learning processes [Michman and Langer 1975] and localization of the denture (i.e. upper or lower jaw) [Seifert et al. 1962] was also acknowledged. Finally, the important correlation between the satisfaction with dentures and the patient's ability to adjust to general health [Emerson and Giddon 1955] as well as the importance of written documentation (including the mutual agreement of objectives before treatment is begun) [Swoope 1972] was also documented based on complete denture related studies primarily.

On the base of premised findings (and collaterally), psychogenic problems related to other kinds of dentures (i.e. removable partial dentures and/or fixed dentures) were also analyzed, and lastly a "holistic view" of psychogenic denture intolerance emerged. In this approach, denture intolerance is recognized as a member of the "large family" of orofacial psychosomatic manifestations. Studies based on this concept investigate both peculiarities and general aspects of denture related psychogenic symptoms from the viewpoint of orofacial psychosomatics as a whole and more general problem [Ross et al. 1953, Schweitzer 1964, Marxkors and Müller-Fahlbusch 1981, Müller-Fahlbusch and Marxkors 1981, Müller-Fahlbusch and Sone 1982, Newton 1984, Demmel and Lamprecht 1996, Wolowski 2000].

Major role of provocative life events like traumas and object loss and the consequent appearance of depression were primarily documented based on this "holistic" approach [Müller-Fahlbusch 1972, 1975, 1976, Hach et al. 1978, Holland-Moritz 1979, Fassbind 1985]. Accordingly, the significance of traumatizing dental events (frequently causing traumas and object loss) and their role as a significant provocative life event was also acknowledged [Schweitzer 1964, Miller 1970, Marxkors 1975]. The importance of learning processes (including classic/operant conditioning or model learning) [Ross et al. 1953, Newton 1984] was emphasized as well. The meaning of symptom chronification processes induced by repeated unsuccessful somatic dental treatments [Müller-Fahlbusch 1975, Wolowski 2000, Fábíán et al. 2004, Fábíán and Fejérdy 2007, 2010] as well as the significance of primary and secondary gain from illness [Ross et al. 1953] were also recognized mainly on the base of this approach.

Similarly, importance of the rather complex patient-nurse-dentist interrelationships [Miller 1970, Fábíán and Fábíán 2000, Fejérdy et al. 2004, Fábíán et al. 2007/a, 2010], meaning of the dental team's mental health [Gerbert et al 1992, Fábíán and Fejérdy 2010], as well as the significance of proper timing of the prosthodontic treatment [Müller-Fahlbusch 1992] were documented. Meaning of the impact of oral health on quality of life [Giddon and Hittelman 1980, Leao and Sheiham 1996, McGrath and Bedi 1998, 1999, Watanabe 1998], patient's oral health behavior [Weinstein et al. 1979, Kiyak 1996] and patient's oral health-

related beliefs [MacEntee 1996, McGuire et al. 2007] were recognized as well. The importance of social [Nakaminami et al. 1989], existential [Graham 2005, Fábián et al. 2006/b] and spiritual/religious [Fábián et al. 2005/b, Fábián 2007/a] aspects of psychogenic denture intolerance were also emphasized based on the latter “holistic” approach primarily.

Definition of Psychogenic Denture Intolerance (PDI)

Considering above described emergence of its concept no wonder if yet there is currently no consensus on the etiology and pathogenesis of psychogenic denture intolerance. Similarly, there is neither a prevalent definition nor a current abbreviation for convenient use in the scientific literature of psychogenic denture intolerance. Thus, a consensus definition of psychogenic denture intolerance is still keeping us waiting. Therefore, psychogenic denture intolerance will be defined in the present work as follows: Patient's refusal accepting or wearing truly prepared (standard, properly made) fixed and/or removable denture(s) because of appearing any kind of psychogenic symptom(s) in relation to the denture or to the treatment procedure [Fábián and Fábián 2000, Fábián et al. 2004, 2007/a, Fábián and Fejérdy 2010]. Further, an abbreviation derived from the first letters of psychogenic denture intolerance (namely "PDI") will also be used in the text.

Psychological Peculiarities of Mouth and Teeth

The orofacial region is an area of human privacy of special personality relevance with overproportional cortical representation, which appears to be particularly predisposed for a large number of different functional and somatoform psychosomatic disorders [Kreyer 2000]. There are several deep-seated psychological reasons for this sensitivity. The oral psychogenic manifestations (including PDI) are based on the unique psychological and dept-psychological function of the mouth, and teeth during infancy (breasting), sexuality, aggressiveness and communication [Fábián and Fábián 2000, Fábián et al. 2000, 2007/a, Fábián and Fejérdy 2010]. There is also a great importance of the symbolic values of the mouth, teeth, tongue, and face [Fábián and Fábián 2000, Fábián et al. 2000, 2007/a]. The mouth (including the teeth and tongue) also serves as a primary zone of interaction with the environment, and as such can have far-reaching emotional significance [Ament and Ament 1970, Benson 2000]. It is surely tremendously important already in the baby's life as an organ through which to receive nourishment, as an organ of pleasure, as an organ of contact with the mother, as an organ of testing, learning and understanding and as an organ of social signal [Ament and Ament 1970]. The mouth should be considered also as a highly charged erotogenic region already from the childhood [Ament and Ament 1970, Benson 2000]. The mouth (including the teeth and tongue) is also a source of contention, and when speech comes, the mouth becomes a seat of power in a new sense as well [Ament and Ament 1970].

Psychoemotional consequences of certain traumas like the experience of forced feeding by parents in the childhood, as well as the forced manipulation of the tongue blade by the physician, or a forced (or even painful) approach by the dentist (already in the early

childhood) should also be considered [Ament and Ament 1970]. The early traumatizing experience of the infant related to losing somebody (the baby must be taken off the breast) and destroying something (intake of food with the use of the teeth) is strongly coupled to the oral region and the teeth too. (Premised coupling may be especially important during the so-called oral stage and oral-sadistic stage of the psychological development [Fábián et al. 2000, Fábián et al. 2006].) Loss of milk teeth (deciduous teeth) is usually the first loss of any part of the body and the first significant injury of body image [Ament and Ament 1970]. Next, wearing orthodontic appliances may lead to shame and body image disturbances. In many cases the dental treatment may also be traumatic for adults, and the dental experience may be perceived as an "assault" which recapitulates the old feelings (i.e. from the childhood) of helplessness, aggression, defenselessness etc. [Ament and Ament 1970]. Tooth loss may profoundly affect the psychosocial well-being of patients, even those who are apparently coping well with dentures [Fiske et al. 1998, Allen and McMillan 2003]. Complete edentulousness is a serious life event which can be perceived as more stressful than retirement [Bergendal 1989, Trullson et al. 2002], because tooth loss (especially edentulousness) also symbolizes the loss of living force, and evokes a symbolic meaning of growing old, evanescence and death [Fábián and Fábián 2000, Fábián et al. 2000, 2006, 2007/a].

The importance of dental appearance to overall appearance is also rated high [Hassel et al. 2008], and tooth loss can have tremendous patient impact and social implication [Roumanas 2009]. Therefore, significant alteration of the patient's self image may occur with an increasingly worsened dental status [Fiske et al. 1998, Trullson et al. 2002, Allen and McMillan 2003]; which may lead (in the absence of dental restorations and/or dentures) to a self-image can be labeled as "becoming a deviating person" [Trullson et al. 2002]. Bad dental status also may lead to feeling of guilt and shame [Trullson et al. 2002]. Notwithstanding that, removable dentures may significantly improve oral health-related quality of life [Adam et al. 2007, Ellis et al. 2007]; living and coping with removable dentures (especially with full dentures) may lead to a self-image can be labeled as "becoming an uncertain person" [Trullson et al. 2002]. (Premised self-image may appear because of the uncertainty and insecurity when having to cope with a removable denture.) In order to manage premised uncertainty and being in control, patient often develops avoiding strategies to ensure that no-one would notice his/her removable denture [Trullson et al. 2002]. Such patient frequently keeps the hand in front of the mouth when talking or laughing, and avoids social interactions, especially when eating. These avoiding strategies contribute to restricted social participation and a further change (worsening) of self-image [Trullson et al. 2002].

Besides premised psychological and psychosocial aspects (see above), certain neurobiological aspects of the orofacial region should also be considered. Importantly, those tissues, which are of mesodermic origin in most part of the body, are of ectodermic (ectomesodermic) origin in the orofacial region indicating a strong coupling of orofacial and nervous structures. Accordingly, stem cells present in the human dental pulp has a tendency to differentiate into functionally active neurons under certain experimental circumstances [Király et al. 2009]; which may be another indication of the above mentioned relatively strong coupling of several orofacial and nervous tissues. Further, innervation of the orofacial region is originated from the brain stem (instead of spinal cord); which may also be a base of a strong influence of psycho-emotional functions on the orofacial tissues and function.

Premised peculiarities may also be responsible for the finding that, facial and masticatory muscles are highly sensitive to psycho-emotional processes, reacting earlier, stronger and longer lasting with muscle spasm to psychoemotional stress comparing to other muscles of the body [Heggendorn et al. 1979]. It is also likely that, increased activity of facial and masticatory muscles have a special extraordinary role in the attenuation (elimination due to motor activity) of psychoemotional stress [Sato et al. in press]. A similar extraordinary role in the attenuation of psychoemotional stress due to autonomic activity (i.e. similar to that of shedding tears) may also be expected in relation with the psychoemotional stress induced alterations of saliva secretion [Fábíán and Fejérdy 2010]. In addition, mouth, teeth and tongue act as organs of sense of touch temperature and taste, leading to a reach representation of these organs in the central nervous system [Fábíán et al. 2007/a]. Taking together all above data it is no wonder that, the orofacial region is affected by psychosomatic manifestations much more frequently comparing to most other parts of the body [Böning 1990].

Psychogenic Pain Manifestations

Concept and Features of Pain

Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage [Merskey 1979]. Pain is a multifaceted and multilevel phenomenon including a specific sensation, a variable emotional state, an aspect of interoception and a specific behavioral motivation [Carli 2009]; all of which are strongly influenced by personality, cultural background, as well as social and economic factors [Dworkin and Sherman 2006, Vanhaudenhuyse et al. 2009]. Previous pain experiences [Vanhaudenhuyse et al. 2009], the significance of the organ involved [Vanhaudenhuyse et al. 2009], local pain threshold of tissues [Ogawa et al. 2004] as well as individual's current pain tolerance, mood (depression) and level of anxiety also influence pain experience significantly [Dworkin and Sherman 2006]. There are three major components of pain such as psychogenic-, neuropathic- and nociceptive pain components [Burton 1969, Gerbershagen 1995, Baldry 2005]. Importantly, pain of vascular and muscular origin also belong to nociceptive pain, because nociceptors of blood vessels or muscles (or other related tissues) are the primary sources of pain impulses in these cases [Baldry 2005]. In some cases pain symptom is solely due to a single pain component; in other cases two or three components may be present concomitantly [Burton 1969, Gerbershagen 1995, Baldry 2005].

Radiation of Pain

Local radiation of pain leading to local (near) referred pain is a frequent concomitant phenomenon of pain, especially when deep-seated pain appears (i.e. painful teeth radiating pain into other teeth or the ear). Distant radiation of pain may also occur, according to the location of dermatomes which are skin areas having sensory innervations from the same central nervous segment. Radiating visceral pain usually does not cover the whole dermatome

surface, but specific smaller areas of the dermatome, which are referred to as Head-zones [Hansen and Schliack 1962, Gerbershagen 1995]. Although the "classical" dermatomes typically belong to the spinal cord segments; brainstem is also considered as a "segment" of the central nervous system in this relation, and skin innervations areas of the branches of trigeminal nerve are considered as dermatomes of the facial region [Gerbershagen 1995]. There are also smaller facial skin areas located roughly concentric around the mouth called „trigeminal projection fields" [Gerbershagen 1995] which are considered as facial Head zones.

Pain Chronification Processes

Pain chronification processes are also highly important in relation with orofacial pain. Whenever increased activity of pain related afferent fibers and other neurons of pain related centers appears, profound intracellular changes occur in few hours (but latest in few days). Premised profound changes are referred to as activity dependent neuronal plasticity which includes activation of immediate early genes and several transcription factors as well as altered synthesis of neurotransmitters, receptor precursors, or secondary messengers like kinases or phosphatases (etc.) [Hunt et al. 1987, Besson 1999, Zimmermann 2004]. Although premised changes are reversible at the beginning, they may become irreversible in the course of time (roughly in months). Whenever irreversible changes appeared, the pain sensation may become independent from the primary cause and may become chronic, because at this stage neurons may fire spontaneously in an irreversible manner.

Atypical Facial Pain

Atypical facial pain is an idiopathic symptom. It is a diagnosis of exclusion, after other conditions have been considered and eliminated. Although any area of the face can be involved, the most commonly affected areas are the maxillary region [Sarhani et al. 2005/b] and other non-muscular non-joint orofacial areas [Köling 1998]. Atypical facial pain may also appear in relation with PDI [Fábián 1999, Fábián et al. 2006/a, Fábián and Fejérdy 2010]. Although atypical facial pain is considered as idiopathic [Sarhani et al. 2005/a,b], psychological distress, anxiety and depression are prevalent among PDI related atypical facial pain patients.

Myofascial Pain

Myofascial pain is characterized by the presence of focal, exquisitely tender muscle areas called trigger points typically found in the taut muscle band and producing a characteristic pain referral pattern on palpation. Besides trigger points, myofascial pain may also lead to the appearance of heterotopic orofacial pain as well as to the pain of the temporomandibular joint and to denture soreness. Major cause of PDI related myofascial pain is the overload (spasm and muscle fatigue) of masticatory muscles because of neuromuscular dysfunction induced by

complex psychogenic processes [Schwartz 1955, Laskin 1969, Dahlström 1989, Sieber et al. 2003]. Disturbed relationships of the teeth and/or occlusal surfaces, altered occlusal guidance as well as poorly occluding and/or ill fitting dentures also may trigger neuromuscular malfunctions.

Glossodynia and Oropyrrosis (Burning Mouth Sy.)

Burning mouth syndrome (BMS) is a distinct clinical entity characterized by a chief complaint of unremitting oral burning concomitant with no oral mucosal clinically observable lesions, or other relevant apparent organic basis. It is usually milder upon awakening and progressively increases in the course of the day [Sarlanı et al. 2005/b]. Insertion of the denture increases the pain intensity in many cases. Although PDI related burning mouth syndrome can be induced via psychogenic pathways; data suggest a complex interaction between several general health factors, psychosocial stressors and denture dysfunction in order to explain burning mouth syndrome in most cases [Swensson and Kaaber 1995, Aneksuk 1989, Grushka and Sessle 1991].

Tooth-Located Psychogenic Pain

Tooth-located pain of primarily psychogenic origin is called atypical odontalgia [Sarlanı et al. 2005/b, Dworkin and Sherman 2006, Baad-Hansen 2008]. The pathomechanism of atypical odontalgia is far not clear yet; however, alteration of catecholamine (epinephrine and norepinephrine) [Nagy et al. 2000] and HSP70/HSPA type stress protein [Fábíán et al. 2009/a] levels in the dental pulp may play an important role. Although there are only few published cases in the literature, tooth located pain of psychogenic origin is a rather frequent symptom of PDI patients [Müller-Fahlbusch 1991, Fábíán 1999, Fábíán and Fejérdy 2010]. The pain can be centered on apparently normal teeth, endodontically treated teeth, abutment teeth [Sarlanı et al. 2005/b] and on dental implants [Kromminga et al. 1991]. Extraction site (alveolar bone) of a previously extracted tooth (or removed implant) is also frequently targeted and referred to as phantom tooth pain [Fábíán 1999, Sarlanı et al. 2005/b].

Psychogenic Neuromuscular Symptoms

Specificities of Orofacial Muscles

Masticatory and facial muscles are highly sensitive muscles having a rather extended representation in the motor cortices comparing to other muscles [Sergl 1996]. Besides motor cortices, other brain centers including limbic system, thalamus, cerebellum, as well as the basal ganglia and brainstem reticular formation are also involved in the neuromuscular regulation of facial and masticatory muscles [Harris and Griffin 1975]. In contrast to most other muscles, which are supplied via the spinal cord, motor nervous supplies of masticatory

and facial muscles are directly from the brainstem via several cranial nerves. Brainstem is a structure mediating numerous important psychoemotional processes such as pain, arousal, attention, mood (i.e. depression), anxiety as well as defensive and reproductive behaviors [Clark et al. 2006]. Therefore, brainstem seems to be an ideal structure to regulate muscles having unique functions strongly interrelated with emotions including speech (chewing muscles, tongue-, lip-, and facial muscles), expressing emotions nonverbally (facial muscles), fighting and aggressiveness (chewing muscles) or sexuality (lips, tongue). Thus, it is no wonder that, facial and masticatory muscles are highly sensitive to psycho-emotional processes [Heggendorn et al. 1979, Graber 1995, Fuhr and Reiber 1995], and increase of their tone (muscle tension) occurs rather frequently [Heggendorn et al. 1979, Graber 1995, Fuhr and Reiber 1995]. Accordingly, their reaction (increase of tension) to stress is stronger and longer lasting comparing to other muscles of the body, which is expected to be a basis of the high frequency of occurrence of myofascial pain [Heggendorn et al. 1979]. Since stress induced increase of muscle tension is usually higher in elevators comparing to depressors [Heggendorn et al. 1979, Graber 1995, Fuhr and Reiber 1995], increased muscular tone may also lead to the decrease of free way space; which may trigger parafunction due to the increasing frequency of occurrence of spontaneous contact between antagonist teeth [Troest 1995].

Occlusal Discomfort and Dysesthesia

Occlusal discomfort may be induced by occlusal interferences especially in sensitive patients with increased vulnerability [Le Bell et al. 2006]. Oral parafunctions and/or increased muscle tension leading to fatigue in masticatory muscles can also result in occlusal discomfort [Mew 2004, Litter 2005] because of the failure of proper proprioceptive feedback [Litter 2005]. In more severe forms occlusal complaint may appear as an occlusal dysesthesia (OD) [Baba et al. 2005, Clark et al. 2005], which is a collective notion of several uncomfortable feelings of occlusion despite the absence of any observable occlusal anomaly or discrepancy [Fábián et al. 2005, Baba et al. 2005]. These patients are preoccupied with their dental occlusion, believing that, it is abnormal [Jagger and Korszun 2004], therefore occlusal dysesthesia is frequently also referred to as "phantom bite" or "occlusal neurosis" in the literature. The condition is remarkable for the nature of the involved explanations and interpretations that the patients give and for their persistence in trying to find a "dental" or "occlusal" solution [Jagger and Korszun 2004] or a "bite correction" [Toyofuku and Kikuta 2006]. However, available evidence suggests that, the symptoms cannot be improved by occlusal treatments; therefore, it is essential to avoid irreversible restorative treatment [Jagger and Korszun 2004, Toyofuku and Kikuta 2006].

Instability of Removable Denture

Stability of dentures is highly important for patient satisfaction with removable dentures. Increased vertical dimension and improper determination of the horizontal dimension of

occlusion as well as occlusal interferences and unfavorable denture bearing tissues may frequently lead to instability of removable dentures (especially of complete dentures) [Griffin and Harris 1975, Gonzalez and Desjardins 1983]. Inadequate border seal [Kivovics et al. 2007], overextension or underextension of the denture base as well as the improper relation of teeth to the ridge and inadequate contour of the artificial gingiva may also lead to denture instability [Gonzalez and Desjardins 1983]. Use of canine guidance instead of bilateral balanced occlusion may also trigger instability of complete dentures [Rehmann et al. 2008]. Certain patients, especially edentulous patients also complain of lack of adequate space of tongue movement and consequent instability of denture [Griffin and Harris 1975]. Changes of saliva secretion [Niedermeier et al. 2000, Wolff et al. 2003] and alterations of neuromuscular function may also be a cause of instability of removable (especially complete) dentures. Since psychogenic factors including psychosocial stress and other psychoemotional conditions may significantly alter both neuromuscular function and saliva secretion, an appearance of instability of dentures as a symptom of psychogenic origin of PDI also frequently occur.

Bruxism

Bruxism can be defined as a nonfunctional contact of the teeth during grinding, gnashing, tapping or clenching. Malocclusions including occlusal early contacts and other occlusal disharmonies as well as improper determination of the vertical dimension and/or other alterations of the maxillomandibular relationship [Biondi and Picardi 1993, Le Bell et al. 2006] may induce bruxism. Ill-fitting removable dentures also frequently induce clenching [Sandler et al. 1995]. Malocclusions and denture related discrepancies are likely to trigger ancient mechanisms similar to those being responsible for the elimination of impacted occlusal foreign bodies in animals (i.e. impacted pieces of bones, or small pebbles) via biting on it with strong biting forces (to crush by pressing) and/or via intermittent isometric grinding movements (to grind, to „erode” it). Importantly, psychosocial factors [Biondi and Picardi 1993, Kampe et al. 1997] and psycho-emotional conflicts [Somer 1991, Biondi and Picardi 1993] may also be responsible for the appearance of bruxism. Increased level of anxiety and lower level of socialization also increases the incidence of bruxing behavior [Kampe et al. 1997]. Premised psychogenic factors are likely to be responsible for PDI coupled bruxism cases.

Accidental Biting on Soft Tissues

Inadequacies of dentures including instability, decreased vertical dimension, insufficient horizontal overlap and shaping of teeth, improper position of teeth, and poor location of plane of occlusion may frequently lead to biting of tongue, cheek and lip [Gonzalez and Desjardins 1983, Piquero et al. 1999] and mucosa ridging [Piquero et al. 1999]. Accidental biting on tongue (and other soft tissues) also may occur as soft tissue parafunction of neuromuscular origin. Therefore, in PDI related psychogenic cases psychosocial stress effects and consequent neuromuscular dysfunction rather than denture mistakes and organic diseases are

the major cause of such manifestations. In such cases stress induced malfunction of preventive reflex mechanisms such as the inhibitory influence of jaw muscle proprioceptors on genioglossus activity (which may lead to accidental bite on tongue) [Sauerland and Mizuno 1970] as well as a decreased (or absence of) inhibitory effect of the mechanical stimulation of tooth on tongue protrusion [Schmitt et al 1973] can be expected as a concrete cause [Griffin and Harris 1975].

Overactive Gag Reflex ("Gagging")

The term gagging refers to a defense reflex, which attempts to eject unwanted, irritating, or toxic materials from the upper gastrointestinal tract. Although gag reflex is a normal defense function of human, overactive gag reflex ("gagging") may be rather disadvantageous for denture wearers, especially for those wearing removable dentures. Gagging may also lead to failure to tolerate making impression and/or other routine dental procedures. There can be numerous mechanisms behind gagging including several gastrointestinal and central nervous defense pathways; but denture related gagging are primarily induced by several conditioning processes, or several neurotic psychological pathways [Savage and McGregor 1970, Newton 1984, Gáspár et al. 2002]. Conditioning processes are typical of gagging caused by learned avoidance reactions originating from former aversive encounters with dental treatment [Eli and Kleinhaus 1997]; whereas neurotic psychological pathways are typical of gagging caused by deep-seated defense mechanisms, which serve as inadequate solutions to several psychodynamic conflicts [Eli and Kleinhaus 1997, Gáspár et al. 2002].

Other Psychogenic Symptoms

Mucosal Allergic Reactions

Allergic and/or irritative reactions of the oral mucosa primarily occur due to wearing fixed- and/or removable dentures prepared from dental cast alloys and/or acrylic resins. Other dental materials like amalgam and rarely composites may also induce such reactions. These reactions usually belong to delayed type (type IV) allergic reactions which develop in three major phases as follows: (1) metal ions or other small molecule allergens (referred to as haptens) bind to proteins and hapten-protein-complexes develop [De Rossi and Greenberg 1998, Büdinger and Hertl 2000, Martin 2004]; (2) hapten-protein-complexes are presented by antigen presenting cells inducing T-lymphocyte activation and consequent proliferation/accumulation of a specific T-lymphocyte population in lymph nodes [De Rossi and Greenberg 1998, Büdinger and Hertl 2000, Martin 2004]. Premised T-lymphocyte population is specifically sensitized and possess hapten-protein-complex-specific antigen receptors on their surfaces [De Rossi and Greenberg 1998, Büdinger and Hertl 2000, Martin 2004]; therefore, this phase is referred to as sensitization phase; (3) if hapten-protein-complexes at issue appear again, trafficking (recirculating) sensitized T-lymphocytes become reactivated and start to release proinflammatory cytokines and chemokines which lead to cellular immune-inflammatory reactions in the tissue [De Rossi and Greenberg 1998,

Büdinger and Hertl 2000, Martin 2004]. Accordingly, this phase is referred to as effector phase [Borelli et al. 1988]. The occurrence of premised effector phase is the primary cause of most denture related mucosal allergic reactions, as well as for various allergic reactions including contact eczemas, allergic dermatitis and certain granulomatous reactions [De Rossi and Greenberg 1998, Büdinger and Hertl 2000, Borelli et al. 1988, Martin 2004]. Since higher cortical-, limbic- and neurosensory systems (receiving and integrating a great diversity of signals arising from inside and outside the body) significantly influence immunity [Chrousos 1998, Ligier and Sternberg 2000]; influence of personality, cognitions and certain psychoemotional stress states on immune responses frequently occur. Primary effector arms of the psychoemotional nervous reactions are the hypothalamus, anterior pituitary gland and adrenal cortex (referred to as hypothalamic-pituitary adrenocortical axis, HPA axis) [Chrousos 1998, Ligier and Sternberg 2000]. Other key elements are located in the brain stem, which elements influence the immune reactions primarily due to the activation of the sympathetic nervous system [Chrousos 1998, Ligier and Sternberg 2000]. Despite the expected role of HSP70/HSPA in the oral tolerance data in the literature suggest that stress induced overexpression of HSP70/HSPA may also play an important role in the appearance of stress related mucosal allergic reactions, because HSP70/HSPA proteins are likely to play a role in haptentation and consequent sensitization [Fábíán et al. 2009/a]. Although allergic reactions are primarily of somatic origin, in PDI related cases psycho-emotional stress induced central nervous processes may play a major role in the maintenance of symptoms.

Chronic Inflammations

There is a high number of several inflammatory processes, which may occur in the oral cavity in relation with preparing or wearing dentures. Some of them may be stress induced and frequently appear also as a major symptom of PDI patients. Psychosocial stress induced chronic inflammations of the oral mucosa, dental pulp, periapical tissues and periodontal tissues belong to this group primarily. Stress induced activation of HSP70/HSPA type stress proteins are likely to play pivotal role in premised stress induced oral inflammatory processes [Fábíán et al. 2007/b, 2008,b,c, 2009/a,b]; notwithstanding that HSP70/HSPA proteins also play a highly important role in the maintenance of orofacial tissues' health [Fábíán et al., 2007/b, 2008,b,c, 2009/a,b]. Oral mucosal inflammation may occur because of cross reactivity of specific antibodies against microbial HSP70 type stress proteins with human HSP70/HSPA proteins and/or because of autoantibodies against HSP70/HSPA proteins. Inflammatory processes of the dental pulp can also be triggered or worsened due to the overexpression of HSP70/HSPA proteins under stress. Disadvantageous amplification of inflammatory response to bacteria (i.e. carious dentine) and/or to several dental materials as well as to tooth preparation due to the immune enhancing effect of HSP70/HSPA proteins could be expected as a possible pathway [Fábíán et al. 2009/a]. Flare up or formation of periapical lesions may also be triggered and or worsened due to stress induced overexpression of HSP70/HSPA molecules in lymphocytes and endothelial cells of inflammatory granulation tissues, as well as in lining epithelium of radicular and residual cysts [Fábíán et al. 2009/a, Fábíán and Fejérdy 2010].

Exacerbations of periodontal inflammations are also likely to occur due to a bone resorptive effect of HSP70/HSPA proteins [Fábián et al. 2009/a,b]; after the turning of gingivitis into a more severe inflammation (i.e. periodontitis) with pocket formation and irreversible destruction of the periodontal bone.

Cognitive-Behavioral Symptoms

Besides various cognitive-behavioral symptoms caused by any psychiatric conditions of PDI patients, the most important behavioral symptom of PDI patient is overcriticism. It frequently appears as esthetical complaints of overcritical patients [Ma et al. 2008]. Overcritical patients usually demands extraordinary efforts (and/or previous guarantees of treatment outcome) at no additional cost [Ma et al. 2008], and expect treatment goals, which are very difficult or impossible to achieve [Ma et al. 2008]. Although appearance of esthetic complaints may be the most frequent, overcriticism may also occur in relation with any other aspect of denture and dental treatment certainly. Importantly, failure to recognize psychogenic factors in such cases, together with prolonged unsuccessful somatic/operative dental therapy may induce profound behavioral changes, and incite certain patients to seek relief through litigation [Harris et al. 1993].

Salivation Problems

Saliva is a major determinant of the oral environment and oral comfort. Salivary components can originate from several sources (i.e. major and minor salivary glands, blood, oral mucosal cells and oral microbes) leading to a rather complex collection of molecules [Fábián et al. 2008/a,d]. Saliva constituents play major role in several oral processes, including formation of acquired pellicle on tooth surfaces, bacterial adhesion and biofilm (plaque) formation, crystal growth homeostasis of teeth, mucosal surface protection, hard tissue surface protection and antimicrobial defense [Guggenheimer and Moore 2003, Fábián et al. 2008/a,d].

Saliva also play important role in the oral wound healing, dental caries formation, calculus formation and gingival inflammation [Fábián et al. 2008/a,d]. Quantitative and/or qualitative changes of saliva are also major cause of both subjective dry mouth sensation (xerostomia) and ptyalorrhea (ptyalism) [Fábián et al. 2008/a,d].

Salivary changes may also lead to halitosis, alteration of taste perception and alteration of phonation. Based on above data it is obvious that, salivary changes are major determinant of the oral milieu; and qualitative or quantitative alterations of saliva secretion may lead to general Oral Discomfort as well as specific oral symptoms. There can be several causes behind alterations of salivary secretion including several systemic diseases and/or their medication as well as insertion of new dentures. Importantly, occurrence of psychosocial and/or psychoemotional stress including anxiety [Somer et al. 1993] and depression [Bushfield et al. 1961, Gottlieb et al. 1961] also strongly influence salivary secretion, therefore may lead to the appearance of several saliva related manifestations including saliva related symptoms of PDI.

Halitosis (Oral Malodor, Foetor Ex Ore)

The most plausible causes of oral malodor (halitosis) are insufficient oral hygiene, periodontal disease, carious lesions [Yaegaki and Coil 1999/a,b, 2000/b, Lang and Filippi 2004, Nalcaci and Baran 2008] or non-oral causes like nose-throat disorders, several medical conditions, several drugs, or nutritional habits [Yaegaki and Coil 2000/a,b, Lang and Filippi 2004]. However, in some cases there are several psychological or psychopathological causes behind the appearance of oral malodor [Yaegaki and Coil 1999/a,b, Lang and Filippi 2004]. It may occur that patients expect but do not exhibit own oral malodor (also referred to as "pseudohalitosis" or "halitofobia") [Yaegaki and Coil 1999/b, 2000/a,b, Giddon and Anderson 2006], which may also appear as a symptom of PDI. Since the bacterial decomposition processes and resulted volatile sulfur compounds are decisively responsible for the appearance of malodor in most cases [Lang and Filippi 2004], stress induced deterioration of salivary defense functions (and consequent changes of oral microbiota) may also lead to oral malodor of psychogenic origin [Fábíán et al. 2008/a,d]. Elimination of gingival bleeding, periodontal pockets or carious lesions, and also professional dental hygiene and proper individual dental hygiene including tongue brushing are crucial to control oral malodor (halitosis) [Sanz et al. 2001] also in cases of psychogenic origin certainly. However, in cases of psychogenic origin psychosomatic therapy of patients is also needed [Yaegaki and Coil 1999/a, 2000, Lang and Filippi 2004].

Alteration of Taste Sensation

Denture induced oral inflammations, bacterial biofilm accumulation because of wearing dentures or release of chemical components from dentures (i.e. acrylate monomer [Har-Zion et al. 2004], nickel [Pfeiffer and Schwickerath 1991]) may appear as real taste stimuli. However, dentures may induce a "real" alteration of taste sensation as well. Such denture induced alterations of taste sensation may occur primarily due to six major pathways as follows: (1) prevention of contact between receptor sites and food [Har-Zion et al. 2004, Nalcaci and Baran 2008]; (2) prevention of contact of the tongue with the palatal rugae, which is considered as highly important for dispersing the food and bringing it into more intimate contact with the taste buds [Har-Zion et al. 2004]; (3) entrapment of a part of the food between the mucosa and the baseplate [Har-Zion et al. 2004]; (4) interference with chewing and usual mobility of the tongue and cheeks and consequently decreased release of taste and smell flavor stimuli [Har-Zion et al. 2004]; (5) interference with the free movement of humid and warm air in the oral and nasal cavities and consequent affection of retronasal olfaction; (6) appearance of pain and pressure or alteration of mucosal touch experience, which also may lead to taste alterations [Har-Zion et al. 2004]. Importantly, psychological stress factors may also lead to the alteration of taste sensation in PDI cases [Fábíán and Fejérdy 2010]. There can be numerous central nervous mechanisms considered behind psychological alteration of taste sensation including conditioned learning processes [Chang and Scott 1984] and modulation of glutamate-, GABA- or substance-P related orosensory

(taste related) synaptic transmission of the rostral nucleus of the solitary tract (rNST) in brainstem [Smith et al. 1998, Grabauskas and Bradley 1998].

Alteration of Speech

Speech may be altered in some patients after insertion of a new prosthesis, especially (but not exclusively) in the case of complete denture patients. Obviously, it is more difficult to adapt pronunciation of consonants to a new oral situation than vocals; because of the crucial impact of morphology and position of lips, teeth, palatal gingiva, palate and tongue during pronunciation of consonants [Molly et al. 2008]. Artificial teeth (of any dentures) primarily disturb the production of plosives and fricatives [Molly et al. 2008], whereas removable dentures (including RPDs and especially full dentures) frequently make the pronunciation of labio-dental and apico-alveolar consonants difficult [Lundqvist 1993, Molly et al. 2008]. However any other consonants [Kaán et al. 1993] (i.e. bilabial-, palatal- velar- and glottal-consonants [Molly et al. 2008]) or vocals [Seifert et al. 1999/a,b] may also be distorted following insertion of a new denture. Moreover the voice itself, and the range of the voice may also be changed because of insertion of new dentures [Seifert et al. 1999/a,b]. Importantly, stress and other psychoemotional factors strongly influence both saliva secretion and psychomotor function both of which play major role in phonation. Therefore, alteration of speech may also appear as a primarily psychogenic symptom of PDI.

Pseudoneurological and General Health Symptoms

Pseudoneurological symptoms are symptoms appearing as if they would be caused by any neurological disorder; however without any detectable somatic/neurological background. Several sensory disturbances such as hypoaesthesia, hyperaesthesia, anaesthesia, paraesthesia and dysaesthesia as well as facial palsy are the most important such symptoms in the orofacial region. Certain pain symptoms, taste related phenomena and tinnitus as well as certain autonomic and motor symptoms (i.e. facial tic, oral parafunctions) may also be classified as a member of this group. Some general symptoms including fatigue, weakness, headache, dizziness, nausea, emotional tension and irritability may occur in relation with denture intolerance as well. Psychogenic fever a common psychosomatic disease with acute or persistent body temperature above normal range under psychosocial stress may also appear as symptom of orofacial psychosomatic manifestations [Oka and Oka 2007] including PDI. Complaints by patients who attribute their symptoms to dental alloys or other dental materials are also often general in nature [Garhammer et al. 2001]; and may include various symptoms like headache, weakness, paraesthesia as well as appearance of blisters and intestinal problems [Garhammer et al. 2001]. Although pseudoneurological and general health symptoms of PDI patients are of psychogenic origin in many cases [Di Felice et al. 1991, Hampf 1987]; any possible somatic causes including both local (peripheral) and central nervous causes should also be considered and ruled out very carefully.

"Amalgam Illness"

Possible adverse health effects due to mercury released by amalgam fillings have been discussed in several studies of patients who attribute various symptoms to the effects of amalgam fillings [Lindberg et al. 1994, Malt et al. 1997, Bailer 2001, Vamnes et al. 2004]. Although the frequency of occurrence of amalgam related symptoms is not rare, in general, no systematic relation of specific symptoms to increased mercury levels could be established in such cases [Lindberg et al. 1994, Malt et al. 1997, Bailer 2001]. Although in certain cases the toxic/irritative effects may not be excluded [Scott et al. 2004, McCullough and Tyas 2008], there are primarily psychological and/or pathopsychological causes behind the "amalgam illness" [Lindberg et al. 1994, Malt et al. 1997, Bailer 2001]. In the absence of any clear toxicological and/or allergological evidence, "amalgam illness" may be seen as a label for a general tendency toward somatization [Lindberg et al. 1994, Malt et al. 1997, Bailer 2001].

Diagnosis and Differential Diagnosis

Cornerstones of Diagnostic Process

One of the most important goals of the diagnosis of psychogenic denture intolerance (PDI) is the early recognition of the disorder and the avoidance of further useless invasive dental treatment, especially because aggravation or spread of symptoms following invasive dental interventions is not uncommon. Early diagnosis (or assumption) of PDI is also crucial for the prevention of symptom chronification which frequently render pain and other symptoms intractable.

The most important diagnostic tool for such purposes is collecting detailed patient's history and careful evaluation of all relevant psychosocial, medical and dental anamnestic data including their understanding in a context of a biopsychosocial model of orofacial disorders [Ross et al. 1953, Engel 1977, Barsby 1994, Pertes 1998, Green and Laskin 2000, Dworkin and Sherman 2006, Fábíán and Fejérdy 2010]. Careful clinical examination of the oral and orofacial region and general health status as well as X-ray, MRI and other specific examinations are also cornerstones of a proper diagnostic process certainly. Referral to other professionals may also be needed, and especially recommended in the case of pain and other (pseudo)neurological symptoms. Since psychogenic symptoms may mimic a great variety of somatic symptoms, a clear-cut diagnosis and proper differential diagnosis of PDI could be rather difficult.

Therefore, it should be emphasized that, the diagnosis of PDI is a presumptive one in many cases. The diagnosis may change later in the course of the disease as the clinical findings change and/or stabilize, therefore continuous monitoring and evaluation of the patient over time is essential.

A detailed differential diagnosis considering possible disorders of the teeth, and/or oral- and maxillofacial tissues should also be carried out, and any other possible somatic causes behind the symptoms should also be excluded.

Patient's History (Anamnesis)

As previously mentioned, one of the most important (if not the most important) diagnostic tool for making diagnosis of PDI is collecting detailed patient's history [Ross et al. 1953, Barsby 1994, Pertes 1998] including a careful evaluation of all relevant psychosocial, medical and dental anamnestic data [Ross et al. 1953, Pertes 1998, Dworkin and Sherman 2006]. History taking is an art in the subtle direction of conversation with the patient. As the interview develops and the data accumulate, insight into the patient's problems becomes more apparent to the dentist, further enabling him to direct the interview along the most useful lines. History-taking is a process usually extending over a number of patient's visit to reach adequate amount of data related to the dental-, medical-, and psychosocial life events [Fábián and Fábián 2000, Fábián et al. 2007/a]. Correlation may exist between the degree of satisfaction with dentures and the patient's ability to adjust to general health problems [Emerson and Giddon 1955, Tickle et al. 1997] and/or psychosocial problems. Therefore, interrelations between psychosocial- medical- and dental history should be evaluated carefully, and the data should be understood in a context of a biopsychosocial model of disorders [Ross et al. 1953, Engel 1977, Barsby 1994, Pertes 1998, Green and Laskin 2000, Dworkin and Sherman 2006, Fábián and Fejérdy 2010].

History of symptoms that are inconsistent with physical findings and history of a precipitating life event after which the symptom began [Müller-Fahlbusch and Sone 1982, Brodine and Hartshorn 2004, Dworkin and Sherman 2006] are hallmarks of PDI cases [Müller-Fahlbusch and Sone 1982, Brodine and Hartshorn 2004]. Moreover, as a result of apparent shortcomings in primary care and preliminary diagnostics, many PDI patients typically look back on long and complicated case histories with numerous negative examinations and futile attempts at somatic therapies and/or somatic interventions [Marxkors and Müller-Fahlbusch 1981, Dworkin and Sherman 2006]. Increased number of dental clinics and hospitals visited by the patient in the previous time also refers to an increased risk of PDI. History of any dental or facial trauma may also be a sign of increased risk of appearance of psychogenic orofacial symptoms [Müller-Fahlbusch and Sone 1982, Brodine and Hartshorn 2004]. Similarly, head and neck traumas as well as survived head and/or neck cancer may also increase the risk of such manifestations significantly. History of symptom substitutions (turning of a symptom into an other one) also frequently appear in the history of PDI patients; however the role of possible environmental or other psychosocial changes should also be considered carefully before assuming an underlying pathopsychological background [Gale and Carlsson 1976].

Evaluation of Medical and Psychosocial Status

Besides orofacial status (including intraoral and dental status, see below), there are two other major facets of patients' status namely the medical- and the psychosocial status. Medical status may strongly influence both diagnostic and treatment procedures as well as the prognosis of PDI. Importantly, there is a strong association between self-reported general health status and the subjectively evaluated oral health status [Emerson and Giddon 1955,

Tickle et al. 1997] indicating that, lower level of self reported general health may predispose to Oral Discomfort and consequent oral psychosomatic manifestations [Emerson and Giddon 1955, Tickle et al. 1997]. Depression, anxiety disorder, somatization and other psychiatric disorders as well as any other diseases leading to significant suffering and/or limitations of psychosocial functioning may lead to significant treatment difficulties and poor prognosis [Dworkin and Sherman 2006].

Psychosocial status also strongly influences both treatment possibilities and the prognosis of the offered treatment. Psychosocial conditions may trigger the appearance and/or may lead to the fixation of orofacial psychosomatic problems [Ross et al. 1953, Dworkin and Sherman 2006]. Fixation of a sick role of the patients may also be rooted in several psychosocial causes [Dworkin and Sherman 2006]. Accordingly, both treatment possibilities and their prognosis are more guarded when self reported activity limitations are high. The prognosis is especially poor, if the medical and/or dental symptoms interfere appreciably with ability to discharge responsibilities (i.e. at home, in school, at work), or limit socializing activities [Dworkin and Sherman 2006]. Moreover, if there are strong extraneous social and/or health influences or any other object loss like bereavement, unemployment, placing on the retired list, diagnosis of life-threatening illness (etc.), the individual's ability to accept dentures may also be seriously compromised [Fábíán and Fejérdy 2010]. The psychoemotional "strength" and supportive potential of family members, friends and other members of the social networks of patients also of high diagnostic and prognostic importance.

Diagnosis Based on Definition

The essence of diagnosis based on definition is that, the symptom(s) should fit in the definition of denture intolerance, which was defined in the present work as: patient's refusal accepting or wearing truly prepared (standard, properly made) fixed and/or removable denture(s) because of appearing any kind of psychogenic symptom(s) in relation to the denture or to the treatment procedure [Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010]. Since psychogenic symptoms may mimic a great variety of (most of) somatic symptoms; to assess that whether a symptom is of psychogenic origin or not, the next five characteristics of psychogenic symptoms [Marxkors and Müller-Fahlbusch 1981] can be considered as follows: (1) well-marked divergence between the symptoms and the clinical findings; (2) unsuccessful previous somatic treatments; (3) fluctuation of the symptoms; (4) conspicuous emotional involvement of the patient into the dental problem; (5) presumable relationship of the symptoms and the psychosocial history [Marxkors and Müller-Fahlbusch 1981]. Symptoms which meet at least four from the above five criteria are very likely to be of psychogenic origin.

Diagnosis of Exclusion

It should be emphasized that, the diagnosis of PDI is also a diagnosis of exclusion [Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010], therefore a careful and accurate dental examination is crucial. A detailed differential diagnosis considering

possible disorders of the teeth, or oral- and maxillofacial tissues should also be carried out. PDI may be expected only at those cases without any significant dental finding. Following the careful dental examination, any other possible somatic causes behind the symptoms should also be excluded. For such purposes, exclusion of systemic diseases that may cause orofacial symptoms should also be carried out and medication induced symptoms should also be considered. Further, referral to other professionals like maxillofacial surgeon, neurologist, psychiatrist, otorhinolaryngologist, ophthalmologist, internist, rheumatologist (etc.) may also be needed, and especially recommended in the case of pain and other (pseudo)neurological symptoms [Sarlani et al. 2005/a.b]. It should be also considered that, the diagnosis may change later in the course of the disease as the clinical findings change and/or stabilize, therefore continuous monitoring and evaluation of the patient over time is essential [Sarlani et al. 2005/a.b].

Prevention of Psychogenic Denture Intolerance

Cornerstones of Prevention

Besides high quality preparative dental skills and technical background, cornerstones of prevention are screening of risk patients, proper treatment planning, proper communication with the patient, screening of the patient-nurse-dentist interrelationships, as well as prevention of dental fear, prevention of treatment-induced pain and prevention of relapse. It is also crucial for the prevention that patients must be made aware of their responsibilities in achieving a satisfactory outcome. It is also a matter of considerable significance that the clinician carefully weighs the option of nontreatment in certain cases. Moreover, dental team members should also be able to accept a patient from the bottom of heart as well as to understand and accept own emotions for a proper and successful treatment and for the avoidance of treatment failure. Therefore, dentists and other team members should become a mature, well-disposed and good person on behalf of the patients as well as on behalf of their own interest. Maintenance of the mental health of dental team is also crucial for the prevention of PDI. Lack of attention to all above aspects may invite treatment failure and may lead to the occurrence of PDI.

Screening of Risk Patients

Screening of risk patient is primarily based on the collection of a detailed patient's history as well as on careful evaluation of medical- psychosocial and orofacial status. Dissatisfaction with the existing denture may also indicate a risk of PDI. Patients with high dental fear and high fear of pain as well as with previous disturbances of patient-dentist relationship may also be considered as risk patients, because dental fear, pain and disturbances of patient-dentist relationship may lead to significant psychological stress that may turn into PDI symptoms. Accordingly, patients with high level of perceived stress may also be considered as risk patients. Besides semi-structured interview [Levin and Landesman 1976, Kaban and Belfer 1981] certain psychological and pathopsychological questionnaires may also be used for

screening purposes efficiently. Patients with increased dental fear could be recognized using the Dental Anxiety Scale (DAS) [Corah 1969], the Dental Fear Survey (DFS) [Kleinknecht et al. 1973, 1984] or the single-item Dental Anxiety Question (DAQ) [Neverlien 1990]. Risk patients in relation with dental fear may also be recognized using the "Expectation Scale" [Fábián G. et al. 2007]. Fear of pain may be measured using the Short form of Fear of Pain Questionnaire (FPQ-SF) measuring four factors such as fear of severe-, minor-, injection- and dental-pain [Asmundson et al. 2008]. Previous disturbances of the patient-dentist relationship may be recognized using the Getz's Dental Beliefs Survey (DBS) [Milgrom et al. 1985]. Exposure to stressful life events during the preceding period may be measured using the Social Readjustment Rating Scale (SRRS) [Holmes and Rahe 1967]. Similarly, patients' perception of stress may be measured using the Perceived Stress Scale (PSS) [Cohen et al. 1983]. Shortest form (Form C) of Cattell's 16 PF questionnaire may also be used to predict the appearance of PDI [Cattell et al. 1970, Reeve et al. 1984].

Proper Treatment Planning

Lack of attention to all facets of treatment planning may invite treatment failure; therefore, assessment of both mental (psychosocial) as well as physical (biomedical) conditions should be evaluated carefully [Bensing 2000, Green and Laskin 2000]. Accordingly, patient's subjective need as well as the normatively determined need for replacement of missing teeth should also be evaluated carefully before any prosthodontic treatment [Mazurat and Mazurat 2003/a,b, Graham et al. 2006]; and it should always be considered that the success of a prosthetic treatment strongly depends on the patients' perception of the value of such replacement [Mazurat and Mazurat 2003/a,b]. It should be also considered that, patients are rather focused on social meaning of the mouth than to physical (physiological) function of the teeth when defining subjective need for replacement of missing teeth [Graham et al. 2006]. Although occlusion of a complete dental arch is preferable normatively in most cases, many people are really satisfied with less than 28 natural ($\pm$ wisdom) teeth. In fact, there is a discrepancy between the normative need and perceived need even for complete denture services in edentulous. Therefore, prompt replacement of absent teeth without a subjective need of patients may lead to an overtreatment and discomfort [Mazurat and Mazurat 2003/a,b] especially when RPDs are used because of financial (or other) reasons [Mazurat and Mazurat 2003/a,b, Bae et al. 2006].

It should be also considered that aesthetics seems to be more important than other functions for a great majority of individuals [Elias and Sheiham 1998]; therefore there can be a lack of subjective need for replacement of missing posterior (particularly molar) teeth [Elias and Sheiham 1998, Mazurat and Mazurat 2003/a,b] especially if only a few teeth are absent. Although aesthetics seems to be more important than other functions for a great majority of individuals, expectations and wishes of patient related to chewing must also not be underestimated [Fábián and Fábián 2000, Mazurat and Mazurat 2003/a,b, Fábián et al. 2007/a]. Although chewing ability can be acceptable for patients functioning from second premolar to second premolar (i.e. when all molars missing) [Mazurat and Mazurat 2003/a,b]; more advanced loss of teeth may induce strong difficulties of chewing. In such cases

improvement of mastication may be a highly important wish of patients certainly [Mazurat and Mazurat 2003/a,b]. Patients' expectations related to speech may also influence satisfaction with dentures strongly. Similarly, numerous other expectations and wishes of patients related to the denture and/or dental treatment may also appear. Importantly, executable and acceptable wishes and expectations of patient should be fulfilled. If inexecutable and/or unacceptable wishes and expectations also appear, the clinician should carefully weigh the option of nontreatment [Levin and Landesman 1976, Stein 1983, Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010].

Besides premised subjective need and conscious wishes related to denture function, unconscious wishes of patients related to both dental treatment and denture (as well as to dentist) should also be considered [Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010]. In contrast to conscious wishes, understanding unconscious wishes (e.g. to look younger, to stop the appearance of aging, to be loved by the dentist etc.) may be more challenging but is similarly important for preventing the manifestation of PDI symptoms [Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010]. It may occur that, the patient's conscious and/or unconscious wishes related to the dental treatment and/or dentures and/or dentist may not harmonize with the reality and possibilities of a prosthetic treatment.

It is a matter of considerable significance that the clinician carefully weighs the option of nontreatment in such cases [Stein 1983, Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010].

It should be also considered that, fixed dentures are usually preferable to a removable one for most patient [Szentpétery et al. 2005]. Therefore properly planned fixed dentures supported either by natural abutment teeth or by properly inserted dental implants may decrease the risk of triggering psychogenic symptoms (comparing to the estimated risk of similar cases treated with RPDs).

However, financial means and compliance of the patient as well as the amount of stress and possible complications induced by preparation of teeth and/or implant-surgical interventions should also be considered very carefully. Because of the high success rate of endosseous dental implants, strategic extraction of compromised tooth/teeth and their replacement with implant supported fixed dentures should also be considered.

Importantly, planning of a time-schedule ensuring enough time to carry out a high quality, precise, "lege artis" dental treatment is one of the most important points to prevent triggering psychogenic symptoms. Timing of the start of prosthetic treatment and the insertion of dentures may also be of particular importance [Müller-Fahlbusch and Sone 1982], because heavy psychosocial and/or psychoemotional stress conditions as well as the active phases of psychiatric disorders significantly increase the risk of PDI [Müller-Fahlbusch and Sone 1982].

Therefore, starting of prosthetic treatment or insertion of dentures under any heavy psychological stress conditions and/or active phases of psychiatric disorders should be avoided [Müller-Fahlbusch and Sone 1982]; and the prosthetic treatment or insertion of denture should be delayed [Müller-Fahlbusch and Sone 1982].

Communication with the Patient

Communication with patient is a key factor of the prevention of PDI. There are three major facets of proper and efficient patient-dentist communication including (1) good interpersonal relationship and mutual trust, (2) exchange of information, and (3) making treatment related decisions [Fábián and Fejérdy 2010]. A good interpersonal relationship and mutual trust can be created due to friendliness, genuineness, readiness to help, empathy, respect, warmth, as well as eliciting feelings, paraphrasing and reflecting. Silence as well as humor may also be very useful in certain cases. Genuineness and congruence of any nonverbal communication is also highly important for creating a good interpersonal relationship and mutual trust. Simple basic techniques such as verbal following (to show he/she is listening to what the other say) [Deneen et al. 1973] and verbal reflection (to restate the essential feelings and contents of the message) [Deneen et al. 1973] as well as greeting the patient and using the patient's name should also not be forgotten. Exchange of information consists of information giving and information seeking, and to meet their needs, both dentists and patients alternate between information giving and information seeking. Patients need to know that what is the cause of their disorder as well as what are the treatment possibilities and possible treatment outcomes. Clear information about treatment procedures and their conditions should also be presented for patients; and patients' level of understanding should be raised until they clearly understand how the proposed treatment will meet their needs [Shigli et al. 2008]. Patients also need to know if the dentist accepts them and takes them seriously. Dentist needs information to establish the right diagnosis and treatment plan. For such purposes, listening to what the patient saying but also to what he/she is unable to say is crucial [Fábián and Fejérdy 2010]. Treatment related decisions-making should be based on mutuality, because not only the dentist but also the patient is responsible for his/her own health. The dentist should give up to hiding behind the facade of a professional authority and should share the problem and problem solving with the patient [Freeman 1999/a,b, Lechner and Roessler 2001]. However, if the dentist has unresolved issues regarding power and control, dependency and insecurity, self-esteem, prejudice, or other similar issues [Cohen 1994] such a "shared" decision-making could be rather untoward. Therefore, dentist's skills should include both an understanding of the patient as well as an understanding of oneself [Fábián and Fejérdy 2010].

Prevention of Dental Fear

Prevention (or at least a reduction) of dental fear should be integrated into dental treatment of every patients regularly, and especially into the treatment of patients with increased dental fear values [Kreyer 1989, Benson 2000, Hermes et al. 2006]. Especially because traumatizing dental events may trigger PDI symptoms [Müller-Fahlbusch and Sone 1982]. The most common reasons of dental fear are bad (painful or fearful) dental experiences [Moore and Birn 1990], lack of control over the social situation in the dental chair [Moore and Birn 1990], lack of control over personal emotional reactions [Moore and Birn 1990], feeling of powerlessness during treatment [Moore and Birn 1990] and social

learning processes with a negative image of dentists [Moore and Birn 1990]. The most important method for reduction of dental fear is communication. Great care should be taken to avoid fear producing terms or phrases describing the dental treatment or the dental instruments. It is also highly important that, to make only those promises that can be backed [Botto 2006]. It is also important that, to give the perception of being in control to the patient [Botto 2006]. For such purposes, the "tell-show-do" technique [Addelston 1959] may be used efficiently. This method includes describing in simple terms what is about to occur ("tell"); than allowing the patient to see, feel, explore and manipulate the tools or instruments ("show"); and than the starting of the procedure ("do") [Addelston 1959, Allen 2006]. Methods for distraction to refocus patient's attention away from the potentially painful/fearful stimulus or procedure may also be used, advantageously [Allen 2006, Botto 2006]. Offering of brief brakes during treatment is the simplest method for such purposes [Allen 2006]. Another simple forms of distraction include jiggling of the patients' cheek or asking them to hold their legs up in the air during a critical procedure (i.e. administrating of local anaesthesia, or making impression) [Botto 2006]. Music expressing positive emotions [Suda et al. 2008] may also be used for distraction during the dental treatment [Vinard and Ravier-rosenblum 1989, Botto 2006]. Audiovisual methods like video games [Allen 2006], two-dimensional DVD-glasses and three-dimensional virtual reality technology [Askay et al. 2009] may also be used. Mind-body therapies including hypnosis, self-hypnosis, photo-acoustic stimulation, relaxation and several biofeedback methods are also suitable for distraction during the dental treatment. In some cases pharmacological methods including premedication with anxiolytics relative analgesia (inhalation of nitrous oxide - oxygen mixture), conscious sedation (intravenously administered benzodiazepines) and occasionally general anaesthesia as well as psychotherapeutic approaches may also be used to reduce dental fear.

Prevention of Treatment Induced Pain

Painful dental treatment may also trigger PDI, especially when patient was emotionally traumatized [Müller-Fahlbusch and Sone 1982]. Therefore, pain catastrophizing behavior of patients (if any) should be recognized and considered. The use of highly efficient anaesthesia (including postoperative anaesthesia) and the use of psychological stress reducing interventions before treatment (i.e. cognitive or behavior therapy), during treatment (i.e. relaxation, hypnosis etc.) and after treatment (i.e. post hypnotic suggestions, self-hypnosis etc.) is also crucial for such patients. Despite local anaesthesia that have made painless dentistry a reality, psychological methods for alleviation of pain (and/or fear of pain) are still needed for sensitive patients [Benson 2000].

Patient-Nurse-Dentist Interrelationships

Psychological factors in patient-nurse-dentist interrelationships are major determinants of treatment outcome. Four major facets of patient-nurse-dentist relationships should be considered, such as the real relationship and the boundaries of relationship as well as the regression and the transference phenomena. The real relationship is an equal and unique

relationship between persons [Freeman 1999/b]. This is a genuine and realistic interaction in which the uniqueness of the dentist is complemented by the uniqueness of the patient and the uniqueness of the nurse [Freeman 1999/b]. The real relationship between the patient and the dentist (and/or nurse) should be protected by boundaries [Reid et al. 2007]. Acceptance of gifts, dinner invitations or unconventional payments for treatment (i.e. exchanging dental services for housepainting service, or for a discount on an automobile purchase etc.) put the dentist and/or the nurse at high risk of violating boundaries with patients [Reid et al. 2007]. Consequently, conflicts of interest may appear, which may erode their ability to make objective decisions on behalf of the patient [Reid et al. 2007]. Importantly, relation between the dentist and the nurse should also be protected by similar boundaries [Fábíán and Fejérdy 2010]. Regression simply describes the psychological state of the patients (and/or nurse and/or dentist) as they change from being in an emotionally controlled to a less well-controlled emotional state, which is associated with a change in relationship status [Freeman 1999/b]. Because of regression, the relationship may show a typical pattern of interaction between a parent and child [Freeman 1999/b] or any other pattern of emotionally deep-rooted relationships. Regression is frequently coupled with transference phenomena. Within the transference patients (and/or nurse, and/or dentist) will re-experience emotions and psycho-emotional patterns of previous relations (primarily from the childhood) [Freeman 1999/b]. The transference is particularly important because the dentist (as a "major target" of transference) may, therefore, be perceived emotionally as a "caring parent" or as a "powerful" or "threatening" person (or "any other" important person) [Freeman 1999/b, Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010]; which may lead to patient's (and/or nurse's) behavior appearing as inadequate and disturbing from the viewpoint of the dentist.

Prevention of Relapse

Prevention of relapse is primarily based on inducing and maintenance of long run changes of patients' health related behavior in respect of somatic (dental and medical) as well as mental health. It should be considered that, a structural plan for patient education has a better chance for success than one based only on intuition. Therefore, a clear identification and establishment of the educational needs of patients is crucial [Wentz 1972]. Subsequently, identification of possible/available motivation stimuli and concrete goal setting should be carried out [Wentz 1972]. Next goals of education plan should be achieved due to motivation and learning reinforcement [Wentz 1972]. Finally, results should be evaluated and controlled regularly [Wentz 1972]. Importantly, whatever structural plane is to be used, it must incorporate some means of providing information other than in an advice-giving format [Freeman 1999,a]. Information must be presented in such a way that patients feel its importance, and in a way that they can "take ownership" of it [Freeman 1999,a]. Depending on their personality, somatic condition, family/social background and symptoms, various methods may be offered for the patients. Most preferable home practice of several mind-body methods, diet- and medicinal herb therapy as well as CAM methods may be offered. Online support groups as well as patients' clubs could be another promising possibility for prevention

of relapse of PDI patients. Although prevention of relapse is primarily based on the patients' self-reliance, patient should be told to return for care if symptoms recur and cannot be quickly controlled by the previously used (home practice) procedures [Laskin and Block 1986].

Mental Health of Dental Team

It should be emphasized that, many dentists (and other members of the dental team) have a working situation with rather hard work conditions and occupational stress. Further, dentists' profession is coupled with a difficult psychoemotional challenge as well [Kreyer 1992]. The active inhibition of dentist's emotional expression (especially when dealing with problem patients) may also lead to increased working stress and risk for a variety of health problems [Berry and Pennebaker 1993]. To reduce psychoemotional stress during working with patients, dental team members should be able to accept a patient from the bottom of heart as well as to understand and accept own emotions. Thus, dentists and other team members should become a mature, well-disposed and good person on behalf of the patients as well as on behalf of their own interest. For such purposes several psychotherapeutic approaches as well as several mind-body methods are available and may be useful for dentists as well. Besides intrapersonal problems, interpersonal problems of team members should also be managed [Kreyer 1992]. In this respect, the most important is to recognize, verbalize and discuss the appearing problem. Verbalization and discussion can be facilitated due to organizing regular formal team meetings as well as informal team meetings.

Treatment Possibilities

Strategy of Psychosomatic Dental Therapy

Majority of psychogenic denture intolerance patients refuse to accept psychological background of their symptoms [Ross et al 1953, Schwichtenberg and Doering 2008], and instead of psychiatrists or psychotherapists, first they visit dentist and insist on the somatic origin of their symptoms [Ross et al. 1953]. Therefore, a simple referral to psychiatrist and/or psychotherapist would not solve the problem in most cases. Consequently, an initial psychosomatic therapy is needed prior to definitive therapy, which is a scope of dental profession's duty [Moulton et al. 1957, Pomp 1974]. The most important goals of initial psychosomatic therapy are avoidance of further useless invasive dental treatment [Brodine and Hartshorn 2004, Reewes and Merrill 2007, Baad-Hansen 2008] as well as obtaining decrease (recovery) of symptoms and motivation of patients to participate in a definitive psychosomatic therapy (which is the highest level care of psychogenic denture intolerance patients).

Gradual escalation of therapy and avoidance of irreversible forms of treatment are "cornerstones" of the initial psychosomatic therapy [Laskin and Block 1986]; which utilizes several placebo and/or palliative methods (i.e. physiotherapies, medication, medicinal herb therapy, diet therapy, complementary and alternative medicine /CAM/ therapy etc.) combined

with certain psychotherapeutic approaches and administration of any mind-body therapies (which are the "basic therapeutics" for psychosomatic disorders [Iversen 1989, Binder and Bider 1989, Krause 1983, 1994, 2000, Fábíán in press]). In contrast to initial therapy, definitive psychosomatic therapy should be carried out by specialized dental professionals as members of a specialized psychosomatic team including experienced dentists and other medical and psychotherapeutical professionals. Definitive psychosomatic therapy should be offered for patients being refractory to the initial psychosomatic therapy, and for patients responding to it but without a stable treatment outcome in a long run (i.e. patients with frequent relapses). Fundamentals of the complex process of dental psychosomatic therapy will be discussed in details below.

Initial Psychosomatic Therapy

As already indicated above, most of oral psychosomatic patients (especially PDI patients) refuse to accept psychological background of their symptoms [Ross et al 1953, Schwichtenberg and Doering 2008]. Instead of psychiatrists or psychotherapists, first they visit dentist and insist on the somatic origin of their symptoms [Ross et al. 1953, Fábíán et al. 2005/a]. In a dental psychosomatic unit, at least three-fourth (76,1 %) of the patients would surely need psychotherapy [Schwichtenberg and Doering 2008] but only 37,8 % of the patients would really begin a psychotherapeutic treatment [Schwichtenberg and Doering 2008]. Those, who do not begin definitive psychotherapy (the majority of such patients), are apt to continue doing "dentist shopping" (i.e. frequent and useless change of dentist) [Schwichtenberg and Doering 2008], with futile attempts at somatic therapies and/or surgical interventions [Kreyer 2000].

These data clearly indicate that, a simple referral to psychiatrist and/or psychotherapist would not solve the problem of the majority of PDI patients. Therefore, an initial psychosomatic therapy (IPT) is needed prior to definitive therapy, which is a scope of dental profession's duty [Moulton et al. 1957, Pomp 1974]. Initial psychosomatic therapy (IPT) is a scope of every dentist's duties currently, because of the absence of an extended network of specialized dental professionals. However, management of oral psychosomatic patients in specialized dental groups appears to be a meaningful perspective for the future [Dahlström et al. 1997, Green and Laskin 2000, Schwichtenberg and Doering 2008]. It may also be considered that, the network of prosthodontists may fulfill this task, but in this case, post-graduate education of prosthodontists should be completed with proper psychological and psychotherapeutical knowledge in a suitable manner.

In acute cases, the most important goal of initial psychosomatic therapy (IPT) is the avoidance of further useless invasive dental treatment [Brodine and Hartshorn 2004, Reewes and Merrill 2007, Baad-Hansen 2008], especially because aggravation or spread of symptoms following invasive dental interventions may occur frequently [Lesse 1956, Köling 1988, Toyofuku and Kikuta 2006]. The prevention of symptom chronification is also crucial, because chronification (especially with prolonged unsuccessful somatic/operative dental therapy in the history) frequently render pain and other symptoms intractable [Harris et al. 1993, Wöstmann 1996, Toyofuku and Kikuta 2006]. Prevention of symptom chronification

can be carried out efficiently with the simple avoidance of repeating unsuccessful dental treatments in many cases, in other cases symptom-centered palliative methods, certain psychotherapeutic approaches and any mind-body therapies should also be used.

Administration of premised symptom-centered palliative methods combined with certain psychotherapeutic approaches and with any mind-body therapies may lead to recovery in many cases, and solve the problem finally (definitively) [Pomp 1974, Harris et al. 1993, Wöstmann 1996, Fábián and Fejérdy 2010]. In other acute cases, there is some significant improvement, but there is no recovery of the symptoms during IPT. In such cases another important aim of initial psychosomatic therapy (IPT) is to keep these patients in a psychosomatic therapeutic relationship and motivate them to participate in a definitive psychosomatic therapy.

In chronic cases the prognosis of IPT is rather poor [Harris et al. 1993, Dahlström et al. 1997, Fábián and Fejérdy 2010]. There is no significant improvement during IPT in roughly half of chronic oral symptoms [Dahlström et al. 1997; Fábián et al. 2004, Schwichtenberg and Doering 2008]. Therefore, majority of chronic patients would need proper definitive psychosomatic therapy. Consequently, besides reaching some improvement of the symptoms (if any), major goal of IPT in chronic case is avoidance of repetition of unsuccessful futile somatic/operative dental treatment [Brodine and Hartshorn 2004] and obtaining motivation of patients to participate in a definitive psychosomatic therapy [Brodine and Hartshorn 2004, Demmel 2007].

As mentioned above, gradual escalation of therapy and avoidance of irreversible forms of treatment [Laskin and Block 1986] are "cornerstones" of the initial psychosomatic therapy. Therefore, the first goal of IPT in the management of psychogenic denture intolerance patients is to provide them with some understanding of their problem [Laskin and Block 1986], even if these patients often have difficulty accepting a psychophysiologic explanation for their disease [Laskin and Block 1986]. During this phase of IPT use of placebo and/or palliative methods (physiotherapies, medication, medicinal herb therapy, diet therapy, CAM therapy etc.) combined with certain psychotherapeutic approaches may be an appropriate way [Green and Laskin 1974, Laskin and Block 1986] to decrease the symptoms, and to develop a good patient-dentist relationship. At this phase of IPT some of the cases may be solved, others may be improved and at least an increase of patients' understanding related to their symptoms may be reached [Fábián and Fejérdy 2010].

Based on patients' improved understanding, second goal of IPT is to administer any mind-body therapies, which are "basic therapeutics" for psychosomatic disorders [Iversen 1989, Binder and Bider 1989, Krause 1994, Fábián et al. 2009/d]. Mind-body therapies (i.e. hypnosis, self-hypnosis, meditation, photo-acoustic stimulation, biofeedback methods etc.) advantageously improve psychogenic symptoms in most of the cases leading either to a recovery or to a significant improvement, coupled with increased motivation of patients to take part in a definitive psychosomatic therapy (see below) [Fábián and Fábián 1998; Fábián et al. 2002, 2005/a, 2009/c,d]. Accordingly, third goal of IPT is to refer patients being refractory to IPT to definitive psychosomatic therapy (see below) and to make sure that these patients enter and participate in the definitive psychosomatic therapy [Fábián and Fejérdy 2010, Demmel 2007].

Although the efficiency of initial psychosomatic therapy (IPT) is strongly depends on the improvement of symptoms; using efficient symptom-centered treatments it should be always

considered that, any psychogenic symptom may represent a defense against psychiatric decompensation [Lesse 1956, Delaney 1976, Marbach 1978, Violon 1980, Kaban and Belfer 1981]. Therefore, symptom-centered interventions should be administered carefully, resulting in slow-going, gradual decrease of the symptoms to prevent decompensation of the patient during IPT.

Definitive Psychosomatic Therapy

Definitive psychosomatic therapy (DPT) is a highest level care of PDI patients utilizing any available dental, medical and psychotherapeutical treatment possibilities in an evidence based manner. Definitive psychosomatic therapy should be offered for patients being refractory to IPT [Laskin and Block 1986], and for patients responding to IPT but without a stable treatment outcome in a long run (i.e. patients with frequent relapses). Definitive psychosomatic therapy (DPT) is clearly not a scope of every dentist's duties. Definitive psychosomatic therapy should be carried out by specialized dental professionals [Dahlström et al. 1997, Hertrich and Joraschky 1996, Dworkin et al. 2002, Fábíán and Fejérdy 2010] as members of a specialized psychosomatic team including experienced dentists and other medical and psychotherapeutical professionals.

Although clinical experiences clearly indicate that there is a major role of specialized dental professionals also in definitive psychosomatic therapy (DPT) in most cases; close collaboration with medical professionals (especially with psychiatrists and neurologists) and with psychotherapists is clearly needed for DPT [Dahlström et al. 1997, Hertrich and Joraschky 1996, Von Korff 2005, Turner et al. 2006, Schwichtenberg and Doering 2008]. Dentists having full-qualification in psychotherapy and/or psychosomatic dentistry seem to be the most adequate professionals to organize definitive psychosomatic therapy for PDI patients, especially because continuation of a psychosomatic supportive dental care is needed in most of the cases [Laskin and Block 1986, Fábíán and Fejérdy 2010].

It should be also considered that, PDI patients usually display a lower level of psychopathology comparing to psychiatric patients [Meldolesi et al. 2000]; but in some cases severe psychopathologies including psychoses may also appear as a background of PDI symptoms [Lesse 1956, Delaney 1976, Marbach 1978, Violon 1980, Kaban and Belfer 1981, Fábíán 1999]. In such cases referral to psychiatrist and definitive psychiatric therapy (instead of psychosomatic therapy) is crucial; although a supportive type psychosomatic dental care is usually also needed for proper definitive treatment and for stabilizing oral functions and oral health.

After-Care

All patients including PDI patients who wear any fixed or removable dentures should have annual examinations [Woelfel 1983]. These examinations should include evaluation of the health of oral/orofacial tissues (with special respect of the supporting teeth and other supporting tissues) as well as evaluation of the condition, retention, stability and other

relevant properties of dentures [Woelfel 1983]. PDI related psychogenic symptoms of patients should also be examined.

Although there are rather few studies about the efficiency of treatment of oral psychosomatic patients, it is likely that at least 50-60% of such patients significantly improve in their complaints [Schwichtenberg and Doering 2008]. Sometimes, especially in not yet chronic cases the success rate may be even higher. However, there is still a relatively large amount of patients without any success or with suffering from several residual symptoms.

Therefore, patients should be made aware that it may not be possible to provide a permanent cure for their problem, but that they can learn to manage in a satisfactory manner [Laskin and Block 1986]. Home practice of most mind-body therapies, diet- and medicinal herb therapy as well as complementary and alternative medicine (CAM) therapies regularly supervised by the dentist (and/or other professionals) are good tools to manage such residual symptoms as well as to maintain clinical results and to prevent relapse [Laskin and Block 1986, Fábíán and Fejérdy 2010]. In the case of relapse (or exacerbation) of the PDI symptoms, patients should be treated as described in respect of initial psychosomatic therapy (IPT).

Conclusion

Even though a prosthesis is fabricated conscientiously and properly, there is no assurance that the patient will be comfortable while wearing it or satisfied with the therapy [Mazurat and Mazurat 2003/a,b]. A normative evaluation by a dentist and a subjective evaluation by the patient related to the denture or to the dental treatment may be rather different [Lechner and Roessler 2001]. The factors not related to operative/technological dental skills that contribute to the success of prosthodontic treatments are becoming more and more important.

PDI related symptoms are usually multifactorial (multicausal) notwithstanding that there is always a major psychogenic component behind the symptoms too [Fábíán and Fábíán 2000, Fábíán et al. 2006, 2007/a]. Therefore, both somatic and psychogenic pathways of pathomechanism should be considered carefully for both diagnosis and treatment. Since psychogenic symptoms may mimic a great variety of somatic symptoms; a clear cut diagnosis and proper differential diagnosis of PDI could be rather difficult. Therefore it should be emphasized that, the diagnosis of PDI is a presumptive one in many cases. The diagnosis may change later in the course of the disease as the clinical findings change and/or stabilize, therefore continuous monitoring and evaluation of the patient over time is essential [Sarlaní et al. 2005/a,b]. A detailed differential diagnosis considering possible disorders of the teeth, and/or oral- and maxillofacial tissues should also be carried out, and any other possible somatic causes behind the symptoms should also be excluded.

Prevention is likely to be the most important tool for the management of psychogenic denture intolerance related problems at a social level. Cornerstones of prevention are screening of risk patients, proper treatment planning, proper communication with the patient, screening of the patient-nurse-dentist interrelationships, as well as prevention of dental fear, prevention of treatment induced pain and prevention of relapse [Fábíán and Fábíán 2000, Fábíán et al. 2007/a, Fábíán and Fejérdy 2010]. It is also crucial for the prevention that,

patients must be made aware of their responsibilities in achieving a satisfactory outcome. It is also a matter of considerable significance that the clinician carefully weighs the option of nontreatment in certain cases [Levin and Landesman 1976, Stein 1983, Fábíán and Fábíán 2000, Fábíán et al. 2007/a].

Majority of psychogenic denture intolerance patients refuse to accept psychological background of their symptoms [Ross et al 1953, Fábíán et al. 2005/a, Schwichtenberg and Doering 2008]; and instead of psychiatrists or psychotherapists, first they visit dentist and insist on the somatic origin of their symptoms [Ross et al. 1953, Fábíán et al. 2005/a]. Therefore a simple referral to psychiatrist and/or psychotherapist would not solve the problem in most cases. Consequently an initial psychosomatic therapy is needed prior to definitive therapy, which is a scope of dental profession's duty [Moulton et al. 1957, Pomp 1974, Fábíán and Fejérdy 2010]. Avoidance of irreversible forms of treatment is crucial for a proper initial psychosomatic therapy certainly. Initial psychosomatic therapy may utilize several placebo and/or palliative methods (i.e. physiotherapies, medication, medicinal herb therapy, diet therapy, CAM therapy etc.) combined with certain psychotherapeutic approaches and administration of any mind-body therapies, which are "basis therapeutics" for psychosomatic disorders.

References

- Adam, RZ; Geerts, GA; Lalloo, R. The impact of new complete dentures on oral health-related quality of life. *SADJ*, 2007 62, 264-268.
- Addelston, H. Child patient training. *CDS Review*, 1959 38, 27-29.
- Allen, PF. and McMillan, AS. A review of the functional and psychosocial outcomes of edentulousness treated with complete replacement dentures. *J. Can. Dent. Assoc.*, 2003 69, 662.
- Ament, P. and Ament, A. Body image in dentistry. *J. Prosthet Dent*, 1970 24, 362-366.
- Aneksuk, V. Burning mouth syndrome. *J. Dent. Assoc. Thai*, 1989 39, 251-258.
- Askay, SW; Patterson, DR; Sharar, SR. Virtual reality hypnosis. *Contemp Hypnosis*, 2009 26, 40-47.
- Asmundson, GJG; Bovell, CV; Carleton, RN; McWilliams, LA. The Fear of Pain Questionnaire - Short Form (FPQ-SF): Factor validity and psychometric properties. *Pain*, 2008 134, 51-58.
- Baad-Hansen, L. Atypical odontalgia - pathophysiology and clinical management. *J. Oral. Rehabil.*, 2008 35, 1-11.
- Baba, K; Aridome, K; Haketa, T; Kino, K; Ohyama, T. Sensory perceptive and discriminative abilities of patients with occlusal dysesthesia. *Nihon Hotesu Shika Gakki Zasshi*, 2005 49, 599-607.
- Bae, KH; Kim, C; Paik, DI; Kim, JB. A comparison of oral health related quality of life between complete and partial removable denture-wearing older adults in Korea. *J. Oral. Rehabil.* 2006 33, 317-322.
- Baldry, PE. *Acupuncture, trigger points and musculoskeletal pain*. (Third edition). Edinburgh - London - New York - Philadelphia - San Francisco - Toronto: Elsevier; 2005; 45-72.

- Balters, W. Die Bedeutung von Zahnverlust und Zahnersatz für den Patienten - von der Psychologie her gesehen. *Dtsch Zahnärztl Z*, 1956/a 11, 112-120.
- Balters, W. Die Bedeutung von Zahnverlust und Zahnersatz für den Patienten - von der Psychologie her gesehen. (Fortsetzung). *Dtsch Zahnärztl Z*, 1956/b 11, 465-468.
- Barsby, MJ. The use of hypnosis in the management of "gagging" and intolerance to dentures. *Br Dent J*, 1994 176, 97-102.
- Bensing, J. Bridging the gap. The separate worlds of evidence-based medicine and patient-centered medicine. *Patient Education Counseling*, 2000 39, 17-25.
- Benson, PE. Suggestion can help. *Ann R Australas Coll Dent Surg*, 2000 15, 284-285.
- Bergendal, B. The relative importance of tooth loss and denture wearing in Swedish adults. *Community Dent Health*, 1989 6, 103-111.
- Berry, DS. and Pennebaker, JW. Nonverbal and verbal emotional expressions and health. *Psychother Psychosom*, 1993 59, 11-19.
- Besson, JM. The neurobiology of pain. *Lancet*, 1999 353, 1610-1615.
- Binder, H. and Binder, K. *Autogenes Training, Basispsychotherapeutikum*. Köln: Deutscher Ärzteverlag; 1989.
- Biondi, M. and Picardi, A. Temporomandibular joint pain-dysfunction syndrome and bruxism: etiopathogenesis and treatment from a psychosomatic integrative viewpoint. *Psychother Psychosom*, 1993 59, 84-98.
- Borelli, S; Seifert, HU; Seifert, B. Sensibilisierung und allergische Reaktionen. In: hupfauf L (ed.) *Teilprothesen*. München: Urban and Schwarzenberg; 1988; 247-263.
- Botto, RW. Chairside techniques for reducing dental fear. In: Mostofsky DI, Forgione AG, Giddon DB editors. *Behavioral dentistry*. Ames (Iowa): Blackwell Munksgaard; 2006; 115-125.
- Böning, J. Psychosomatic and psychopathological aspects in dental-orofacial medicine with special reference to old age. *Z Gerontol*, 1990 23, 318-321.
- Brodine, AL. and Hartshorn, MA. Recognition and management of somatoform disorders. *J. Prosthet Dent*, 2004 91, 268-273.
- Burton, RC. The problem of facial pain. *JADA*, 1969 79, 93-101.
- Bushfield, BL; Wechsler, H; Barnum, WJ. Studies of salivation in depression. *Arch. Gen. Psych.*, 1961 5, 472-477.
- Büdinger, L. and Hertl, M. Immunologic mechanisms in hypersensitivity reactions to metal ions: an overview. *Allergy*, 2000 55, 108-115.
- Carli, G. An update on pain physiology: the relevance of Craig's and Jäning's hypotheses for hypnotic analgesia. *Contemp Hypnosis*, 2009 26, 4-14.
- Cattell, RB; Eber, HW; Tatsuoka, M. *Handbook for the 16 PF Questionnaire*. Champaign, Illinois: Institute of Personality and Ability Testing; 1970.
- Chang, FCT. and Scott, TR. Conditioned taste aversions modify neuronal responses to gustatory stimuli. *J. Neurosci.*, 1984 4, 1850-1862.
- Chrousos, GP. Stressors, stress, and neuroendocrine integration of the adaptive response. The 1997 Hans Selye memorial lecture. *Ann. NY Acad. Sci.*, 1998 851, 311-335.
- Clark, DL; Boutros, NN; Mendez, MF. *The brain and behavior. An introduction to behavioral neuroanatomy*. Cambridge: Cambridge University Press; 2006; 165-177.
- Clark, GT; Minakuchi, H; Lotaif, AC. Orofacial pain and sensory disorders in the elderly. *Dent Clin North Am*, 2005 49, 343-362.
- Cohen, PA. Recognizing and managing the difficult patient. *CDS Review*, 1994, 11-16.

- Cohen, S; Kamarck, T; Mermelstein, R. A global measure of perceived stress. *J Health Soc Behav*, 1983 24, 385-396.
- Corah, NL. Development of a Dental Anxiety Scale. *J. Dent Res.*, 1969 48, 596.
- Dahlsröm, L. Electromyographic studies of craniomandibular disorders: a review of the literature. *J. Oral. Rehabil.* 1989 16, 1-20.
- Dahlström, L; Lindvall, AM; Milthon, R; Widmark, G. Management of chronic orofacial pain: attitudes among patients and dentists in a Swedish county. *Acta Odontol. Scand.*, 1997 55, 181-185.
- De Rossi, SS. and Greenberg, MS. Intraoral contact allergy: A literature review and case reports. *JADA*, 1998 129, 1435-1441.
- Delaney, JF. Atypical facial pain as a defense against psychosis. *Am. J. Psychiat.*, 1976 133, 1151-1154.
- Demmel, H-J. and Lamprecht, F. Zahnheilkunde. In: von Uexküll T, editor. *Psychosomatische Medizin*. München, Wien, Baltimore: Urban and Schwarzenberg; 1996; 1125-1130.
- Demmel, HJ. Die Überweisung psychosomatisch kranker Patienten. *Zahnärztl Mitteilungen*, 2007 97, 40-44.
- Deneen, LJ; Heid, DW; Smith, AA. Effective interpersonal and management skills in dentistry. *JADA*, 1973 87, 878-880.
- Di Felice, R; Samson, J; Carlino, P; Giuliani, M; Fiore-Donno, G. Psychogenic oral paraesthesia. *Rev. Odontostomatol. (Paris)*, 1991 20, 189-194.
- Doering, S. and Wolowski, A. Psychosomatik als interdisziplinäres Fach. *Zahnärztl Nachr Sachsen-Anhalt*, 2007 12, 7.
- Dolder, E. Zur Psychologie des Zahn-Verlustes und des Zahn-Ersatzes. *Dtsch Zahnärztl Z*, 1956 11, 469-477.
- Drost, R. Sogenannte prothesenunverträglichkeit (oder: Die nichtacceptierte Prothese). Psychosomatische Aspekte - I. Teil -. *ZWR*, 1978/a 87, 848-852.
- Drost, R. Sogenannte prothesenunverträglichkeit (oder: Die nichtacceptierte Prothese). Psychosomatische Aspekte - II. Teil -. *ZWR*, 1978/b 87, 907-913.
- Dworkin, SF. and Sherman J. Chronic orofacial pain: biobehavioral perspectives. In: Mostofsky DI, Forgione AG, Giddon, DB editors. *Behavioral dentistry*. Ames (Iowa): Blackwell Munksgaard; 2006; 99-113.
- Dworkin, SF; Turner, JA; Mancl, L; Wilson, L; Massoth, D; Huggins, KH; LeResche, L; Truelove, E. A randomized clinical trial of a tailored comprehensive care treatment program for temporomandibular disorders. *J. Orofac. Pain*, 2002 16, 259-276.
- Eli, I. and Kleinhauz, M. Hypnosis and dentistry. *Hypn Int Monographs*, 1997 3, 59-69.
- Elias, AC. and Sheiham, A. The relationship between satisfaction with mouth and number and position of teeth. *J. Oral. Rehabil.*, 1998 25, 649-661.
- Ellis, JS; Pelekis, ND; Thomason, JM. Conventional rehabilitation of edentulous patients: the impact on oral health-related quality of life and patient satisfaction. *J. Prosthodont*, 2007 16, 37-42.
- Emerson, WA. and Giddon, DR. Psychologic factors in adjustment to full denture prosthesis. *J. Dent Res.*, 1955 34, 683-684.
- Engel, G. The need for a new medical model: a challenge for biomedicine. *Science*, 1977 196, 129-136.

- Fábián, G; Müller, O; Kovács, Sz; Nguyen, MT; Fábián, TK; Csermely, P; Fejérdy, P. Attitude toward death: Does it influence dental fear? *Ann. NY Acad. Sci.*, 2007 1113, 339-349.
- Fábián, TK. Treatment possibilities of denture intolerance. Use of photo-acoustic stimulation and hypnotherapy in a difficult clinical case. [Adalékok a protézis intolerancia gyógyításához. Fény-hang kezeléssel kombinált hipnoterápia alkalmazásának szokatlan módja egy eset kapcsán.] *Hipno Info*, 1999 38, 81-88.
- Fábián, TK. Mental health and pastoral counseling [Mentálhigiéné és lelkipogondozás]. In: Fábián TK, Vértes G. editors *Psychosomatic dentistry [Fogorvosi pszichoszomatika]*. Budapest: Medicina; 2007/a; 158-168.
- Fábián, TK. Psychosomatics of salivation problems [A nyálszekréciós zavarok pszichoszomatikája]. In: Fábián TK, Vértes G editors. *Psychosomatic dentistry [Fogorvosi pszichoszomatika]*. Budapest: Medicina; 2007/b; 121-126.
- Fábián, TK. *Mind-body connections. Pathways of psychosomatic coupling under meditation and other altered states of consciousness*. New York: Nova Science Publishers; in press.
- Fábián, TK. and Fábián, G. Stress of life, stress of death: Anxiety in dentistry from the viewpoint of hypnotherapy. *Ann. NY Acad. Sci.*, 1998 851, 495-500.
- Fábián, TK. and Fábián, G. Dental stress. In: Fink G, editor in chef. *Encyclopedia of Stress*. Vol. 1. San Diego: Academic Press; 2000; 657-659.
- Fábián, TK. and Fejérdy, P. Denture intolerance [Fogpótlás intolerancia]. In: Fábián TK, Vértes G editors. *Psychosomatic dentistry [Fogorvosi pszichoszomatika]*. Budapest: Medicina; 2007; 127-146.
- Fábián, TK and Fejérdy P. *Psychogenic denture intolerance. Theoretical background, prevention and treatment possibilities*. New York: Nova Science Publishers; 2010.
- Fábián, TK; Vértes, G; Tóth, Zs. Some depth-psychological aspects of the psychogenic symptoms of the orofacial tissues. *Fogorv Szle*, 2000 93, 262-267.
- Fábián, TK; Vértes, G; Szabó, A; Varga, K. Photo-acoustic stimulation and hypnotherapy. An effective combination for treatment of oral psychosomatic disorders. *Hypn Int. Monographs*, 2002 6, 199-207.
- Fábián, TK; Kaán, B; Fejérdy, L; Tóth, Zs; Fejérdy, P. Effectiveness of psychotherapy in the treatment of denture intolerance. Evaluation of 25 cases. *Fogorv Szle*, 2004 97, 163-168.
- Fábián, TK; Krause, WR; Krause, M; Fejérdy, P. Photo-acoustic stimulation and hypnotherapy in the treatment of oral psychosomatic disorders. *Hypnos*, 2005/a 32, 198-202.
- Fábián, TK; Vértes, G; Fejérdy, P. Pastoral psychology, spiritual counseling in dentistry. Review of the literature. *Fogorv Szle*, 2005/b 98, 37-42.
- Fábián, TK; Mierzwińska-Nastalska, E; Fejérdy P. Photo-acoustic stimulation. A suitable method in the treatment of psychogenic denture intolerance. *Protet Stomatol*, 2006/a 56, 335-340.
- Fábián, TK; Kovács, Sz; Müller, O; Fábián, G; Marten, A; Fejérdy, P. Some aspects of existential psychotherapy in dentistry. *Fogorv Szle*, 2006/b 99, 246.
- Fábián, TK; Fábián, G; Fejérdy, P. Dental Stress. In: Fink G, editor in chef. *Encyclopedia of Stress. 2-nd enlarged edition*, Vol. 1. Oxford: Academic Press; 2007/a; 733-736.
- Fábián, TK; Fejérdy, P; Nguyen, MT; Söti, Cs; Csermely, P. Potential immunological functions of salivary Hsp70 in mucosal and periodontal defense mechanisms. *Arch Immunol Ther. Exp.*, 2007/b 55, 91-98.

- Fábíán, TK; Fejérdy, P; Csermely, P. Chemical biology of saliva in health and disease. In: Begley T, editor in chief. *Wiley encyclopedia of chemical biology*. Hoboken: John Wiley and sons; 2008/a. DOI: 10.1002/9780470048672.webc643.
- Fábíán, TK; Söti, Cs; Nguyen, MT; Csermely, P; Fejérdy, P. Expected functions of salivary HSP70 in the oral cavity. (Review article). In: Morell E, Vincent C editors. *Heat shock proteins: New research*. Hauppauge: Nova Science Publishers; 2008/b; 321-340.
- Fábíán, TK; Söti, Cs; Nguyen, MT; Csermely, P; Fejérdy, P. Expected functions of salivary HSP70 in the oral cavity. *Int. J. Med. Biol. Frontier*, 2008/c; 289-308.
- Fábíán, TK; Fejérdy, P; Csermely, P. Salivary genomics, transcriptomics and proteomics: The emerging concept of the oral ecosystem and their use in the early diagnosis of cancer and other diseases. *Current Genomics*, 2008/d 9, 11-21.
- Fábíán, TK; Gótai, L; Beck, A; Fábíán, G; Fejérdy, P. The role of molecular chaperones (HSPAs/HSP70s) in oral health and oral inflammatory diseases: A review. *Eur. J. Inflammation*, 2009/a 7, 53-61.
- Fábíán, TK; Csermely, P; Fábíán, G; Fejérdy, P. Spondyloarthropathies and bone resorption: A possible role of heat shock protein (Hsp70). (Review). *Acta Physiol Hungarica*, 2009/b 96, 149-155.
- Fábíán, TK; Gótai, L; Krause, WR; Fejérdy, P. Zahnärztliche Hypnoseforschung an der Semmelweis Universität Budapest. *Deutsche Z Zahnärztl Hypn*, 2009/c 8, 9-14.
- Fábíán, TK; Kovács, KJ; Gótai, L; Beck, A; Krause, WR; Fejérdy, P. Photo-acoustic stimulation: Theoretical background and ten years of clinical experience. *Contemp Hypn*, 2009/d 26, 225-233.
- Fassbind, O. Psychogene Prothesenunverträglichkeit: Therapiemöglichkeiten. *Schweiz Mschr Zahnmed*, 1985 95, 595-598.
- Fejérdy, P; Fábíán, TK; Krause, W-R. Mentalhygienische Aufgaben von Krankenschwestern in der Zahnmedizin. Kommunikation mit den Patienten und dem Zahnarzt. *Dtsch Z Zahnärztl Hypn*. 2004 4; 32-34.
- Figgenger, L. Non-acceptance of prosthetic appliances at the focus of forensic consequences. *J. Forensic Odontostomatol.*, 1996 14, 28-29.
- Fiske, J; Davis, DM; Frances, C; Gelbier, S. The emotional effects of tooth loss in edentulous people. *Brit. Dent J.*, 1998 184, 90-93.
- Freeman, R. Strategies for motivating the non-compliant patient. *Brit. Dent J.* 1999/a 187, 307-312.
- Freeman, R. A psychodynamic understanding of the dentist-patient interaction. *Brit. Dent J.*, 1999/b 186, 503-505.
- Fuhr, K. and Reiber, T. Klinische Funktionsdiagnostik. In: Koeck B, editor. *Funktionsstörungen des Kauorgans*. München - Wien - Baltimore: Urban and Schwarzenberg; 1995; 75-78.
- Gale, EN. and Carlsson, SG. Look carefully; a short note on symptom substitution. *Behav. Res. Ther.*, 1976 14, 77.
- Garhammer, P; Schmalz, G; Hiller, KA; Reitingner, T; Stolz, W. Patients with local adverse effects from dental alloys: frequency, complaints, symptoms, allergy. *Clin. Oral Invest*, 2001 5, 240-249.
- Gáspár, J; Fejérdy, L; Fábíán, TK. The psychical aspects of overactive gag reflex (gagging): a case report. *Fogorv Szle*, 2002 95, 199-203.

- Gerbershagen, HU. *Die schwierige Schmerzpatient in der Zahnmedizin*. Stuttgart: Thieme; 1995; 2-15, 27, 39, 116-117.
- Gerbert, B; Bernzweig, J; Bleecker, T; Bader, J; Miyasaki, C. How dentists see themselves, their profession, the public. *JADA*, 1992 123, 72-78.
- Giddon, DB. and Anderson, NK. The oral and craniofacial area and interpersonal attraction. In: Mostofsky DI, Forgione AG, Giddon DB editors. *Behavioral dentistry*. Ames (Iowa): Blackwell Munksgaard; 2006; 3-17.
- Giddon, DB. and Hittelman, E. Psychologic aspects of prosthodontic treatment for geriatric patients. *J. Prosthet Dent*, 1980 43, 374-379.
- Golebiewska, M; Sierpińska, T; Namiot, D; Likeman, PR. Affective state and acceptance of dentures in elderly patients. *Gerodontology*, 1998 15, 87-92.
- Gonzalez, JB. and Desjardins, RP. Management of patients with new prostheses. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment in prosthodontics*. Philadelphia: Lea and Febiger; 1983; 506-535.
- Gottlieb, G; Paulson, G; Raleigh, NC. Salivation in depressed patients. *Arch. Gen. Psych.*, 1961 5, 468-471.
- Grabauskas, G. and Bradley, R. Ionic mechanisms of GABAA biphasic synaptic potentials in gustatory nucleus of the solitary tract. *Ann. NY Acad. Sci.*, 1998 855, 486-487.
- Graber, G. Neurologische und psychosomatische Aspekte der Myoarthropathien des Kauorgans. *Zahnärztl Welt Rundschau*, 1971 21, 997-1000.
- Graham, G. Two case-histories of ladies who were unable to wear their dentures without feeling sick and retching. *Hypnos*, 2005 32, 210-212.
- Graham, R; Mihaylov, S; Jepson, N; Allen, PF; Bond, S. Determining "need" for a removable partial denture: a qualitative study of factors that influence dentist provision and patient use. *Brit. Dent J.*, 2006 200, 155-158.
- Green, CS. and Laskin, DM. Long-term evaluation of conservative treatment for myofascial pain-dysfunction syndrome. *JADA*, 1974 89, 1365-1368.
- Green, CS. and Laskin, DM. Temporomandibular disorders: Moving from a dentally based to a medically based model. *J. Dent Res.*, 2000 79, 1736-1739.
- Griffin, CJ. and Harris, R. The regulatory influences of the trigeminal system. In: Griffin CJ, Harris R editors. *The temporomandibular joint syndrome. The masticatory apparatus of man in normal and abnormal function*. (Monographs in oral science, Vol. 4.) Basel - München - Paris - London - New York - Sydney: Karger; 1975; 65-86.
- Grushka, M. and Sessle, BJ. Burning mouth syndrome. *Dent Clin. North Am.*, 1991 35, 171-184.
- Guckes, AD; Smith, DE; Swoope, CC. Counseling and related factors influencing satisfaction with dentures. *J. Prosthet Dent*, 1978 39, 259-267.
- Guggenheimer, J. and Moore, PA. Xerostomia. Etiology, recognition and treatment. *JADA*, 2003 134, 61-69.
- Hach, B; Lehl, S; Niedermeier, W. Psychopathological and psychopathometric findings in patients with dental prosthesis intolerance. *Dtsch Zahnärztl Z*, 1978 33, 238-244.
- Hampf, G. Dilemma in treatment of patients suffering from orofacial dysaesthesia. *Int. J. Oral. Maxillofac. Surg.*, 1987 16, 397-401.
- Hansen, K. and Schliack, H. *Segmentale Innervation*. Stuttgart: Thieme, 1962.
- Harris, M; Feinmann, C; Wise, M; Treasure, F. Temporomandibular joint and orofacial pain: clinical and medicolegal management problems. *Brit. Dent J.*, 1993 174, 129-136.

- Harris, R. and Griffin, CJ. Neuromuscular mechanisms and the masticatory apparatus. In: Griffin CJ, Harris R editors. *The temporomandibular joint syndrome. The masticatory apparatus of man in normal and abnormal function.* (Monographs in oral science, Vol. 4.) Basel - München - Paris - London - New York - Sydney: Karger; 1975; 45-64.
- Har-Zion, G; Brin, I; Steiner, J. Psychophysiological testing of taste and flavour reactivity in young patients undergoing treatment with removable orthodontic appliances. *Eur. J. Orthodontics*, 2004 26, 73-78.
- Hassel, AJ; Wegener, I; Rolko, C; Nitschke, I. Self-rating of satisfaction with dental appearance in an elderly German population. *Int. Dent J.*, 2008 58, 98-102.
- Heggendorn, H; Voght, HP; Graber, G. Experimentelle Untersuchungen über die orale Hyperaktivität bei psychischer Belastung, im besonderen bei Aggression. *Schweiz Mschr Zahnheilk*, 1979 89, 1148-1161.
- Hermes, D; Saka, B; Bahlmann, L; Matthes, M. Behandlungsangst in der Mund-, Kiefer- und Gesichtschirurgie. *Mund Kiefer Gesichts Chir*, 2006 10, 307-313.
- Hertrich, K. and Joraschky, P. Psychotherapie in der zahnärztlichen Praxis: Möglichkeiten der Integration und Kooperation. In: Sergl HG, editor. *Psychologie und Psychosomatik in der Zahnheilkunde.* München - Wien - Baltimore: Urban and Schwarzenberg; 1996; 261-279.
- Hirsch, B; Levin, B; Tiber, N. Effects of patient involvement and esthetic preference on denture acceptance. *J. Prosthet Dent*, 1972 28, 127-132.
- Hirsch, B; Levin, B; Tiber, N. Effects of dentist authoritarianism on patient evaluation of dentures. *J. Prosthet Dent*, 1973 30, 745-748.
- Holland-Moritz, R. Systematische Befunderhebung bei Zahnersatzunverträglichkeit. *Dtsch Zahnärztl Z*, 1979 34, 786-788.
- Holland-Moritz, R. Zahnersatzunverträglichkeit und Schleimhautbrennen. *Dtsch Zahnärztl Z*. 1980 35, 948-952.
- Holmes, TH. and Rahe, RH. The Social Readjustment Rating Scale. *J. Psychosom. Res.*, 1967 11, 213.
- Hunt, CC. and Paintal, AS. Spinal reflex regulation of fusimotor neurons. *J. Physiol. Lond*, 1958 143, 195-212.
- Ismail, Y; Zullo, T; Kruper, D. The use of the psychological screening inventory as a predictor of problem denture patients. *J. Dent Res.*, 1974 54 (spec. issue. A), L128.
- Iversen, G. and Geleitwort. In: Binder H, Binder K editors. *Autogenes Training, Basispsychotherapeutikum.* Köln: Deutscher Ärzteverlag; 1989.
- Jagger, RG; Korszun, A. Phantom bite revised. *Br. Dent J.*, 2004 197, 241-243.
- Kaán, M; Bolla, K; Keszler, B. Speech characteristics of persons wearing full upper and lower prostheses. *Fogorv Szle*, 1993 86, 45-53.
- Kaban, LB. and Belfer, ML. Temporomandibular joint dysfunction: an occasional manifestation of serious psychopathology. *J. Oral. Surg.*, 1981 39, 742-746.
- Kampe, T; Edman, G; Bader, G; Tagdae, T; Karlsson, S. Personality traits in a group of subjects with long-standing bruxing behaviour. *J. Oral. Rehabil.*, 1997 24, 588-593.
- Király, M; Porcsalmy, B; Pataki, A; Kádár, K; Jelítai, M; Molnár, B; Hermann, P; Gera, I; Grimm, WD; Ganss, B; Zsembéry, A; Varga, G. Simultaneous PKC and cAMP activation induces differentiation of human dental pulp stem cells into functionally active neurons. *Neurochem. Int.*, 2009 55, 323-332.

- Kivovics, P; Sajgó, P; Herendi, G; Hermann, P. Evaluation of comfort, sound formation and food relations of the oral surface of mould cast metal plates, based on patient questionnaires. *Fogorv Szle*, 1990 83, 371-373.
- Kiyak, HA. Measuring psychosocial variables that predict older persons' oral health behavior. *Gerodontology*, 1996 13, 69-75.
- Kleinknecht, RA; Klepac, RK; Alexander, LD. Origins and characteristics of fear of dentistry. *J. Am. Dent. Assoc.*, 1973 86, 842-848.
- Kleinknecht, RA; Thorndike, RM; McGlynn, FD; Harkavy, J. Factor analysis of the Dental Fear Survey with cross-validation. *J. Am. Dent. Assoc.*, 1984 108, 59-61.
- Kotkin, H. Diagnostic significance of denture complaints. *J. Prosthet Dent.*, 1985 53, 73-77.
- Köling, A. Neurologist, otolaryngologist...? Which specialist should treat facial pain? *Lakartidningen*, 1998 95, 2320-2325.
- Kranz, H. Zahnverlust und Zahnersatz als psychologisches Problem. *Dtsch Zahnärztl Z*, 1956 11, 105-112.
- Krause, WR. Grundlagen der Biofeedback-Therapie und ihre therapeutischen Anwendungsmöglichkeiten. *Z ärztl Fortbild*, 1983 77, 493-494.
- Krause, WR. Hypnose und Autogenes Training (Selbsthypnose) in der Rehabilitation. Ergänzung in: Schultz, JH. *Hypnose-Technik. Praktische Anleitung zum Hypnotisieren für Ärzte*. 9. Auflage - bearbeitet und ergänzt von G. Iversen und W.-R. Krause. Stuttgart - Jena - New York: Gustav Fischer Verlag; 1994. 71-79.
- Krause, WR. Hypnose und Autogenes Training. *Mitteldeutsch Hausarzt*, 2000 2, 20-22.
- Kreyer, G. Advances in dental psychology. *Z Stomatol*, 1989 86, 123-130.
- Kreyer, G. Zur Inzidenz und Wertigkeit von psychischen, somatischen und psychosomatischen Belastungsfaktoren bei österreichischen Zahnbehandlern. *Z Stomatol*, 1992 89, 319-331.
- Kreyer, G. Psychosomatics of the orofacial system. *Wien Med Wochenschr*, 2000 150, 213-216.
- Kromminga, R; Müller-Fahlbusch, H; Habel, G. Interdisciplinary investigations on the criticism of the term "anesthesia dolorosa". *Dtsch Zahnärztl Z*, 1989 44, 960-961.
- Landt, H. Adaptationsprobleme bei der oralen prothetischen Rehabilitation des alternden Menschen. In: Körber E, editor. *Die zahnärztlich-protetische Versorgung des älternden Menschen*. München, Wien: Carl Hanser Verlag; 1978; 69-92.
- Lang, B; Filippi, A. Halitosis -- Part 1: epidemiology and pathogenesis. *Schweiz Monatsschr Zahnmed*, 2004 114, 1037-1050.
- Langer, A; Michman, J; Seifert, I. Factors influencing satisfaction with complete dentures in geriatric patients. *J. Prosthet Dent*, 1961 11, 1019-1031.
- Laskin, DM. Etiology of the pain-dysfunction syndrome. *JADA*, 1969 79, 147-153.
- Laskin, DM. and Block, S. Diagnosis and treatment of myofacial pain-dysfunction (MPD) syndrome. *J. Prosthet Dent*, 1986 56, 75-85.
- Le Bell, Y; Niemi, PM; Jämsä, T; Kylmä M; Alanen, P. Subjective reactions to intervention with artificial interferences in subjects with and without a history of temporomandibular disorders. *Acta Odontol. Scand.*, 2006 64, 59-63.
- Leao, A. and Sheiham, A. The development of a socio-dental measure of dental impacts on daily living. *Community Dent. Health*, 1996 13, 22-26.
- Lechner, SK. and Roessler, D. Strategies for complete denture success: beyond technical excellence. *Compend Contin. Educ. Dent*, 2001 22, 553-559.

- Lesse, S. Atypical facial pain syndromes of psychogenic origin. Complications of their misdiagnosis. *J. Nerv. Ment Dis.*, 1956 124, 346-351.
- Levin, B. and Landesmann, HM. A practical questionnaire for predicting denture success of failure. *J. Prosthet Dent.*, 1976 35, 124-130.
- Ligier, S. and Stenberg, EM. Immune response. In: Fink G, editor in chief. *Encyclopedia of stress*. (vol. 2.). San Diego: Academic Press; 2000; 523-530.
- Lindberg, NE; Lindberg, E; Larsson, G. Psychologic factors in the etiology of amalgam illness. *Acta Odontol. Scand.*, 1994 52, 219-228.
- Litter, B. Phantom bite revisited. *Brit. Dent J.*, 2005 198, 149.
- Lowenthal, U. and Tau, S. Effects of ethnic origin, age, and bereavement on complete denture patients. *J. Prosthet Dent*, 1980 44, 133-136.
- Lundqvist, S. *Speech and other oral functions. Clinical and experimental studies with special reference to maxillary rehabilitation on osseointegrated implants. (Ph.D. thesis).* Göteborg, 1993.
- Ma, H; Sun, HQ, Ji, P. How to deal with esthetically overcritical patients who need complete dentures: A case report. *J. Contemp Dent. Pract.*, 2008 9, 22-27.
- MacEntee, MI. Measuring the impact of oral health in old age: a qualitative reaction to some quantitative views. *Gerodontology*, 1996 13, 76-81.
- Malt, UF; Nerdrum, P; Oppedal, B; Gundersen, R; Holte, M; Löne, J. Physical and mental problems attributed to dental amalgam fillings: A descriptive study of 99 self-referred patients compared with 272 controls. *Psychosom. Med.*, 1997 59, 32-41.
- Marbach, JJ. Phantom bite syndrome. *Am. J. Psychiatry*, 1978 135, 476-479.
- Martin, SF. T lymphocyte-mediated immune responses to chemical haptens and metal ions: Implications for allergic and autoimmune disease. *Int. Arch. Allergy Immunol.*, 2004 134, 186-198.
- Marxkors, R. Psychische Konditionierung. *ZWR*, 1975 84, 461-462.
- Marxkors, R. and Müller-Fahlbusch, H. Zur Diagnose psychosomatischer Störungen in der zahnärztlich-prothetischen Praxis. *Dtsch Zahnärztl Z.* 1981 36, 787-790.
- Massler, M. Oral manifestations during the female climacteric (the postmenopausa syndrome). *Oral Surg Oral Med Oral Pathol*, 1951 4, 1234-1243.
- Mazurat, NM. and Mazurat, RD. Discuss before fabricating: Communicating the realities of partial denture therapy. Part I: Patient expectations. *J. Can. Dent. Assoc.*, 2003/a 69, 90-94.
- Mazurat, NM. and Mazurat, RD. Discuss before fabricating: Communicating the realities of partial denture therapy. Part II: Clinical outcomes. *J. Can. Dent. Assoc.*, 2003/b 69, 90-94.
- McCullough, MJ. and Tyas, MJ. Local adverse effects of amalgam restorations. *Int. Dent J.*, 2008 58, 3-9.
- McGrath, C. and Bedi, R. A study of the impact of oral health on the quality of life of older people in the UK - findings from a national survey. *Gerodontology*, 1998 15, 93-98.
- McGrath, C. and Bedi, R. The importance of oral health to older people's quality of life. *Gerodontology*, 1999 16, 59-63.
- McGuire, L; Millar, K; Lindsay, S. A treatment trial of an information package to help patients accept new dentures. *Behav. Res. Ther.*, 2007 45, 1941-1948.

- Meldolesi, GN; Picardi, A; Accivile, E; Toraldo di Francia, R; Biondi, M. Personality and psychopathology in patients with temporomandibular joint pain-dysfunction syndrome. *Psychother Psychosom*, 2000 69, 322-328.
- Merskey, H. Pain terms: a list with definitions and notes on usage. Recommended by the IASP sub-committee on taxonomy. *Pain*, 1979 6, 249-252.
- Mew, J. Phantom bite. *Brit. Dent J.*, 2004 197, 660.
- Michman, J. and Langer, A. Postinsertion changes in complete dentures. *J. Prosthet Dent*, 1975 34, 125-134.
- Milgrom, P; Weinstein, P; Kleinknecht, RA; Getz, T. *Treating fearful dental patients: a clinical handbook*. Reston (Virginia): Reston Publishing Company; 1985; 138-142.
- Miller, AA. Psychological Considerations in Dentistry, *JADA*, 1970 81, 941-947.
- Molly, L; Nackaerts, O; Vandewiele, K; Manders, E; van Steenberghe, D; Jacobs, R. Speech adaptation after treatment of full edentulism through immediate loaded implant protocols. *Clin Oral Implant Res*, 2008 19, 86-90.
- Moore, R. and Birn, H. Phenomenon of dental fear. *Tandlaegebladet*, 1990 94, 34-41.
- Moulton, R; Ewen, S; Thieman, W. Emotional factors in periodontal disease. *Oral. Surg. Oral. Med. Oral. Pathol.*, 1957 5, 833-860.
- Müller-Fahlbusch, H. Über situative Provokation endogener depressiver Phase. *Med. Welt*, 1972 23, 919.
- Müller-Fahlbusch, H. Nervenärztliche Befunde bei Prothesenunverträglichkeitserscheinungen. *ZWR*, 1975 84, 574-578.
- Müller-Fahlbusch, H. Nervenärztliche Aspekte der Prothesenunverträglichkeit. *Dtsch Zahnärztl Z*, 1976 31, 13-17.
- Müller-Fahlbusch, H. Psychogenic pain sensation in teeth, mouth and jaws. *Dtsch Zahnärztl Z*, 1991 46, 109-111.
- Müller-Fahlbusch, H. Therapie nach erkannter psychosomatischer Störung. *Dtsch Zahnärztl Z*, 1992 47, 157-161.
- Müller-Fahlbusch, H. and Marxkors, R. *Zahnärztliche Psychagogik. Vom Umgang mit dem Patienten*. München, Wien: Carl Hanser Verlag; 1981.
- Müller-Fahlbusch, H. and Sone, K. Präprotetische Psychagogik. *Dtsch Zahnärztl Z*, 1982 37, 703-707.
- Nagy, G; Bartha, Y; Keresztes, T; Olveti, E; Madléna, M. Quantitative analysis of catecholamines in human dental pulp. *J. Endod*, 2000 26, 596-598.
- Nakaminami, T; Omae, T; Akanishi, M; Maruyama, T. Psychosomatic aspects of the patients with stomatognathic dysfunction. 2. *Nihon Hotetsu Shika Gakkai Zasshi*, 1989 33, 395-400.
- Nalcaci, R. and Baran, I. Factors associated with self-reported halitosis (SRH) and perceived taste disturbance (PTD) in elderly. *Arch. Gerontol. Geriatr*, 2008 46, 307-316.
- Neverlien, PO. Assessment of a single-item dental anxiety question. *Acta Odontol. Scand.*, 1990 48, 365-369.
- Newton, AV. The psychosomatic component in prosthodontics. *J. Prosthet. Dent.*, 1984 52, 871-874.
- Niedermeier, W. Physiology and pathophysiology of the minor salivary glands. *Dtsch. Z Mund Kiefer Gesichtschir*, 1991 15, 6-15.
- Niedermeier, W; Becker, H; Christ, F; Habermann, PG. Effect of menopausal syndromes on symptoms of denture intolerance. *Schweiz Mschr Zahnheilk*, 1979 89, 1011-1018.

- Niedermeier, W; Huber, M; Fischer, D; Beier, K; Müller, N; Schuler, R; Brinninger, A; Fartasch, M; Diepgen, T; Matthaeus, C; Meyer, C; Hector, MP. Significance of saliva for the denture-wearing population. *Gerodontology*, 2000 17, 104-118.
- Ogawa, T; Tanaka, M; Ogimoto, T; Okushi, N; Koyano, K; Takeuchi, K. Mapping, profiling and clustering of pressure pain threshold (PPT) in edentulous oral mucosa. *J. Dent.istry*, 2004 32, 219-228.
- Oka, T. and Oka, K. Age and gender differences of psychogenic fever: a review of the Japanese literature. *BMC BioPsychoSocial Med.*, 2007 1, 11.
- Pertes, RA. Differential diagnosis of orofacial pain. *Mt Sinai J. Med.*, 1998 65, 348-354.
- Pfeiffer, P. and Schwickerath, H. Nickel solubility and metallic taste. *ZWR*, 1991 100, 762-770.
- Piquero, K; Ando, T; Sakurai, K. Buccal mucosa ridging and tongue indentation: incidence and associated factors. *Bull Tokyo Dent Coll*, 1999 40, 71-78.
- Pomp, AM. Psychotherapy for the myofascial pain-dysfunction syndrome: a study of factors coinciding with symptom remission. *JADA*, 1974 89, 629-632.
- Reeve, PE; Watson, CJ; Stafford, GD. The role of personality in the management of complete denture patients. *Brit. Dent. J.*, 1984 156, 356-362.
- Reeves 2nd, JL. and Merrill, RL. Diagnostic and treatment challenges in occlusal dysesthesia. *J. Calif. Dent. Assoc.*, 2007 35, 198-207.
- Reid, KI; Mueller, PS; Barnes, SA. Attitudes of general dentists regarding the acceptance of gifts and unconventional payment from patients. *JADA*, 2007 138, 1127-1133.
- Ross, GL; Bentley, HJ; Greene Jr., GW. The psychosomatic concept in dentistry. *Psychosom Med*, 1953 15, 168-173.
- Roumanas, ED. The social solution - denture esthetics, phonetics, and function. *J. Prosthodont*, 2009 18, 112-115.
- Sandler, NA; Ziccardi, V; Ochs, M. Differential diagnosis of jaw pain in the elderly. *JADA*, 1995 126, 1263-1272.
- Sanz, M; Roldán, S; Herrera, D. Fundamentals of breath malodor. *J Contemp Dent Pract*, 2001 2, 1-17.
- Sarlani, E; Balciunas, BA; Grace, EG. Orofacial pain - Part I. Assessment and management of musculoskeletal and neuropathic causes. *AACN Clin. Issues*, 2005/a 16, 333-346.
- Sarlani, E; Balciunas, BA; Grace, EG. Orofacial pain - Part II. Assessment and management of vascular, neurovascular, idiopathic, secondary, and psychogenic causes. *AACN Clin Issues*, 2005/b 16, 347-358.
- Sato, C; Sato, S; Takashina, H; Ishii, H; Onozuka, M; Sasaguri, K. Bruxism affects stress responses in stressed rats. *Clin Oral Invest*, in press. DOI: 10.1007/s00784-009-0280-6
- Sauerland, EK. and Mizuno, N. A protective mechanism for the tongue. Suppression of the genioglossal activity induced by stimulation of trigeminal proprioceptive afferents. *Experientia*, 1970 26, 1226-1227.
- Savage, RD. and MacGregor, AR. Behavior therapy in prosthodontics. *J. Prosthet Dent*, 1970 34, 126-132.
- Schmitt, A; Yu, SKJ; Sessle, BJ. Excitatory and inhibitory influences from laryngeal and orofacial areas on tongue position in the cat. *Arch. Oral. Biol.*, 1973 18, 1121-1130.
- Schwartz, LL. Pain associated with the temporomandibular joint. *JADA*, 1955 51, 394-397.
- Schweitzer, JM. *Oral rehabilitation problem cases. Treatment and evaluation*. Vol. 2. Saint Louis: C.V. Mosby Company; 1964; 611-625.

- Schwichtenberg, J. and Doering, S. Success of referral in a psychosomatic-psychotherapeutic outpatient unit of a dental school. *Z Psychosom Med. Psychother*, 2008 54, 285-292.
- Scott, A; Egner, W; Gawkrödger, DJ; Hatton, PV; Sherriff, M; van Noort, R; Yeoman, C; Grummit, J. The national survey of adverse reactions to dental materials in the UK: a preliminary study by the UK Adverse Reactions Reporting Project. *Brit. Dent J.*, 2004 196, 471-477
- Seifert, I; Langer, A; Michmann, J. Evaluation of psychologic factors in geriatric denture patients. *J. Prosthet Dent*, 1962 12, 516-523.
- Seifert, E; Runte, C; Selders, D; Lamprecht-Dinnesen, A; Bollmann, F. Der Einfluss der Zahnprothese auf die Stimme. *HNO*, 1999/a 47, 485-489.
- Seifert, E; Runte, C; Riebandt, M; Lamprecht-Dinnesen, A; Bollmann, F. Can dental prostheses influence vocal parameters? *J. Prosthet Dent*, 1999/b 81, 579-585.
- Shigli, K; Shivappa Angadi, G; Hedge, P; Hebbal, M. Patients' knowledge and understanding of the implications of wearing dentures. Report of a survey conducted at a dental institute in the south of India. *Prim. Dent Care*, 2008 15, 85-89.
- Sieber, M; Grubenmann, E; Ruggia, GM; Palla, S. Relation between stress and symptoms of craniomandibular disorders in adolescents. *Schweiz Monatsschr Zahnmed*, 2003 113, 648-654.
- Silverman, S; Silverman, SI; Silverman, B; Garfinkel, L. Self-image and its relation to denture acceptance. *J. Prosthet Dent*, 1976 35, 131-141.
- Smith, DV; Li, CS; Davis, BJ. Excitatory and inhibitory modulation of taste responses in the hamster brainstem. *Ann. NY Acad. Sci.*, 1998 855, 450-456.
- Somer, E. Hypnotherapy in the treatment of the chronic nocturnal use of a dental splint prescribed for bruxism. *Int. J. Clin. Exp. Hypn*, 1991 39, 145-154.
- Somer, E; Ben-Aryeh, H; Laufert, D. Salivary composition, gender, and psychosocial stress. *Int. J. Psychosom*, 1993 40, 17-21.
- Stein, RS. Mutual protective complex of dental restorations. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment of prosthodontics*. Philadelphia: Lea and Febiger; 1983; 306-326.
- Suda, M; Morimoto, K; Obata, A; Koizumi, H; Maki, A. Emotional responses to music: towards scientific perspectives on music therapy. *NeuroReport*, 2008 19, 75-78.
- Sugawara, N; Shiozawa, I; Masuda, T; Takei, H; Tsuruta, J; Ogura, N; Hasegawa, S. A survey on condition of outpatients at prosthodontics II, University Hospital, Faculty of Dentistry, Tokyo Medical and Dental University. *Kokubyo Gakkai Zasshi*, 1998 65, 251-259.
- Svensson, P. and Kaaber, S. General health factors and denture function in patients with burning mouth syndrome and matched control subjects. *J. Oral. Rehabil.*, 1995 22, 887-895.
- Swoope, CC. Identification and management of emotional patients. *J. Prosthet Dent*, 1972 27, 434-440.
- Swoope, CC. Predicting denture success. *J. Prosthet Dent*, 1973 30, 860-865.
- Szentpétery, AG; John, MT; Slaze, GD; Setz, JM. Problems reported by patients before and after prosthodontic treatment. *Int. J. Prosthodont*, 2005 18, 124-131.
- Tau, S. and Lowental, U. Some personality determinants of denture preference. *J. Prosthet Dent*, 1980 44, 10-12.

- Tickle, M; Craven, R; Worthington, HV. A comparison of the subjective oral health status of older adults from deprived and affluent communities. *Community Dent Oral Epidemiol*, 1997 25, 217-222.
- Toyofuku, A. and Kikuta, T. Treatment of phantom bite syndrome with milnacipran - a case series. *Neuropsych. Dis. Treatment*, 2006 2, 387-390.
- Troest, T. Form und Funktion im stomatognathen System. In: Koeck B, editor. *Funktionsstörungen des Kauorgans*. München - Wien - Baltimore: Urban and Schwarzenberg; 1995; 11-26.
- Trulsson, U; Engstrand, P; Berggren, U; Nannmark, U; Brånemark, PI. Edentulousness and oral rehabilitation: experiences from the patients' perspective. *Eur. J. Oral. Sci.*, 2002 110, 417-424.
- Turner, JA; Mancl, L; Aaron, LA. Short- and long-term efficacy of brief cognitive-behavioral therapy for patients with chronic temporomandibular disorder pain: A randomized, controlled trial. *Pain*, 2006 121, 181-194.
- Vanhaudenhuyse, A; Boly, M; Laureys, S; Faymonville, ME. Neurophysiological correlates of hypnotic analgesia. *Contemp Hypnosis*, 2009 26, 15-23.
- Vinard, H. and Ravier-Rosenblum, C. The psychosomatic effects of the sonic environment in the dental office. *Rev. Odontostomatol. (Paris)*, 1989 18, 101-108.
- Violon, A. The onset of facial pain. A psychological study. *Psychother Psychosom*, 1980 34, 11-16.
- Von Korff, M; Balderson BHK; Saunders, K; Miglioretti, DL; Lin, EHB; Berry, S; Moore, J; Turner, JA. A trial of an activating intervention for chronic back pain in primary care and physical therapy settings. *Pain*, 2005 113, 323-330.
- Watanabe, I. Masticatory function and life style in aged. *Nippon Rogen Igakkai Zasshi*, 1998 35, 194-200.
- Weinstein, P; Milgrom, P; Ratener, P; Morrison, K. Patient dental values and their relationship to oral health status, dentist perceptions and quality of care. *Community Dent Oral. Epidemiol.*, 1979 7, 121-127.
- Wentz FM. Patient motivation: a new challenge to the dental profession for effective control of plaque. *JADA*, 1972 85, 887-891.
- Woelfel, JB. Denture base materials and their effects on oral tissues. In: Laney WR, Gibilisco JA editors. *Diagnosis and treatment in prosthodontics*. Philadelphia: Lea and Febiger; 1983; 235-252.
- Wolff, A; Gadre, A; Begleiter, A; Moskona, D; Cardash, H. Correlation between patient satisfaction with complete dentures and denture quality, oral condition, and flow rate of submandibular/sublingual salivary glands. *Int. J. Prosthodont*, 2003 16, 45-48.
- Wolowski, A. Orale Implantologie bei Patienten mit orofazialer Somatisierungsstörung. *Stomatologie*, 2000 97, 11-16.
- Wöstmann, B. Psychogene Zahnersatzunverträglichkeit. In: Sergl HG, editor. *Psychologie und Psychosomatik in der Zahnheilkunde*. München - Wien - Baltimore: Urban and Schwarzenberg; 1996; 187-213.
- Yaegaki, K. and Coil, JM. Clinical application of a questionnaire for diagnosis and treatment of halitosis. *Quintessence Int.*, 1999/a 30, 302-306.
- Yaegaki, K. and Coil, JM. Clinical dilemmas posed by patients with psychosomatic halitosis. *Quintessence Int.*, 1999/b 30, 328-333.

-
- Yaegaki, K. and Coil, JM. Genuine halitosis, pseudo-halitosis, and halitophobia: classification, diagnosis, and treatment. *Compend Contin Educ. Dent*, 2000/a 21, 880-886, 888-889, 890.
- Yaegaki, K. and Coil, JM. Examination, classification, and treatment of halitosis; Clinical perspectives. *J. Can. Dent. Assoc.*, 2000/b 66, 257-261.
- Zimmermann, M. Physiologie von Nozizeption und Schmerz. In: Basler HD, Franz C, Kröner-Hervig B, Rehfisch HP editors. *Psychologische Schmerztherapie*. Berlin: Springer; 2004; 17-58.

Chapter XIII

Pulsed Electromagnetic Field (PEMF) Therapy for Peri-Implant Healing and Bone Remodeling

***Tibor Károly Fábián^{*1}, Anita Beck^{1,2}, Gábor Fábián³,
Pál Fejérdy¹ and Péter Hermann¹***

¹Clinic of Prosthetic Dentistry,

²Department of Oral Biology,

³Clinic of Pediatric Dentistry and Orthodontics

Semmelweis University Budapest, Faculty of Dentistry, Budapest, Hungary, EU

Abstract

Data in the literature revealed that bone formation around dental implants can be promoted due to PEMF treatment significantly. Since PEMF decreases the rate of residual ridge resorption following tooth extraction, PEMF is likely to improve healing of implants inserted into extraction sites as well. PEMF may also be used advantageously for accelerating bone healing following advanced implant surgery. Although PEMF stimulation during the first two weeks after surgery seems to be the most efficient; late phase bone healing (even four weeks after surgery) may also be accelerated with PEMF administration. Moreover, PEMF stimulation speeds up bone remodeling processes which effect is likely to be a crucial factor of the phenomenon that PEMF accelerates tooth movement (and coupled periodontal processes). PEMF also influence the function of periodontal ligament fibroblasts, which also may play a significant role in the acceleration of bone remodeling around orthodontically moved teeth as well as around teeth subjected to high mechanical load during mastication (i.e. abutment teeth) or because of parafunction. Although PEMF was found to be beneficial to a wide variety of therapeutic processes also in dentistry; the exact mechanisms by which PEMF affects alveolar bone and other periodontal tissues is far not fully understood yet. However,

* E-mail address: fab@fok.usn.hu.

certain mechanisms are already recognized and the increased amount of knowledge may be utilized also for the clinical practice. In this chapter scientific data of clinical relevance are collected and discussed.

Introduction

How a cell detects a PEMF signal is unknown currently, although some mechanisms have been proposed to account for the interaction of PEMF with cells. PEMF may interact with moving charges and change their velocities as in the classic interaction of magnetic fields with any moving charge [Blank 1995, Goodman and Blank 1998]. Resonance-interaction of PEMF fields with Earth's magnetic field [Goodman and Blank 1998] as well as the phenomenon of ion-cyclotron resonance [Liboff et al. 1987, Lednev 1991, Blanchard and Blackman 1994] may also be responsible for several biological responses. Since PEMF penetrate the cell (unlike electric fields) interaction can occur anywhere inside (or outside) the cell [Goodman and Blank 1998]. It is very likely that PEMF signal exhibit sufficient energy to elicit a transient activation/alteration of membrane bound receptors or their inner associated proteins [Luben 1982, Varani et al. 2003, Schnoke and Midura 2007] and may also alter density of certain receptors of the cells [Varani et al. 2003]. Therefore, PEMF is believed to influence the function of receptor coupled signal transduction pathways [Luben 1982, Goodman and Blank 1998, Varani et al. 2003, Schnoke and Midura 2007]. A wide variety of cell-membrane structural and functional properties are also altered by PEMF including binding of Ca^{2+} to anionic fixed charges in the cell membrane [Lednev 1991], conformational changes of ion channels [Balcavage et al. 1996] and trans membrane ion transport [Lednev 1991]. PEMF significantly increases intracellular free Ca^{2+} level and decreases Ca^{2+} dependent signal transduction responses to receptor ligands present in the serum as well [Satake et al. 1990]. Alteration of cell communication via alteration of local factor production and gap junction signaling is also very likely [Lohmann et al. 2003]. Further, PEMF is likely to induce phosphorylation of several proteins of intracellular signal transduction pathways [Schnoke and Midura 2007]. PEMF induced changes of the activity of several ATPases and enzymes is also very likely [Blank 1995, Goodman and Blank 1998]. PEMF is also likely to influence gene activation [McClellan et al. 1990, Blank and Goodman 1997], transcription [Blank and Goodman 1997, Tuinstra et al. 1997, Solazzo et al. 2010] as well as translation and protein synthesis [Goodman et al 1993, Goodman and Blank 1998] pathways of the cell.

Effect of PEMF on Osteoclasts

Osteoclasts are responsible for both bone demineralization and collagen degradation in the bone [Chang et al. 2005, Fábíán et. al. 2009/a, Barnaba et al. in press]. Osteoclasts are bone resorptive large multinucleated cells derived from bone marrow hematopoietic stem cells of monocyte-macrophage lineage [Chang et al. 2005, Fábíán et. al. 2009/a, Barnaba et al. in press]. The fact that immune cells and osteoclast precursor cells are derived from the same hematopoietic precursor cells may be responsible for the phenomenon that osteoclastogenesis is promoted by macrophage colony-stimulating factor (M-CSF) [Chang et

al. 2005] and also that pro-inflammatory cytokine network responsible for the activation of immune cells exert an activating effect also on osteoclast precursor cells [Fábián et. al. 2009/a]. Production of proinflammatory cytokines is an important factor also for the maturation of osteoclasts in the bone, because tumor necrosis factor alpha (TNF- α) interleukin 1 β (IL-1 β) and interleukin 6 (IL-6) induce the expression of RANKL (receptor activator of nuclear factor- κ B ligand; a member of tumor necrosis factor superfamily) in osteoblasts which is essential for final differentiation steps of osteoclasts (as well as for their bone resorbing capacity) [Fábián et. al. 2009/a]. Osteoblasts have RANKL on their surfaces and via contact dependent signaling they activate RANK /receptor activator of nuclear factor- κ B/ on the surface of osteoclast progenitors, resulting in osteoclast maturation and bone resorption [Chang et al. 2005, Schwartz et al. 2009]. In addition, osteoblasts also secrete soluble RANKL [Chang et al. 2005, Schwartz et al. 2009]. Provided that macrophage colony-stimulating factor (M-CSF) is present, RANKL is both necessary and sufficient for osteoclastogenesis [Chang et al. 2005]. Osteoblasts however also produce osteoprotegerin /OPG/ which acts as a decoy (scavenger) receptor binding RANKL, thereby preventing it from binding RANK on the surface of osteoclasts resulting in decreased osteoclast maturation and bone resorption [Chang et al. 2005, Schwartz et al. 2009].

PEMF is likely to cause a delay of osteoclast differentiation [Chang et al. 2004/a, 2005, Barnaba et al. in press] coupled with decreased proinflammatory cytokine (TNF- α , IL-1 β , IL-6) production [Chang et al. 2004/a] and decreased RANKL and MC-SF production [Chang et al. 2005] of differentiating cells in the first week (7-9 days) of daily PEMF exposure. PEMF also increases the production of osteoprotegerin (OPG) in osteoblasts without any change of secretion of RANKL [Schwartz et al. 2009], leading to the increase of OPG/RANKL ratio resulting in decreased osteoclast maturation and decreased bone resorption [Schwartz et al. 2009]. However, the effect may be transitory in nature and may disappear thereafter [Barnaba et al. in press]; and the effect seems to be coupled with PEMF stimulation of lower energy [Chang et al. 2003, 2005]. (In contrast, stimulation with PEMF of higher energy may lead to increased osteoclast differentiation [Chang et al. 2003, 2005] coupled with increased proinflammatory cytokine (TNF- α , IL-1 β) production [Chang et al. 2003] and decreased secretion of OPG and decrease of OPG/RANKL ratio [Chang et al. 2005].) Besides premised effects of PEMF on osteoclast differentiation, PEMF may also increase the apoptotic rate of osteoclasts which may also lead to decrease of bone degradation under certain circumstances [Chang et al. 2006].

Effect of PEMF on Osteoblasts

Osteoblasts produce bone matrix during development, after bone injury, and during normal bone remodeling [Chang et al 2005, Fábián et. al. 2009/a]. In contrast to osteoclasts derived from bone marrow hematopoietic stem cells of monocyte-macrophage lineage [Chang et al. 2005, Fábián et. al. 2009/a, Barnaba et al. in press], osteoblasts arise from multipotential mesenchymal (stromal) cells in bone marrow [Chang et al 2005, Fábián et. al. 2009/a]. PEMF exposure of bone marrow-derived mesenchymal stem cells (cultured in osteogenic medium) likely decreases the rate of their proliferation [Tsai et al. 2009, Jansen et al. 2010] and speeds up their osteogenic differentiation [Tsai et al. 2009, Jansen et al. 2010] in the first week (7-9

days) of daily exposure [Tsai et al. 2009, Jansen et al. 2010]. Expressed OPG/RANKL ratio also increases during this period significantly [Schwartz et al. 2009, Jansen et al. 2010]. Importantly, these effects may be transitory in nature and may disappear thereafter [Tsai et al. 2009, Jansen et al. 2010] in good accordance with the effect of PEMF on osteoclast maturation (see above). Notwithstanding that premised effects maybe be transitory, the shift towards earlier maturation is resulting in a significantly enhanced calcium deposition in the extracellular matrix also thereafter (i.e. after the 7-9-th days of stimulation as well) [Jansen et al. 2010].

There is also a stimulatory effect of PEMF on matured cell cultured osteoblasts. Daily exposure with PEMF increases proliferation of osteoblasts in vitro during the first week (2-7 days) of active proliferation stage of cell culture [Diniz et al. 2002, Chang et al. 2004/b], but importantly this effect disappears thereafter [Diniz et al. 2002, Chang et al. 2004/b]. PEMF also increases the expression of OPG and decreases the expression of RANKL (with a consequent increase of OPG/RANKL ratio) in the first 5 days of daily stimulation but not thereafter [Chang et al. 2004/b]. PEMF also enhances mineralized extracellular matrix formation of cultured osteoblasts in the first 15 days but not thereafter [Diniz et al. 2002]. Increase of extracellular matrix formation is likely to be based on upregulation of important genes related to bone formation (including genes responsible for the synthesis of both collagenous and noncollagenous matrix components) and downregulation of genes responsible for the degradation of extracellular matrix [Solazzo et al. 2010]. Besides above, daily PEMF stimulation also enhances cell proliferation and viability of osteoblasts co-cultured with macrophage cells stimulated with lipopolysaccharide (to release high level of reactive oxygen species as a model of inflammatory environment) during the first 7 days of culture [Lin and Lin in press]. PEMF stimulation also induces release of a soluble factor from osteoblasts, which significantly increase vascular endothelial proliferation (a major factor of angiogenesis) in a paracrine manner [Hopper et al. 2009]. PEMF also likely to inhibit parathyroid hormone (PTH) induced release of osteoclast activating factor [Luben et al 1982, Spadaro and Bergstrom 2002], which prevent PTH induced Ca^{2+} mobilization via inhibiting resorption of the bone by osteoclasts [Spadaro and Bergstrom 2002]. (Please consider that the response of osteoclasts to PTH primarily depends on the activating factor generated by osteoblasts which have PTH receptors lacking in osteoclasts [Spadaro and Bergstrom 2002].)

PEMF Therapy and Peri-Implant Healing

Data in the literature revealed that bone formation around dental implants (i.e. bone coverage and osteointegration of implants) can be promoted due to PEMF treatment significantly [Matsumoto et al. 2000, Chan and Bergman 2008, Grana et al. 2008]. Since PEMF decreases the rate of residual ridge resorption following tooth extraction [Ortman et al. 1992], PEMF is likely to improve periimplant healing of implants inserted into extraction sites as well. In general, daily PEMF exposure of 1 to 2 hours may be recommended based on the scientific literature (2 x 30 min [Grana et al. 2008]; 1 hour [Friedricks et al. 2000, Inoue et al. 2002]; 2 hours [Chan and Bergman 2008]). However, authors experience indicated that PEMF stimulation of 10-15 minutes three times a week also appears to be promising for clinical use. Although stimulations during the first two weeks after surgery seems to be the

most efficient [Matsumoto et al. 2000]; late phase bone healing (even four weeks after surgery) may also be accelerated with PEMF administration [Inoue et al. 2002, Chan and Bergman 2008]. PEMF stimulation of late phase bone healing (1 hour per day for 8 weeks [Inoue et al. 2002]) very likely leads to somewhat greater new bone formation, increased mineral apposition rate and decreased porosity of the cortex adjacent to the healing area [Inoue et al. 2002].

It is likely that, energy value of PEMF stimulation as well as pattern of waveform characteristics including shape and amplitude of the bursts, pulse frequency of the bursts, duration of the bursts, and length of burst on-off periods (burst on-off frequency) are rather important factors of PEMF induced promotion of bone healing [Leisner et al. 2002, Miruda et al. 2005]. It is also likely that high-amplitude narrow-pulse portions of PEMF stimulation are especially advantageous to produce a beneficial bone response [Pienkowski et al. 1992]; whereas symmetry/asymmetry relations of the stimulus pulse waveform may be of less importance [Pienkowski et al. 1992]. Further, low energy fields (0.2 – 0.3 mT [Matsumoto et al. 2000]) seems to be more efficient than high energy fields (0.8 mT [Matsumoto et al. 2000]) to promote bone healing procedures [Rubin et al. 1993, Matsumoto et al. 2000]. Authors experience indicated that PEMF stimulation with very low energy fields (0.05 – 0.1 mT) also appears to be promising for clinical use. It is also very likely that low values (15 Hz) of burst on-off frequencies seems to be more efficient than high values (50 – 150 Hz) of burst on-off frequencies [Rubin et al. 1993]; although higher burst on-off frequencies also promote bone healing significantly [Rubin et al. 1993, Matsumoto et al. 2000].

PEMF may also be used advantageously for accelerating bone healing following advanced implant surgery. PEMF enhances healing of osteotomy gaps [Pienkowski et al. 1992, Darendeliev et al. 1997, Friedrichs et al. 2000, Inoue et al. 2002, Grana et al. 2008]. PEMF can also accelerate bone formation (and total recovery of bone tissue) in bone defects filled with hydroxyapatite (HA) [Ottani et al. 2002] or with demineralized bone-matrix [Takano-Yamamoto et al. 1992]. Similarly, PEMF can also accelerate bone graft incorporation [Kold et al. 1987]. Further, PEMF accelerate consolidation of regenerate bone during distraction osteogenesis [Fredericks et al. 2003] and accelerate healing of both fresh [Grace et al. 1998] and ununited bone fractures [Gossling et al. 1992]. PEMF also alleviate postoperative pain [Hedén and Pilla 2008, Strauch et al. 2009] and decreases postoperative edema [Strauch et al. 2009] which could also be rather advantageous following implant surgery.

Besides above, PEMF also improves wound healing of the gastrointestinal mucosa [Mentes et al. 1996] and the skin [Patino et al. 1996, Scardino et al. 1998], therefore it may be hypothesized that PEMF improves healing of the oral mucosa around osseointegrated implants. PEMF also improves regeneration of injured nerves [Raji and Bowden 1983, Raji 1984, Zienowicz et al. 1991, Kanje et al. 1993] and increases angiogenesis [Tepper et al. 2004, Hopper et al. 2009] which could be important additional factors of periimplant bone and soft tissue healing as well. Further, PEMF reverses the proliferative defects of lymphocytes from aged subjects [Cossarizza et al. 1989/a,b], and induces the expression (and consequent release) of immune activator/modulator HSP70/HSPA type stress proteins [Goodman and Blank 1998, Heredia-Rojas et al. 2010] which are also likely to improve proper bone [Fabián et al. 2009/a,b] and mucosal healing [Fabián et al. 2003, 2007, 2009/b, 2011] around the implants.

PEMF Therapy and Bone Remodeling

Bone is remodeled continuously throughout life through the resorption of old bone by osteoclasts and the subsequent formation of new bone by osteoblasts [Chang et al. 2004/a, Fábíán et. al. 2009]. These two events are tightly coupled to each other and are responsible for the renewal of bone while maintaining its anatomical and structural integrity [Chang et al. 2004/a, Fábíán et. al. 2009]. As mentioned above, PEMF stimulation with low energy fields usually leads to the depression of osteoclast function [Chang et al. 2003, 2005] which may slow down bone remodeling processes. However stimulation with PEMF of higher energy [Chang et al. 2003, 2005] and/or with certain specific patterns of PEMF waveform characteristics [Leisner et al. 2002, Miruda et al. 2005] can lead to increased osteoclast differentiation [Stark and Sinclair 1987, Chang et al. 2003, 2005] coupled with increased proinflammatory cytokine (TNF- α , IL-1 β) production [Chang et al. 2003] and decreased secretion of OPG and decrease of OPG/RANKL ratio [Chang et al. 2005]. Moreover, PEMF stimulation induced moderate increase of osteoclast function coupled with a somewhat more increased osteoblast function likely speeds up bone remodeling processes [Fábíán and Söti 2007].

Speed up of bone remodeling is likely to be crucial factor of the phenomenon that PEMF accelerates tooth movement and coupled periodontal processes [Stark and Sinclair 1987, Chen 1991, Darendeliler et al. 1995, 2007, Showkatbakhsh et al 2010]. Under normal conditions of orthodontic tooth movement (without use of PEMF) osteoclasts proliferate and initial resorption of superficial bone occurs in the area of periodontal ligament compression [Darendeliler et al. 2007]. Accordingly, in the area of periodontal ligament tension, the periodontal fibers unwind, fibroblasts appear and osteoblasts form a non-mineralized collagenous matrix (osteoid) which is mineralized later, trapping some osteocytes in lacunae within the newly formed bone [Darendeliler et al. 2007]. PEMF induced speed up of tooth movement is likely to be primarily due to a further increase of the number of osteoclasts [Stark and Sinclair 1987, Chen 1991] and other active cells [Kim 1990, Chen 1991] without induction of any unfavorable ultrastructural changes of the stimulated cells at issue [Chen 1991].

PEMF is also likely to influence the function of periodontal ligament fibroblasts [Kim 1990, Satake et al. 1990], which may play a significant role in the acceleration of bone remodeling around orthodontically moved teeth [Satake et al. 1990, Darendeliler et al. 2007] as well as around teeth subjected to high mechanical load during mastication (i.e. abutment teeth) and/or because of parafunction [Kim 1990]. PEMF stimulation of periodontal ligament fibroblasts significantly increases their intracellular free Ca^{2+} level and decreases their Ca^{2+} dependent signal transduction responses to receptor ligands present in the serum [Satake et al. 1990], with a duration of roughly a day following stimulation [Satake et al. 1990]. PEMF also induce release of local factors from periodontal ligament fibroblasts, including those influencing cell attachment and (in vitro) spreading of both periodontal ligament fibroblasts and osteoblasts [Kim 1990]. Although meaning of premised effects of PEMF on periodontal ligament fibroblast is far not clear yet, premised data may indicate that PEMF significantly influences function of periodontal ligament fibroblast playing significant role in periodontal bone remodeling processes.

Contraindication of PEMF Therapy

In general, PEMF is contraindicated in the case of patients with any severe systemic diseases and with strongly compromised health [Fábián and Sóti 2007, Fábián and Fejérdy 2010]. In the case of such patients, use of PEMF should be avoided in the absence of a clear agreement of the medical professional(s) responsible for the treatment of the systemic disease(s) at issue [Fábián and Sóti 2007, Fábián and Fejérdy 2010]. Similarly, PEMF is usually contraindicated in the cases of patients with any hemorrhagic diathesis, precancerosis, and malignancy as well as in the case of patients having any oncological treatment in the anamnesis (at least in the absence of a clear agreement of the oncologist or other responsible medical professional) [Fábián and Sóti 2007, Fábián and Fejérdy 2010]. Importantly, use of PEMF should be strictly avoided near to a nevus [Fábián and Sóti 2007, Fábián and Fejérdy 2010]. Use of PEMF should also be avoided in the condylar area during the growth period [Wilmot et al. 1993]. In the absence of clear evidences about the innocuity, PEMF may be contraindicated for patients with pacemaker (especially in the case of unipolar electrode configuration [Gwechenberger et al. 2006]) or with any another implanted electric instruments [Gwechenberger et al. 2006, Fábián and Sóti 2007, Fábián and Fejérdy 2010]. Although application of physiotherapy should be avoided near to the ear, eye, brain or spinal cord (i.e. temporal or paravertebral use) in the absence of clear evidence about the innocuity of the chosen treatment modality [Fábián and Sóti 2007, Fábián and Fejérdy 2010], PEMF seems to be a rather safe treatment modality from these points of view.

Conclusion

PEMF was found to be beneficial to a wide variety of therapeutic processes in the last few decades. Although exact mechanisms by which PEMF affects alveolar bone and other periodontal tissues is far not fully understood yet; certain important mechanisms were recognized recently. Therefore a significantly increased amount of knowledge may be utilized to improve clinical efficiency of PEMF in dentistry. Although PEMF stimulation can be seen as a noninvasive tool, the few contraindications of PEMF should be considered carefully for a safe and successful use.

References

- Balcavage, W; Alvager, T; Swez, J; Goff, C; Fox, M; Abdullyava, S; King, M. A mechanism for action of extremely low frequency electromagnetic fields on biological systems. *Biochem. Biophys. Res. Communication*, 1996 222 (suppl.), 374-378.
- Barnaba, SA; Ruzzini, L; Di Martino, A; Lanotte, A; Sgambato, A; Denaro, V. Clinical significance of different effects of static and pulsed electromagnetic fields on human osteoclast cultures. *Rheumatol Int*, in press, DOI: 10.1007/s00296-010-1724-7.

- Blanchard, J. and Blackman, CF. Clarification and application of an ion parametric resonance model for magnetic field interactions with biological systems. *Bioelectromagnetics*, 1994 15, 205-216.
- Blank, M. Na/K-adenosine-triphosphatase. *Adv. Chem.*, 1995 250, 339-348.
- Blank, M. and Goodman, R. Do electromagnetic fields interact directly with DNA? *Bioelectromagnetics*, 1997 18, 111-115.
- Chan, AY; Bergman, H. Performance verification of a prototype non-invasive intra-oral bone growth stimulator for titanium dental implants. *Conf. Proc. IEEE Eng. Med. Biol. Soc.* 2008, 5624-5627.
- Chang, K; Chang, WHS; Wu, ML; Shih, C. Effects of different intensities of extremely low frequency pulsed electromagnetic fields on formation of osteoclast-like cells. *Bioelectromagnetics*, 2003 24, 431-439.
- Chang, K; Chang, WHS; Yu, YH; Shih, C. Pulsed electromagnetic field stimulation of bone marrow cells derived from ovariectomized rats affects osteoclast formation and local factor production. *Bioelectromagnetics*, 2004/a 25, 134-141.
- Chang, WHS, Chen LT; Sun, JS; Lin, FH. Effect of pulse-burst electromagnetic field stimulation on osteoblast cell activities. *Bioelectromagnetics*, 2004/b 25, 457-465.
- Chang, K; Chang, WHS; Huang, S; Huang, S; Shih, C. Pulsed electromagnetic fields stimulation affects osteoclast formation by modulation of osteoprotegerin, RANK ligand and macrophage colony-stimulating factor. *J. Orthop. Res.*, 2005 23, 1308-1314.
- Chang, K; Chang, WHS; Tsai, MT; Shih, C. Pulsed electromagnetic fields accelerate apoptotic rate in osteoclasts. *Connect Tissue Res*, 2006 47, 222-228.
- Chen, Q. Effect of pulsed electromagnetic field on orthodontic tooth movement through transmission electronmicroscopy. *Zhonghua Kou Qiang Yi Xue Za Zhi*, 1991 26, 7-10, 61.
- Cossarizza, A; Monti, D; Bersani, F; Cantini, M; Cadossi, R; Sacchi, A; Franceschi, C. Extremely low frequency pulsed electromagnetic fields increase cell proliferation in lymphocytes from young and aged subjects. *Biochem. Biophys. Res. Commun.*, 1989/a 160, 692-698.
- Cossarizza, A; Monti, D; Bersani, F; Paganelli, R; Montagnani, G; Cadossi, R; Cantini, M; Franceschi, C. Extremely low frequency pulsed electromagnetic fields increase interleukin-2 (IL-2) utilization and IL-2 receptor expression in mitogen-stimulated human lymphocytes from old subjects. *FEBS Lett*, 1989/b 248, 141-144.
- Darendeliler, MA; Sinclair, PM; Kusy, RP. The effects of samarium-cobalt magnets and pulsed electromagnetic fields on tooth movement. *Am. J. Orthod. Dentofacial. Orthop.*, 1995 107, 578-588.
- Darendeliler, MA; Zea, A; Shen, G; Zoellner, H. Effects of pulsed electromagnetic field vibration on tooth movement induced by magnetic and mechanical forces: a preliminary study. *Australian Dent. J.*, 2007 52, 282-287.
- Darendeliler, MA; Darendeliler, A; Sinclair, PM. Effects of static magnetic and pulsed electromagnetic fields on bone healing. *Int. J. Adult Orthodont Orthognath Surg.*, 1997 12, 43-53.
- Diniz, P; Shomura, K; Soejima, K; Ito, G. Effects of pulsed electromagnetic field (PEMF) stimulation on bone tissue like formation are dependent on the maturation stages of the osteoblasts. *Bioelectromagnetics*, 2002 23, 398-405.

- Fábián, TK. and Fejérdy P. *Psychogenic denture intolerance. Theoretical background, prevention and treatment possibilities*. New York: Nova Science Publishers; 2010. 162-170.
- Fábián, TK. and Söti, Cs. Low induction value pulsed electromagnetic field stimulation. [Alacsony indukció értékű mágneses impulzus tér stimuláció] In: Vértés G, Fábián TK editors. *Psychosomatic dentistry [Fogorvosi Pszichoszomatika]* Budapest: Medicina; 2007; 201-210.
- Fábián, TK; Gáspár, J; Fejérdy, L; Kaán, B; Bálint, M; Csermely, P; Fejérdy, P. Hsp70 is present in human saliva. *Med. Sci. Monit.*, 2003 9, BR62-65.
- Fábián, TK; Fejérdy, P; Nguyen, MT; Söti, Cs; Csermely, P. Potential immunological functions of salivary Hsp70 in mucosal and periodontal defense mechanisms. *Arch. Immunol. Ther. Exp.*, 2007 55, 91-98.
- Fábián, TK; Csermely, P; Fábián, G; Fejérdy, P. Spondyloarthropathies and bone resorption: A possible role of heat shock protein (Hsp70). (Review). *Acta Physiol. Hungarica*, 2009/a 96, 149-155.
- Fábián, TK; Gótai, L; Beck, A; Fábián, G; Fejérdy, P. The role of molecular chaperones (HSPAs/HSP70s) in oral health and oral inflammatory diseases. A review. *Eur. J. Inflammation*, 2009/b 7, 53-61.
- Fábián, TK; Söti, Cs; Nguyen, MT; Csermely, P; Fejérdy, P. Expected functions of salivary HSP70 in the oral cavity. In: Berhardt LV editor. *Advances in Medicine and Biology*. Vol. 19. New York: Nova Science Publishers; 2011; 289-308.
- Fredericks, DC; Nepola, JV; Baker, JT; Abbott, J; Simon, B. Effects of pulsed electromagnetic fields on bone healing in a rabbit tibial osteotomy model. *J. Orthop. Trauma*, 2000 14, 93-100.
- Fredericks, DC; Piehl, DJ; Baker, JT; Abbott, J; Nepola, JV. Effects of pulsed electromagnetic field stimulation on distraction osteogenesis in the rabbit tibial leg lengthening model. *J. Pediatr. Orthop.*, 2003 23, 478-483.
- Gossling, HR; Bernstein, RA; Abbott, J. Treatment of ununited tibial fractures: a comparison of surgery and pulsed electromagnetic fields (PEMF). *Orthopedics*, 1992 15, 711-719.
- Goodman, R. and Blank, M. Magnetic field stress induces expression of hsp70. *Cell Stress and Chaperones*, 1998 3, 79-88.
- Goodman, EM; Greenbaum, B; Marron, MT. Altered protein synthesis in a cell-free system exposed to a sinusoidal magnetic field. *Biochim. Biophys. Acta*, 1993 1202, 107-112.
- Grace, KL; Revell, WJ; Brookes, M. The effects of pulsed electromagnetism on fresh fracture healing: osteochondral repair in the rat femoral groove. *Orthopedics*, 1998 21, 297-302.
- Grana, DR; Marcos, HJ; Kokubu, GA. Pulsed electromagnetic fields as adjuvant therapy in bone healing and peri-implant bone formation: an experimental study in rats. *Acta Odontol Latinoam*, 2008 21, 77-83.
- Gwechenberger, M; Rauscha, F; Stix, G; Schmid, G; Strametz-Juranek, J. Interference of programmed electromagnetic stimulation with pacemakers and automatic implantable cardioverter defibrillators. *Bioelectromagnetics*, 2006 27, 365-377.
- Hedén, P. and Pilla, AA. Effects of pulsed electromagnetic fields on postoperative pain: A double-blind randomized pilot study in breast augmentation patients. *Aesth Plast Surg.*, 2008 32, 660-666.

- Heredia-Rojas, JA; de la Fuente, AOR; González, JMA; Rodríguez-Flores, LE; Rodríguez-Padilla, C; Santoyo-Stephano, MA; Castaneda-Graza, E; Taméz-Guerra, RS. Effect of 60 Hz magnetic fields on the activation of hsp70 promoter in cultured INER-37 and RMA E7 cells. *In Vitro Cell. Dev. Biol. – Animal*, 2010 46, 758-763.
- Hopper, RA; VerHalen, JP; Tepper, O; Mehrara, BJ; Detch, R; Chang, EI; Baharestani, S; Simon, BJ; Gurtner, GC. Osteoblasts stimulated with pulsed electromagnetic fields increase HUVEC proliferation via a VEGF-A independent mechanism. *Bioelectromagnetics*, 2009 30, 189-197.
- Inoue, N; Ohnishi, I; Chen, D; Deitz, LW; Schwardt, JD; Chao, EY. Effect of pulsed electromagnetic fields (PEMF) on late-phase osteotomy gap healing in a canine tibial model. *J. Orthop. Res.*, 2002 20, 1106-1114.
- Jansen, JHW; van der Jagt, OP; Punt, BJ; Verhaar, JAN; van Leeuwen, JPTM; Weinans, H; Jahr, H. Stimulation of osteogenic differentiation in human osteoprogenitor cells by pulsed electromagnetic fields: an in vitro study. *BMC Musculoskeletal Disorders*, 2010 11, 188.
- Kanje, M; Rusovan, A; Siskin, B; Lundborg, G. Pretreatment of rats with pulsed electromagnetic fields enhances regeneration of the sciatic nerve. *Bioelectromagnetics*, 1993 14, 353-359.
- Kim, KT. Biochemical study of human periodontal ligament: preparation of cell attachment materials induced by pulsed electromagnetic fields. *Bull Kanagawa Dent Coll*, 1990 18, 89-98.
- Kold, SE; Hickman, J; Meisen, F. Preliminary study of quantitative aspects and the effect of pulsed electromagnetic field treatment on the incorporation of equine cancellous bone grafts. *Equine Vet. J.*, 1987 19, 120-124.
- Lednev, VV. Possible mechanism for influence of weak magnetic fields on biosystems. *Bioelectromagnetics*, 1991 12, 71-76.
- Leisner, S; Shahar, R; Aizenberg, I; Lichovsky, D; Levin-Harrus, T. The effect of short-duration, high-intensity electromagnetic pulses on fresh ulnar fractures in rats. *J. Vet. Med.*, 2002 49, 33-37.
- Liboff, AR; Rozek, RJ; Sherman, ML; McLeod, BR; Smith, SD. Ca^{2+} cyclotron resonance in human lymphocytes. *J. Bioelectricity*, 1987 6, 12-22.
- Lin, HY; Lin, YJ. In vitro effects of low frequency electromagnetic fields on osteoblast proliferation and maturation in an inflammatory environment. *Bioelectromagnetics*, 2011, in press, DOI: 10.1002/bem.2066.
- Lohmann, CH; Schwartz, Z; Liu, Y; Li, Z; Simon, BJ; Sylvia, VL; Dean, DD; Bonewald, LF; Donahue, HJ; Boyan, BD. Pulsed electromagnetic fields affect phenotype and connexin 43 protein expression in MLO-Y4 osteocyte-like cells and ROS 17/2.8 osteoblast like cells. *J. Orthop. Res.*, 2003 21, 326-334.
- Luben, RA; Cain, CD; Chen, MCY; Rosen, DM; Adey, WR. Effects of electromagnetic stimuli on bone and bone cells in vitro: Inhibition of responses to parathyroid hormone by low-energy low-frequency fields. *Proc. Natl. Acad. Sci. USA*, 1982 79, 4180-4184.
- Matsumoto, H; Ochi, M; Abiko, Y; Hirose, Y; Kaku, T; Sakaguchi, K. Pulsed electromagnetic fields promote bone formation around dental implants inserted into the femur of rabbits. *Clin. Oral Implant Res.*, 2000 11, 354-360.

- McClellan, JA; Boublikova, P; Palcek, E; Lilley, DMJ. Superhelical torsion in cellular DNA responds directly to environmental and genetic factors. *Proc. Natl. Acad. Sci. USA*, 1990 87, 8373-8377.
- Mentes, BB; Tascilar, O; Tatlicioglu, E; Bor, MV; Isman, F; Turkozkan, N; Celebi, M. Influence of pulsed electromagnetic fields on healing of experimental colonic anastomosis. *Dis. Colon Rectum*, 1996 39, 1031-1038.
- Midura, RJ; Ibiwoye, MO; Powell, KA; Sakai, Y; Doehring, T; Grabiner, MD; Patterson, TE; Zborowski, M; Wolfman, A. Pulsed electromagnetic field treatments enhance the healing of fibular osteotomies. *J. Orthop. Res.*, 2005 23, 1035-1046.
- Ortman, LF; Casey, DM; Deers, M. Bioelectric stimulation and residual ridge resorption. *J. Prosthet Dent*, 1992 67, 67-71.
- Ottani, V; Raspanti, M; Martini, D; Tretola, G; Ruggeri Jr, A; Franchi, M; Giuliani Piccari, G; Ruggeri, A. Electromagnetic stimulation on the bone growth using backscattered electron imaging. *Micron*, 2002 33, 121-125.
- Patino, O; Grana, D; Bolgiani, A; Prezzavento, G; Mino, J; Merlo, A; Benaim, F. Pulsed electromagnetic fields in experimental cutaneous wound healing in rats. *J. Burn Care Rehabil*, 1996 17, 528-531.
- Pienkowski, D; Pollack, SR; Brighton, CT; Griffith, NJ. Comparison of asymmetrical and symmetrical pulse waveforms in electromagnetic stimulation. *J. Orthop. Res.*, 1992 10, 247-255.
- Raji, AM. An experimental study of the effects of pulsed electromagnetic field (Diapulse) on nerve repair. *J. Hand Surg. Br.*, 1984 9, 105-112.
- Raji, AR. and Bowden, RE. Effects of high-peak pulsed electromagnetic field on the degeneration and regeneration of the common peroneal nerve in rats. *J. Bone Joint Surg. Br.*, 1983 65, 478-492.
- Rubin, CT; Donahue, HJ; Rubin, JE; McLeod, KJ. Optimization of electric field parameters for the control of bone remodeling: exploitation of an indigenous mechanism for the prevention of osteopenia. *J. Bone Miner Res.*, 1993 8, S573-S581.
- Satake, T; Yasu, N; Kakai, Y; Kawamura, T; Sato, T; Nakano, T; Amino, S; Ishiwata, Y; Saito, S. Effect of pulsed electromagnetic fields (PEMF) on human periodontal ligament in vitro. Alterations of intracellular Ca^{2+} . *Kanagawa Shikagu*, 1990 24, 735-742.
- Scardino, MS; Swaim, SF; Sartin, EA; Steiss, JE; Spano, JS; Hoffman, CE; Coolman, SL; Peppin, BL. Evaluation of treatment with a pulsed electromagnetic field on wound healing, clinicopathologic variables, and central nervous system activity of dogs. *Am. J. Vet. Res.*, 1998 59, 1177-1181.
- Schnoke, M. and Midura, RJ. Pulsed electromagnetic fields rapidly modulate intracellular signaling events in osteoblastic cells: Comparison to parathyroid hormone and insulin. *J. Orthop. Res.*, 2007 25, 933-940.
- Schwartz, Z; Fisher, M; Lohmann, CH; Simon, BJ; Boyan, BD. Osteoprotegerin (OPG) production by cells in the osteoblast lineage is regulated by pulsed electromagnetic fields in cultures grown on calcium phosphate substrates. *Annals Biomed Engineering*, 2009 37, 437-444.
- Showkatbakhsh, R; Jamilian, A; Showkatbakhsh, M. The effect of pulsed electromagnetic fields on the acceleration of tooth movement. *World J. Orthod.*, 2010 11, e52-e56.

- Solazzo, V; Palmieri, A; Pezzetti, F; Massari, L. Carinci, F. Effects of pulsed electromagnetic fields on human osteoblastlike cells (Mg-63). A pilot study. *Clin. Orthop. Relat. Res.*, 2010 468, 2260-2277.
- Spadaro, JA. and Bergstrom, WH. In vivo and in vitro effects of a pulsed electromagnetic field on net calcium flux in rat calvarial bone. *Calcif Tissue Int.*, 2002 70, 496-502.
- Stark, TM. and Sinclair, PM. Effect of pulsed electromagnetic fields on orthodontic tooth movement. *Am. J. Orthod. Dentofacial Orthop.*, 1987 91, 91-104.
- Strauch, B; Herman, C; Dabb, R; Ignarro, LJ; Pilla, AA. Evidence-based use of pulsed electromagnetic field therapy in clinical plastic surgery. *Aesthet Surg. J.*, 2009 29, 135-143.
- Takano-Yamamoto, T; Kawakami, M; Sakuda, M. Effect of a pulsing electromagnetic field on demineralized bone-matrix-induced bone formation in a bony defect in the premaxilla of rats. *J. Dent Res.*, 1992 71, 1920-1925.
- Tepper, OM; Callaghan, MJ; Chang, EI; Galiano, RD; Bhatt, KA; Baharestani, S; Gan, J; Simon, B; Hopper, RA; Levine, JP; Gurtner, GC. Electromagnetic fields increase in vitro and in vivo angiogenesis through endothelial release of FGF-2. *FASEB J.*, 2004 18, 1231-1233.
- Tsai, MT; Li, WJ; Tuan, RS; Chang, WH. Modulation of osteogenesis in human mesenchymal stem cells by specific pulsed electromagnetic field stimulation. *J. Orthop. Res.*, 2009 27, 1169-1174.
- Tuinstra, R; Greenbaum, B; Goodman, EM. Effects of magnetic fields on cell-free transcription in *E. Coli* and HeLa extracts. *Bioelectrochem. Bioenerg.*, 1997 43, 7-12.
- Varani, K; Gessi, S; Merighi, S; Iannota, V; Cattabriga, E; Pancaldi, C; Cadossi, R; Borea, PA. Alteration of A3 adenosine receptors in human neutrophils and low frequency electromagnetic fields. *Biochem. Pharmacol.*, 2003 66, 1897-1906.
- Wilmot, JJ; Chiego Jr., DJ; Carlson, DS; Hanks, CT; Moskwa, JJ. Autoradiographic study of the effects of pulsed electromagnetic fields on bone and cartilage growth in juvenile rats. *Arch. Oral. Biol.*, 1993 38, 67-74.
- Zienowicz, RJ; Thomas, BA; Kurtz, WH; Orgel, MG. A multivariate approach to the treatment of peripheral nerve transection injury: the role of electromagnetic field therapy. *Plast Reconstr. Surg.*, 1991 87, 122-129.

Index

A

- abrasion, 66, 67, 81, 82, 137
- abrasives, xii, 152
- abuse, 144, 173, 195, 196
- abutment, viii, ix, xiv, 3, 4, 7, 8, 9, 15, 16, 17, 20, 47, 52, 61, 62, 63, 64, 65, 67, 72, 75, 76, 77, 78, 79, 206, 219, 243, 248
- abutments, vii, ix, 1, 4, 5, 7, 17, 20, 21, 45, 46, 52, 64, 72, 76, 77, 78, 79, 80
- access, 131, 140
- accessibility, 157
- acetone, 118
- achilles tendon, 190
- acid, 98, 154
- acid etched, 154
- acquired immunodeficiency syndrome, 103
- acrylate, 80, 81, 118, 209, 212
- acrylic, 8, 9, 76, 78, 81, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 131, 132, 147, 153, 154, 160, 161
- acupuncture, xiii, 165, 166, 170, 171, 175, 194, 196
- acute leukemia, 106
- acute myeloid leukemia, 106
- acute stress, 183
- AD, 31, 162, 233
- ADA, 146
- adaptation, 46, 54, 55, 56, 57, 59, 93, 95, 96, 98, 137, 147, 201, 237
- additives, 121, 147
- adenosine, 250, 254
- adherence, x, 86, 90, 93, 128, 157, 161, 162
- adhesion, xii, 47, 48, 72, 105, 106, 116, 117, 119, 121, 122, 126, 128, 145, 146, 152, 153, 154, 155, 156, 157, 160, 161, 162, 211
- adhesive materials, 146, 147, 148
- adhesive strength, 54, 84, 117, 118, 119
- adhesives, xi, 143, 144, 145, 146, 147, 148, 149, 153
- adjustment, xii, 72, 95, 131, 143, 145, 159, 170, 196, 230
- adolescents, 23, 106, 130, 239
- adults, xi, 10, 12, 105, 106, 129, 130, 138, 139, 141, 159, 161, 187, 196, 203, 228, 229, 240
- adverse effects, 119, 122, 173, 175, 232, 236
- adverse event, 186
- aesthetic, ix, 26, 46, 61, 62, 71, 75, 76, 79, 81, 82, 83, 130, 131
- aesthetic character, 62
- aesthetics, ix, xi, 3, 21, 23, 47, 61, 62, 67, 81, 82, 115, 218
- affective disorder, 172, 197
- African Americans, 185
- after-care, 22
- age, 29, 41, 94, 98, 139, 236
- aggression, 119, 203
- aggressiveness, 202, 207
- aging population, x, 85, 130
- agonist, 185
- AIDS, 86, 88, 89, 90, 91, 92, 105, 156, 160, 185
- algorithm, 108
- all ceramic, ix, 45, 46, 48, 53, 56
- all-ceramic restorations, 53
- allergens, 146, 209
- allergic, xi, 116, 166, 174, 177, 191, 209, 236
- allergic reaction, 209
- allergy, x, 85, 230, 232
- alloys, ix, 62, 66, 67, 69, 72, 77, 82, 83, 209
- altered state of consciousness, 176, 177, 178, 179
- alternative medicine, xiii, 165, 166, 183, 223, 227
- aluminum oxide, 81
- alveolar ridge, 64, 80, 146
- amalgam, 209, 214, 236
- Amalgam, 214
- amelogenesis imperfecta, 21, 26
- ammonia, 97
- ammonium, 121

- amplitude, 37, 247
 amylase, 178
 analgesic, 172
 analgesic effect, 172
 anamnesis, 168, 249
 anastomosis, 192, 253
 anchor, 68, 72, 181
 anchorage, 8, 9, 17, 18, 19, 20, 21, 23, 24, 25, 138, 140
 anemia, 90, 91, 92
 anesthetic, 96
 anger, 178, 182
 angiogenesis, 170, 195, 246, 247, 254
 angle, 8, 64, 66, 67, 69, 70, 71, 72, 117
 antagonist teeth, 207
 anterior cingulate cortex, 182
 antibiotic, 154
 antibodies, 94, 210
 antidepressant, 173, 184
 antidepressants, 171, 172
 antidepressive, 172, 182
 antifungal, 87, 104, 105, 122, 153, 155, 156, 157, 159, 163
 antigen, 209
 anti-inflammatory drugs, 171
 antimicrobial agents, xi, 116, 121, 122, 128
 antimicrobials, 121
 anti-nociceptive effects, 178
 antioxidant, 174
 antioxidant activity, 174
 antipsychotic, 110
 anxiety, 131, 166, 172, 174, 177, 178, 180, 181, 182, 188, 192, 195, 196, 200, 204, 205, 207, 208, 211, 216, 237
 anxiety disorder, 216
 anxiolytics, 171, 172, 221
 apex, 20
 aphthous lesions, 96
 apical, 18, 50
 apico-alveolar, 213
 aplastic anemia, 92
 apnea, 181
 appointments, 99
 approximal, ix, 16, 18, 19, 62, 75, 76, 77
 arousal, 176, 178, 207
 ARS, 112
 arthroplasties, 153
 arthroplasty, 162
 articulator, 64
 articulators, 76
 artificial gingiva, 208
 artificial teeth, x, xi, 8, 29, 59, 62, 76, 78, 79, 115, 116, 118, 119
 ascorbic acid, 95
 aspergillosis, x, 86
 aspiration, xii, 151, 152, 158
 aspiration pneumonia, 158
 aspirations, 99
 assault, 203
 assessment, 64, 91, 110, 138, 152, 157, 184, 218
 asthma, 190
 asthmatic patient, 180
 asymmetry, 247
 asymptomatic, 91
 atmosphere, 81
 atmospheric pressure, 146
 atoms, 68
 atrophy, 91, 134
 attached soft tissue, 20
 attachment, 8, 9, 19, 62, 135, 189, 248, 252
 attachments, 132, 135
 attention, xii, 2, 40, 43, 72, 80, 92, 93, 97, 99, 151, 155, 176, 207, 217, 218, 221
 attitudes, 24, 230
 atypical facial pain, 173, 177, 205
 atypical odontalgia, 173, 206
 authoritarianism, 234
 authority, 220
 autoantibodies, 94, 210
 autoimmune, x, 86, 88, 93, 94, 96, 102, 236
 autoimmune disease, x, 86, 88, 93, 102, 236
 autonomic, 176, 177, 179, 180, 204, 213
 autonomic activity, 204
 autonomic nervous system, 180
 autonomy, 167
 autopolymerized resins, 119, 126
 avoidance, 16, 209, 214, 217, 223, 224, 225
 awareness, viii, 27, 28, 39, 40, 41, 169
 axial, 64, 65, 75, 76
 axis, vii, 1, 2, 5, 6, 8, 9, 16, 52, 58, 64, 70, 75, 182, 187, 210

B

- back pain, 240
 bacteria, 54, 87, 158, 161, 178, 210
 bacterial infection, 95
 balanced occlusion, 77
 bar, 8, 9, 19, 135, 141
 barriers, 87
 bars, 9
 basal ganglia, 206
 base, x, xii, 10, 29, 66, 67, 78, 92, 97, 101, 104, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 128, 147, 151, 161, 162, 163, 201, 203, 208, 229, 240

- basement membrane, 94
baseplate, vii, 1, 2, 5, 6, 7, 8, 9, 212
BD, 23, 26, 111, 252, 253
behavior, 29, 40, 43, 57, 166, 167, 176, 185, 189, 195, 208, 221, 222, 229
behavior therapy, 167, 221
behavioral change, 211
behaviors, 43, 109, 207
bending, 48, 53, 56
benefits, vii, 54, 81, 117, 121, 135, 136, 147, 157, 166
benzene, 145
benzodiazepines, 172, 221
beveling, 54
bilateral balanced occlusion, 208
bioavailability, 101
biocompatibility, 66, 78, 119, 127
Biocompatibility, 119, 125, 126, 127
biofeedback, xiii, 22, 165, 166, 176, 177, 179, 185, 190, 192, 194, 221, 225
biofeedback training, 192
biofilms, 87, 103, 105, 113, 153, 158, 160, 163, 164
biological responses, 244
biological rhythms, 183
biological systems, 249, 250
biomaterials, xii, 83, 152, 154, 163
biomechanics, 21
biopsy, 91, 94
biopsychosocial model, 214, 215
birth control, 130
bisphosphonates, x, 86, 92, 101, 112
bite, 25, 28, 33, 53, 57, 131, 134, 135, 185, 186, 192, 196, 207, 209, 234, 236, 237
black cohosh, 173, 174, 193
black tea, 175
bleeding, 86, 92, 97, 98, 101, 212
blepharospasm, 180
blood, 30, 119, 157, 168, 169, 174, 175, 195, 204, 211
blood flow, 169, 174
blood pressure, 169, 174, 195
blood vessels, 119, 175, 204
bloodstream, 104
body exercises, 177, 182
body image, 13, 176, 203
body temperature, 213
bond, xi, 47, 48, 54, 55, 115, 118, 119, 124, 125, 145
bonding, 46, 48, 54, 55, 56, 65, 66, 76, 84, 117, 118
bonds, 53, 145
bone, x, xi, xiii, 18, 19, 20, 75, 77, 86, 88, 92, 97, 101, 102, 108, 130, 131, 132, 133, 134, 137, 139, 140, 146, 147, 161, 170, 171, 173, 175, 189, 191, 206, 211, 232, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254
bone cells, 252
bone defects, 170, 171, 247
bone density, 20
bone deposition, 170
bone form, xiii, 170, 191, 243, 246, 247, 251, 252, 254
bone formation, xiii, 170, 191, 243, 246, 247, 251, 252, 254
bone fractures, 170, 247
bone graft, 247, 252
bone growth, 250, 253
bone healing, xiii, 243, 247, 250, 251
bone loss, xi, 130, 132, 133, 134, 137, 139, 175
bone marrow, 88, 147, 161, 244, 245, 250
bone marrow transplant, 88, 161
bone remodeling, xiii, 97, 171, 189, 243, 245, 248, 253
bone resorption, 77, 131, 133, 137, 140, 232, 245, 251
bones, 208
border seal, 208
brackets, 17, 18, 20
bradykinesia, 98
brain, vii, 28, 29, 41, 43, 168, 169, 174, 178, 182, 183, 185, 188, 193, 203, 206, 210, 229, 249
brain functions, 28, 41
brain stem, 203, 210
brainstem, 205, 206, 213, 239
Brazil, xi, 85, 115, 129, 130, 139, 140
break, 76
breakage, 54
breast augmentation, 251
breast cancer, 173
breathing, xiii, 165, 166, 176, 179, 181, 182, 189
brief dynamic psychotherapy, 168
brittleness, 82
bruxism, 10, 98, 176, 177, 184, 192, 208, 229, 239
budding, 157
burning mouth, x, 85, 172, 173, 191, 206, 239

C

- Ca²⁺, 244, 246, 248, 252, 253
CAD, 47, 67, 84
CAD/CAM, 47, 67, 84
caffeine, 174, 175, 194, 195
calcium, 97, 107, 145, 246, 253, 254
calcium channel blocker, 107
calculus, 211
CAM, 47, 67, 84, 166, 180, 182, 183, 222, 223, 225, 227, 228

- canals, 80
cancer, 88, 97, 103, 110, 111, 112, 155, 159, 160, 161, 187, 197, 232
Candida, xii, 87, 88, 89, 91, 100, 103, 104, 105, 106, 111, 112, 121, 122, 126, 127, 128, 145, 146, 151, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 163, 164
candidiasis, 87, 88, 89, 90, 91, 92, 93, 95, 96, 97, 100, 103, 104, 105, 106, 108, 154, 155, 156, 162, 163, 164
cannabinoids, 186
cannabis, 174
cap, 65
carbamazepine, 99
carbon, xi, 54, 115
carbon fiber, xi, 115
carboxyl, 145
carboxymethyl cellulose, 145
carcinogenic, 66
carcinoma, 108, 110
cardiac pacemaker, 169
cardiac problems, 180, 181, 182
cardiovascular disease, 103
caries, xi, 77, 78, 129, 130, 141, 211
carious, 18, 19, 46, 50, 58, 210, 212
cartilage, 254
case study, 108
casting, 69, 72, 73, 75, 76, 80
castings, x, 62, 66, 67
categorization, 64
catheter, 127
cation, 145
causal relationship, 40
cavity preparation, 54
cell culture, 246
cell death, 126
cell metabolism, 170
cell surface, 161
cellulose, 144
cementation, 4, 54, 78, 79
cemented, 19, 62, 65, 132
central nervous system, 169, 174, 178, 194, 204, 205, 253
ceramic, ix, 25, 26, 45, 46, 47, 48, 50, 53, 54, 55, 56, 57, 58, 59, 62, 65, 67, 79, 81, 82, 83
ceramic materials, ix, 45, 46, 48, 53, 58, 81
Ceramics, 47
cerebellum, 206
cerebral blood flow, 42
cerebral cortex, 42
cerebrospinal fluid, 162
chair-side, 48
challenges, 82, 144, 148, 194, 238
chaperokine, 178
chaperones, 232, 251
chastetree, 174
cheilitis, 154
chemical, 4, 48, 90, 118, 120, 124, 127, 145, 212, 232, 236
chemical properties, 145
chemokines, 209
chemotherapy, x, 86, 88, 90, 92, 93, 100, 101, 106, 110, 111, 112
chewing, viii, 3, 27, 28, 31, 32, 33, 34, 37, 39, 40, 41, 43, 53, 58, 62, 65, 70, 96, 117, 130, 131, 132, 146, 147, 148, 207, 212, 218
Chicago, 83
childhood, 202, 222
children, xi, 129, 130, 162
chlorhexidine, 89, 121, 157, 158, 160, 161, 162
cholinesterase, 110
cholinesterase inhibitors, 110
chronic, x, xi, 64, 86, 88, 93, 96, 97, 100, 102, 129, 130, 143, 154, 162, 171, 174, 176, 177, 179, 182, 184, 189, 191, 196, 205, 210, 225, 227, 230, 239, 240
chronic diseases, 86
chronic fatigue, 174, 182, 184, 191
chronic fatigue syndrome, 191
chronic illness, xi, 129, 130, 182
chronic renal failure, 97
chronification, 205, 224
circulation, 168, 169
circumference, 68, 156
clasp, 8, 29, 84
clasps, 8, 62, 75
classes, vii, 1, 5, 7, 72, 73
classification, vii, 1, 2, 5, 6, 7, 10, 11, 74, 155, 241
cleaning, 89, 90, 93, 94, 95, 98, 105, 127, 156, 158, 159
cleft lip, 21, 25, 26
clenching, 57, 208
clients, 167
clinical application, 56, 62
clinical crown, 18, 79
clinical examination, 64, 214
clinical presentation, 154
clinical symptoms, 93, 170
clinical trials, 136, 191
closure, 4, 26, 80, 113, 176
clustering, 238
CMC, 145
coagulation, 97, 157
cobalt, 186, 250
cocoa, 174
coding, 43

- coefficient of friction, 66, 69, 74, 81
coffee, 174, 194
cognition, 29, 40, 43, 176, 193
cognitive, viii, 27, 28, 40, 41, 43, 99, 167, 174, 175, 176, 181, 185, 189, 195, 196, 211, 221, 240
cognitive ability, ix, 28
cognitive function, viii, 27, 28, 40, 41, 174, 175
cognitive impairment, 28
cognitive perspective, 189
cognitive process, 40, 43
cognitive processing, 43
cognitive therapy, 167
cognitive-behavioral therapy, 167, 181, 196, 240
cohesion, 72, 145, 146
cohesive strength, 48, 145
collaboration, viii, 15, 16, 21, 23, 226
collagen, 170, 244
colonization, xii, 87, 88, 100, 104, 105, 106, 111, 112, 151, 152, 153, 154, 159, 162, 163
color, 82, 118, 123
combination therapy, 109
comfort, viii, ix, 27, 28, 61, 62, 67, 72, 75, 77, 134, 144, 183, 211, 216, 227, 235
commercial, 117, 125
commissure, 96
communication, xiii, 167, 199, 200, 202, 217, 220, 221, 227, 244
communication skills, 200
communities, 240
community, 41, 107, 195
compatibility, 119, 120
compensation, 65, 76
complete blood count, 92
complexity, 42
compliance, 3, 19, 22, 219
complications, vii, xiii, 3, 10, 11, 19, 64, 78, 81, 86, 87, 92, 94, 95, 97, 98, 99, 100, 101, 102, 106, 135, 170, 199, 219
composite, ix, xi, 17, 45, 46, 47, 48, 53, 54, 56, 57, 58, 59, 74, 81, 82, 115, 123, 124
composite resin, 57, 58, 59, 124
composites, 46, 47, 56, 57, 74, 76, 82, 123, 209
composition, 47, 87, 93, 144, 239
compost, 121
compounds, xi, 115, 126, 212
compression, 20, 92, 98, 189, 248
computer, 32
concentration, viii, 27, 34, 76, 87, 93, 96, 117, 157, 174, 179
conception, 175
conditioning, 185, 209
conductance, 179
conduction, 77, 189
configuration, 59, 71, 249
congruence, 167, 220
connector, 48, 53, 54, 56, 59
connectors, ix, 45, 48, 55
conscious awareness, 40, 41
conscious sedation, 221
consciousness, 29, 43, 176, 177, 178, 179, 186, 187, 188, 193, 231
consensus, 107, 117, 176, 202
consolidation, 247
consonants, 213
constituents, 173, 174, 211
construction, ix, 46, 47, 53, 54, 61, 64, 68, 69, 70, 71, 72, 73, 74, 82, 92, 116
consumption, 174
contact dermatitis, 119
contaminant, 91
contamination, xi, 46, 100, 116, 121, 122, 164
contour, 118, 208
contraceptives, 174
control group, 98, 122, 154
controlled trials, 139, 191
controversial, 96, 97, 154
controversies, xi, 143
conus, 62, 64, 65, 66, 67, 68, 70, 71, 72, 73, 76, 78, 83, 84
convergence, 55
conversion, 116, 120, 121, 126
cooling, 66
cooperation, 99
coordination, 201
copolymer, 145
copper, 147
correlation, xi, xii, 58, 127, 129, 130, 151, 152, 163, 164, 201
correlations, 40
corrosion, 53, 66, 67, 71
cortex, viii, 27, 29, 33, 34, 40, 42, 193, 196, 210, 247
corticosteroids, x, 86, 88, 95, 96
cortisol, 180, 182, 195
cosmetic, 25
cost, viii, ix, xi, 15, 23, 45, 46, 100, 118, 129, 131, 132, 135, 136, 137, 141, 211
costs, xi, 66, 84, 129, 130, 135, 136, 137, 138
counseling, 231
covalent bond, 145
coverage, ix, 19, 20, 45, 46, 55, 56, 58, 77, 121, 144, 170, 246
cracks, 48
cranial nerve, 207
cranial nerves, 207
CRF, 97
criticism, 65

cross reactivity, 210
 crossbite, 21
 cross-sectional study, 160
 cross-validation, 235
 crown, ix, 4, 16, 17, 18, 19, 21, 26, 54, 55, 56, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 79, 80, 81, 82, 83, 84
 crowns, ix, 4, 8, 9, 10, 16, 20, 26, 54, 58, 59, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84
 crystal growth, 211
 crystalline, 121
 crystals, 68
 CSF, 244
 CT, 193, 253, 254
 culture, 87, 125, 156, 246
 cure, xi, 129, 130, 181, 227
 curing cycle, 125, 126
 cusps, 8
 cusps, 47
 CV, 228
 cycles, 53, 68, 119, 120, 158
 cycling, 47, 53, 58, 59
 cyclosporine, 97
 cysts, 210
 cytochrome, 174, 175
 cytokines, 209, 245
 cytology, 91
 cytotoxicity, 119, 120, 125, 126

D

daily living, 40, 235
 damages, 78
 danger, 170, 180
 data analysis, 40
 DCA, 149
 dead spaces, 76
 decay, 26
 decisions-making, 220
 decomposition, 212
 decontamination, 159
 defects, 25, 66, 170, 171, 247
 defense mechanisms, 209, 231, 251
 deficiencies, 62
 deficiency, 97, 120, 136, 179
 deformation, 68, 69, 70, 71, 73, 76, 79, 80, 117, 123
 degradation, 119, 244, 245, 246
 delamination, 47, 48
 delayed type dental support, viii, 2, 6, 9
 demarcation line, x, 62, 64, 65, 79
 dementia, 28, 40, 41, 43, 98, 99, 109, 174, 179
 demineralization, 244

demographic change, xi, 129
 dendritic cell, 175
 dendritic cells, 175
 dental alloys, ix, 45, 46, 213, 232
 dental care, viii, xiii, 10, 12, 16, 28, 140, 147, 161, 165, 166, 167, 168, 183, 226
 dental caries, 130, 141, 211
 dental ceramics, 53, 58
 dental clinics, ix, 45, 46, 215
 dental fear, 217, 220, 227, 229, 231, 237
 dental floss, 17
 dental implants, xiii, 3, 13, 20, 25, 26, 62, 103, 109, 140, 141, 168, 170, 191, 206, 219, 243, 246, 250, 252
 dental phobia, 176, 188
 dental plaque, xii, 66, 76, 151, 164
 dental pulp, 203, 206, 210, 234, 237
 dental restorations, 13, 57, 77, 203, 239
 dental support, 7, 9
 dentin, 46, 54, 58, 59
 dentist, viii, xii, xiii, 3, 11, 12, 15, 23, 86, 88, 90, 92, 143, 147, 153, 158, 165, 167, 171, 172, 173, 183, 199, 200, 201, 202, 215, 217, 219, 220, 221, 223, 224, 225, 226, 227, 228, 232, 233, 234, 240
 dento-musocal support, 7, 8, 9
 denture adhesives, xi, 143, 144, 145, 146, 147, 148, 149
 denture base, x, xii, 29, 78, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 128, 147, 151, 161, 162, 208
 denture bearing areas, 144
 denture intolerance, xiii, 11, 24, 165, 166, 177, 178, 184, 186, 187, 199, 200, 201, 202, 213, 214, 216, 223, 225, 227, 228, 231, 237, 251
 denture satisfaction, 28, 141
 deposition, 94, 120, 170, 246
 depression, 153, 172, 174, 175, 176, 177, 178, 180, 181, 182, 184, 191, 195, 196, 201, 204, 205, 207, 211, 229, 248
 depressive symptoms, 174
 deprivation, 182
 depth, 50, 80, 167, 195, 231
 dermatitis, 119, 210
 dermatome, 204
 dermatomes, 204
 desensitization, 192
 destruction, 211
 desynchronization, 178
 detectable, xii, 152, 213
 detection, xii, 102, 105, 152, 155, 159
 developing countries, 130, 137
 diabetes, 86, 88, 93, 106, 164
 diabetic patients, 93, 155

dialysis, 97, 128, 153, 160
 diameter, 10, 18, 20, 64, 68, 69, 70, 72
 diastema, 17
 diet, 222, 223, 225, 227, 228
 differential diagnosis, 91, 214, 216, 227
 digestion, 98
 dimethacrylate, 118
 diode laser, 195
 disability, 41
 discomfort, 2, 29, 95, 131, 166, 169, 170, 207, 218
 discrimination, 43
 diseases, x, xi, 85, 86, 88, 90, 91, 92, 93, 97, 101, 102, 127, 129, 130, 144, 158, 161, 168, 173, 180, 211, 216, 217, 232, 249
 disinfection, 90, 92, 157, 158, 159, 161, 162, 163
 disorder, 26, 43, 97, 98, 99, 109, 175, 186, 187, 196, 213, 214, 220, 240
 displacement, 70, 120
 dissatisfaction, xi, 129, 131
 dissatisfied, 132
 dissociation, 176
 distal end, 7, 20
 distalized, 21
 distilled water, 158
 distraction, 221, 247, 251
 distress, 130, 178, 205
 distributed load, 9
 distribution, 42, 47, 48, 164
 diuresis, 174
 divergence, 216
 diversity, 210
 dizziness, 213
 DNA, 87, 155, 250, 253
 doctors, 171
 dogs, 194, 253
 DOI, 195, 232, 238, 249, 252
 dopamine, 98
 dopaminergic, 98, 174, 182
 dorsal, 30, 33, 34, 40, 43, 182, 187
 dosage, 101, 172
 dosing, 101
 double bonds, 54
 double crown, ix, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 74, 75, 77, 78, 79, 80, 81, 82, 83, 84
 double crowns, ix, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 74, 75, 77, 78, 79, 81, 82
 drifting, 16
 DRS, xii, 152, 153, 155, 156, 157, 158, 159
 drug interaction, 110, 174, 175
 drug metabolism, 174, 192
 drug resistance, 103, 104, 157
 drugs, x, 85, 86, 87, 88, 97, 98, 99, 101, 107, 144, 155, 157, 171, 172, 173, 174, 177, 212

dry mouth, xii, 96, 143, 211
 drying, 175
 durability, ix, 47, 55, 61, 62, 67, 78
 DVD-glasses, 221
 dynamic-static friction, 68, 71
 dysaesthesia, 213, 233
 dysgeusia, 97
 dyskinesia, 98, 99
 dysphagia, 96, 97, 98, 99

E

early contact, 4, 169, 208
 early contacts, 4, 208
 eating, 28, 43, 130, 131, 145, 203
 eccentric positions, 77
 ecology, xiii, 152, 159
 economic, xi, 12, 106, 111, 130, 132, 136, 204
 ecosystem, 232
 ectodermal dysplasia, 21, 26
 ectodermic, 203
 ectomesodermic, 203
 ectopic eruption, 21
 eczemas, 210
 edema, 247
 edentulism, 86, 130, 134, 137, 138, 237
 edentulous area, 4, 7, 9, 20
 edentulous patients, viii, 2, 5, 10, 11, 27, 29, 62, 75, 78, 84, 94, 98, 103, 131, 134, 136, 138, 141, 153, 154, 208, 230
 edentulous ridge, 9, 18, 23
 edentulous space, vii, 1, 5, 8, 23, 46
 edentulousness, 22, 25, 62, 203, 228
 editors, 11, 12, 13, 24, 184, 185, 186, 187, 188, 189, 190, 191, 192, 194, 195, 229, 230, 231, 232, 233, 234, 239, 240, 241, 251
 education, 98, 222, 224
 EEG, 30, 178, 184, 197
 efflux mechanisms, 104
 Egypt, 45
 elastic, 17, 19, 20, 21, 26, 48, 65, 68, 69, 70, 71, 80, 95, 117, 121, 124
 elastic deformation, 68, 69, 70, 71, 80
 elasticity modulus, 46
 elastomer, 79
 elastomers, 79
 elderly, viii, xi, xii, 11, 27, 28, 40, 41, 64, 84, 94, 98, 109, 120, 127, 129, 130, 139, 140, 151, 152, 153, 156, 158, 159, 160, 161, 162, 163, 164, 229, 233, 234, 237, 238
 elderly population, xi, 129, 130, 161, 162
 electric field, 244, 253
 electric fields, 244

- electrodes, 31, 169
 electromagnetic, 22, 24, 168, 170, 175, 184, 186, 188, 189, 190, 191, 192, 193, 194, 195, 197, 249, 250, 251, 252, 253, 254
 electromagnetic field therapy, 184, 197, 254
 electromagnetic fields, 186, 188, 189, 190, 191, 192, 193, 195, 249, 250, 251, 252, 253, 254
 electromagnetism, 251
 electron, 57, 108, 253
 electron microscopy, 108
 elevators, 207
 elongated, 19, 70, 155
 elongation, 170
 e-mail, 1, 15, 27, 45, 61, 85, 115, 129, 143, 151, 165, 199, 243
 EMG, viii, 27, 28, 29, 30, 31, 32, 34, 35, 36, 37, 40, 41, 179, 190
 emotion, 186
 emotional, 132, 174, 178, 180, 184, 201, 202, 203, 204, 207, 208, 210, 213, 216, 220, 222, 223, 229, 232, 239
 emotional conflict, 208
 emotional disorder, 184
 emotional experience, 204
 emotional problems, 132, 174
 emotional processes, 204, 207
 emotional reactions, 180, 220
 emotional state, 204, 222
 emotions, 176, 207, 217, 221, 222, 223
 empathy, 167, 220
 emulsifying, 144
 enamel, 50, 54
 encounters, 209
 endocrine, x, 86, 176
 endorphins, 169
 endosseus, 4, 219
 endothelial, 195, 210, 246, 254
 endothelial cells, 210
 end-stage renal disease, 108
 endurance, 53, 58
 energy, 112, 121, 124, 154, 158, 163, 168, 169, 175, 177, 181, 182, 187, 189, 244, 245, 247, 248, 252
 enlargement, 20, 97
 entrapment, 212
 environment, 53, 116, 119, 195, 202, 211, 240, 246, 252
 enzymes, 175, 244
 epidemiologic, 140
 epidemiology, 103, 104, 152, 163, 235
 epilepsy, 180, 184, 193
 epileptic, 180
 epinephrine, 206
 episodic memory, 41
 epithelial, x, 86, 93, 100, 106, 154
 epithelial cells, 93, 106, 154
 epithelium, 94, 210
 Epstein-Barr virus, 97
 equilibrated splints, 170
 equipment, 80, 179
 erosion, 119
 eruption, 18, 19, 23, 24, 25, 26
 erythema, 91, 119, 155
 erythematous, 91, 108, 155
 esthetic complaints, 211
 esthetic restoration, 18, 46
 esthetical complaints, 211
 etiology, 96, 103, 108, 160, 161, 162, 202, 236
 EU, 1, 15, 126, 165, 199, 243
 Europe, 13
 evidence, 10, 48, 120, 133, 134, 138, 158, 168, 207, 214, 226, 229, 249
 evolution, 56, 86, 94, 98
 examinations, 29, 30, 78, 146, 153, 214, 215, 226
 excessive strength, 48
 excitability, 170
 exclusion, 205, 216
 execution, 21, 40, 42
 exercise, 68, 185
 expansion fluid, 80
 expectation, 131, 139, 141
 expectations, 3, 12, 21, 136, 180, 201, 218, 236
 expertise, 145
 exploitation, 71, 253
 exposure, 16, 76, 102, 113, 158, 183, 245, 246
 exposure of the pulp, 16
 extension, 9, 47, 79
 extensor, 28
 extracellular matrix, 246
 extraction, xiii, 4, 10, 12, 13, 18, 133, 170, 219, 243, 246
 extracts, 173
 extrusion, viii, 15, 16, 18, 19, 25, 26

F

- fabrication, ix, 16, 45, 46, 74, 144
 facial muscles, 206
 facial pain, 173, 177, 205, 229, 230, 235, 236, 240
 facial palsy, 213
 facial tic, 213
 faith, 186
 family members, 216
 fantasy, 176
 fasting, xiii, 165, 167, 191
 fatigue, 53, 57, 58, 69, 169, 173, 174, 175, 177, 179, 182, 183, 184, 191, 205, 207, 213

fear, 176, 200, 217, 220, 221, 227, 229, 231, 235, 237
 feelings, 167, 203, 207, 220
 femur, 191, 252
 fever, 213, 238
 fiber, ix, xi, 45, 46, 47, 48, 53, 54, 56, 57, 115, 117, 123, 124
 fiber content, 47, 53
 fibers, 18, 19, 47, 48, 54, 58, 75, 117, 118, 123, 205, 248
 fibroblasts, xiv, 189, 243, 248
 fibromyalgia, 191, 196
 fibrosis, 96
 filler, 48, 54, 82, 145
 fillers, 54, 80, 82
 filling materials, 127
 fillings, 46, 214, 236
 film thickness, 55
 financial, 3, 81, 86, 134, 135, 218, 219
 fine friction, 70
 finish line, 46
 finishing lines, 54
 Finland, 153
 fireproof mass, 80
 first molar, 53
 fit, xii, 30, 74, 76, 90, 143, 145, 216
 fitting, 67, 146, 147, 152, 159, 208
 fixation, 19, 132, 216
 fixed denture, vii, 1, 2, 3, 4, 5, 6, 7, 132, 200, 201, 219
 flash light stimuli, 178
 flavonoids, 175
 flavor, 212
 flavour, 234
 flaws, 48, 53
 flex, 53
 flexibility, 49, 182
 flexing, 53
 flexural, x, 47, 59, 115, 116, 117, 118, 121, 122, 123, 124, 126, 128
 flora, 154, 159
 flow rate, 108, 142, 179, 240
 fluctuations, 98, 109
 fluid, 80
 fluoroalkyl methacrylates, 128
 fMRI, 41, 43
 food, 31, 99, 141, 142, 145, 146, 183, 203, 212, 235
 foramen, 132, 135
 force, vii, ix, 1, 5, 7, 8, 9, 17, 20, 21, 28, 33, 34, 38, 39, 40, 48, 52, 53, 57, 59, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 79, 82, 83, 84, 124, 131, 134, 135, 139, 145, 146, 148, 203
 forced eruption, 18, 19

forecasting, 103
 formaldehyde, 119, 125
 formation, x, xii, 19, 62, 87, 151, 152, 153, 154, 163, 175, 197, 206, 210, 211, 235, 246, 247, 248, 250
 formula, 33
 fracture, ix, 45, 46, 47, 48, 53, 54, 56, 57, 58, 59, 78, 92, 116, 117, 118, 123, 251
 fracture resistance, ix, 45, 47, 48, 53, 56, 58
 fracture toughness, 58, 123
 fractures, 24, 54, 77, 116, 117, 118, 122, 170, 188, 247, 251, 252
 fragility, 100
 fragments, 96
 framework, ix, 5, 7, 8, 47, 48, 53, 56, 74, 75, 76, 82, 93
 free association, 167
 free associations, 167
 free energy, 154, 163
 free gingiva, ix, 62, 75, 76, 79, 82
 free way space, 207
 Freud, 168, 187
 friction, 65, 66, 67, 68, 69, 70, 71, 74, 78, 81, 168
 frontal region, 7, 22
 functional impression, 8, 9
 fungal, xii, 87, 90, 93, 97, 99, 103, 151, 152, 154, 155, 156, 159, 160, 161, 163
 fungal infection, xii, 87, 90, 97, 99, 103, 152, 155, 163
 fungi, x, 86, 87, 103, 158

G

GABA, 212
 gagging, 209, 229, 232
 gain from illness, 201
 galvanic, 67
 gap, 10, 54, 56, 64, 68, 69, 70, 71, 72, 73, 80, 189, 229, 244, 252
 gastrointestinal tract, 209
 gate-control theory, 169
 gel, 89, 93, 97, 144, 157, 161
 gender differences, 201, 238
 gene activation, 244
 general anaesthesia, 221
 genes, 205, 246
 genetic factors, 253
 genioglossus activity, 209
 genital, 96
 genomics, 232
 genus, 87, 88
 genus *Candida*, 87, 88
 geometrical parameters, 71
 geometry, 54

Germany, 73, 84, 158
 gestures, 182
 ginger, 191
 gingival, x, 4, 23, 46, 52, 54, 62, 64, 65, 68, 69, 71, 72, 79, 80, 97, 107, 108, 211, 212
 gingival crevice, 54
 gingival edge, 71, 79
 gingival overgrowth, 107, 108
 gingivitis, 161, 211
 ginseng, 174, 196, 197
 gland, 112, 179
 glass fiber, xi, 54, 57, 58, 115, 117, 118, 123, 124
 glass fibers, 54, 58, 117, 118, 123
 glass-ceramic, 47
 glasses, 221
 glaucoma, 180
 glaze, 122
 glucose, 93, 174
 glutamate, 212
 gnathodynamometry, 146
 goal setting, 222
 God, 180
 gold, ix, xii, 62, 66, 67, 69, 77, 82, 83, 152, 155
 graduate education, 224
 granulation tissues, 210
 granulomatous reactions, 210
 gravity, 77
 Great Britain, 147
 grinding, ix, 61, 62, 64, 81, 208
 grooves, 49, 52, 53, 65
 group therapy, 192
 group-therapy, 179
 growth, 26, 127, 146, 168, 169, 178, 249, 254
 guidance, 65, 206, 208
 guided tissue regeneration., 19
 guidelines, 111, 118, 158, 160
 guilt, 203
 gum, 32, 43, 96, 144

H

habituation, 178
 hairy leukoplakia, 97, 108
 halitosis, 211, 212, 237, 240, 241
 hallucinations, 195
 haptens, 209, 236
 hard tissues, 155
 hardness, 59, 66, 68, 81, 120, 121
 harmful effects, 133, 146
 harmonization, 82
 hay fever, 190
 hazards, 147, 171, 173, 195
 HE, 108, 110, 185, 195

head and neck cancer, 110, 111, 112, 163
 headache, 189, 190, 213
 Head-zones, 205
 healing, xiii, 93, 130, 133, 167, 170, 171, 183, 186, 189, 191, 192, 195, 243, 246, 247, 250, 251, 252, 253
 health, ix, xi, 10, 12, 28, 62, 67, 70, 75, 86, 94, 103, 109, 129, 130, 131, 137, 139, 140, 146, 147, 166, 168, 173, 180, 181, 182, 188, 190, 191, 192, 193, 195, 201, 206, 210, 213, 214, 215, 216, 220, 222, 223, 226, 229, 231, 232, 239, 249
 health care, 86, 130, 131, 166
 health care system, 130, 131, 166
 health condition, 28, 137, 139, 173, 180
 health effects, 214
 health problems, 215, 223
 health status, 214, 215
 heart rate, 179
 heat shock protein, 186, 195, 232, 251
 height, 18, 55
 helium, 112
 helplessness, 203
 hematological, x, 86, 88, 91, 101, 104, 106
 hematologist, 92
 hematopoietic stem cells, 244, 245
 hematopoietic stem-cell transplant, 106, 111
 hemoglobin, viii, 27, 29
 hemophilia, 97
 hemorrhagic diathesis, 168, 249
 herbal intoxication, 173
 herbal medicine, 173
 herb-drug interactions, 174, 175
 hereditary, 97
 herpes, 106, 189, 192
 herpes simplex, 106, 192
 herpes zoster, 189
 high shine, 81
 high strength, 66
 history, 177, 214, 215, 216, 217, 224, 235
 HIV, 88, 89, 91, 92, 104, 105, 106, 154, 155, 156, 162, 163, 185, 187, 195
 HIV/AIDS, 88, 89, 91, 92, 105, 185
 HIV-1, 187, 195
 HM, 12, 23, 58, 107, 128, 164, 236
 home practice, 222
 homeostasis, 119, 211
 horizontal, 4, 19, 20, 33, 64, 84, 207, 208
 hormonal, 177
 host, 103, 140, 153
 hot flashes, 173, 174, 193
 HPA axis, 182, 210
 HSP70/HSPA, 170, 178, 206, 210, 211, 247

-
- human, 29, 40, 73, 80, 87, 103, 104, 105, 121, 123, 126, 130, 141, 154, 162, 175, 176, 186, 188, 189, 192, 193, 195, 196, 202, 203, 209, 210, 234, 237, 249, 250, 251, 252, 253, 254
- human behavior, 29, 195
- human body, 87
- human health, 192
- human immunodeficiency virus, 104, 105, 162
- human neutrophils, 254
- human subjects, 192
- humoral immunity, 188
- Hungary, 1, 12, 15, 165, 199, 243
- Hunter, 42, 104
- hybrid, 82, 141
- hybridization, 87
- hydrocortisone, 95
- hydrophobicity, 122, 161
- hydroxyapatite, 247
- hygiene, xii, 19, 65, 75, 76, 82, 90, 97, 98, 99, 100, 105, 120, 151, 152, 159, 160, 161, 162, 212
- hyperaemia, 155
- hyperaesthesia, 213
- hyperparathyroidism, 108
- hyperplasia, 108, 155
- hyperprolactinemia, 185
- hypersensitivity, 189, 229
- hypnosis, xiii, 22, 165, 166, 176, 177, 184, 185, 188, 190, 191, 193, 221, 225, 228, 229
- hypnotherapy, 186, 187, 188, 189, 231
- hypoaesthesia, 213
- hypoallergenic, 119, 126
- hypochondria, 175, 179
- hypodontia, 21, 23
- hypotension, 169
- hypothalamo-pituitary-adrenal axis, 187
- hypothalamus, 210
- hysteric, 179
- images, 178
- imagination, 176, 177, 178
- imidazoles, 157
- immediate, 20, 25, 132, 133, 140, 144, 205, 237
- immediate early genes, 205
- immediate load, 20, 133, 140, 237
- immediate loading, 133, 140
- immersion, xi, 89, 116, 121, 127, 157, 162
- immune, x, 66, 86, 87, 120, 163, 170, 176, 178, 183, 209, 210, 236, 244, 247
- immune reaction, 210
- immune response, 66, 163, 210, 236
- immune system, 87, 183
- immunity, 163, 188, 210
- immunoglobulin, 178
- immunohistochemistry, 108
- immunosuppressants, x, 86, 88, 96
- immunosuppression, x, 86, 87, 88, 90, 93, 97
- immunosuppressive, 88, 93, 96, 111, 175
- immunosuppressive drugs, 93, 96
- impact strength, xi, 116
- implant, viii, x, xiii, 2, 3, 4, 5, 7, 10, 11, 15, 16, 17, 18, 20, 21, 23, 24, 25, 26, 46, 86, 92, 94, 95, 97, 98, 108, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 144, 154, 160, 171, 189, 206, 219, 237, 243, 247, 251
- implant placement, viii, 10, 15, 16, 23, 25, 26
- implants, xi, xii, xiii, 4, 10, 11, 12, 13, 17, 19, 21, 23, 24, 25, 86, 97, 102, 105, 130, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 143, 147, 153, 154, 155, 168, 169, 170, 236, 243, 246, 247
- impregnated, xi, 115, 117, 121, 123, 124
- impregnation, 47
- impression, 73, 76, 79, 144, 209, 221
- improvements, 62, 130
- impulses, 168, 169, 204
- in vitro, 56, 58, 67, 74, 104, 123, 128, 148, 153, 157, 160, 162, 195, 246, 248, 252, 253, 254
- in vivo, 58, 127, 163, 191, 195, 254
- incidence, x, 10, 12, 86, 87, 88, 91, 92, 97, 98, 99, 100, 102, 106, 107, 108, 111, 122, 146, 161, 200, 208, 238
- incision, 20
- incisive ability, 145
- incisor, 25, 47, 72
- incisors, 23, 26, 72
- incorporation, x, 115, 121, 122, 127, 128, 247, 252
- indentation, 12, 238
- India, 143, 182, 239
- indication, ix, x, 8, 9, 61, 63, 65, 75, 78, 82, 85, 87, 92, 101, 117, 203
- indirect retainer, 8, 9
- individual differences, 188, 193

I

- IASP, 237
- ibuprofen, 22
- ID, 42, 162
- ideal, 117, 185, 207
- identification, 87, 91, 103, 104, 155, 163, 222
- identity, 176
- idiopathic, 24, 194, 205, 238
- IgA, 94
- IgG, 94
- ill fitting dentures, 146, 148, 149, 206
- illness-related depression, 180
- image, 22, 131, 176, 183, 201, 203, 221, 228, 239
- imagery, 40, 180, 188

individualization, 82
 individuals, viii, xi, 3, 10, 27, 28, 41, 87, 88, 97, 98, 99, 129, 132, 156, 218
 induction, 24, 106, 111, 176, 184, 193, 248, 251
 industrialized countries, 138, 159
 industry, xii, 143
 ineffectiveness, 106
 inefficiency, 132
 infancy, 202
 infection, x, xii, 86, 87, 88, 89, 90, 91, 93, 100, 104, 105, 111, 112, 120, 151, 152, 154, 156, 157, 160, 162, 163, 175, 192
 infection control, 127, 175
 infectious, xii, 87, 88, 90, 99, 101, 102, 106, 151, 152, 180
 infectious agents, 88
 infectious disorders, 180
 inflammation, 76, 108, 155, 169, 170, 171, 191, 193, 210, 211
 inflammations, 210, 211, 212
 inflammatory, 4, 96, 171, 174, 177, 209, 210, 232, 245, 246, 251, 252
 inflammatory disease, 96, 232, 251
 information seeking, 220
 informed consent, 29
 infrared spectroscopy, viii, 27, 28, 29, 30
 infrastructure, 132
 ingestion, 149
 ingredients, 144, 145
 inhibition, 43, 130, 157, 223
 initiation, 48, 107
 injection, 218
 injured nerves, 247
 injuries, 10, 95
 injury, x, 86, 100, 189, 193, 197, 203, 245, 254
 inlay, ix, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59
 inorganic fillers, 54, 82
 insecurity, 203, 220
 insertion, xiii, 4, 5, 16, 18, 20, 21, 22, 52, 54, 74, 78, 146, 199, 211, 213, 219
 insomnia, 174, 177, 179
 instability, 134, 208
 institutions, xii, 151, 152, 160
 insulin, 164, 253
 integration, viii, 15, 16, 40, 229
 integrity, ix, 45, 46, 55, 87, 122, 248
 intelligence, 179
 intensity, 34, 57, 67, 71, 112, 118, 176, 181, 191, 206, 252
 interdental papilla, 76
 interdentally, 7
 interface, 48, 58, 117, 171

interference, 19, 174, 176, 181, 212
 interim restoration, 17
 interleukin, 245, 250
 internist, 173, 217
 internists, 171
 interocclusal, 25, 65, 70, 77
 interpersonal relations, 220
 interradicular bone, 20
 interrelations, 215
 intervention, 20, 138, 167, 190, 235, 240
 intoxication, 173, 195
 intraoral electrostimulation therapy, 169
 intravenously, 221
 intrusion, 18, 19, 20, 25, 64
 investment, 73, 80
 iodine, 111
 ion channels, 244
 ion transport, 244
 Iowa, 187, 229, 230, 233
 irradiation, 100, 112, 158, 162, 189, 193
 irritability, 178, 213
 irritation, 47, 82, 119, 146
 irritative, 166, 209, 214
 Israel, 171, 189
 issues, 65, 79, 107, 110, 220
 itching, 177, 179

J

Japan, 27, 41
 jaw, viii, xi, 27, 28, 29, 30, 31, 32, 34, 35, 37, 38, 40, 41, 42, 43, 65, 92, 102, 109, 112, 113, 130, 138, 140, 141, 168, 176, 181, 196, 201, 209, 238
 jaw closing, 35
 jaw functions, viii, 27, 28, 40, 41, 43
 jaw motion, viii, 27, 28, 29, 30, 31, 32, 34, 35, 37, 38, 40, 41
 jaw movement tracking, 29
 jaw movements, 42, 109, 168
 joint pain, 10, 171, 184, 187, 229, 237

K

keeping fast, 182
 keratinized soft tissue, 20
 keratoconjunctivitis, 107
 Korea, 10, 228

L

laboratory, ix, x, 57, 61, 62, 67, 74, 78, 79, 80, 83, 90, 92, 115, 116, 118, 119, 122, 135, 137, 159

laboratory tests, 92
 laceration, 92, 101
 lack of control, 220
 lactation, 173, 174
 lacunae, 248
 lacunar corrosion, 71
 laser, 112, 168, 170, 175, 183, 187, 189, 191, 192, 193, 195, 196
 lasers, 170, 189, 193
 laws, 68, 72
 leaching, 126
 lead, xii, 2, 16, 17, 19, 20, 22, 28, 54, 76, 77, 87, 131, 147, 151, 155, 166, 178, 183, 203, 205, 207, 208, 209, 211, 212, 216, 217, 218, 222, 223, 225, 245, 248
 leadership, 140
 learning, 40, 43, 167, 176, 179, 201, 202, 212, 221, 222
 learning process, 176, 201, 212, 221
 legs, 221
 lesions, 46, 92, 94, 95, 96, 97, 99, 101, 155, 160, 162, 206, 210, 212
 leukemia, 88, 92, 106
 leukoplakia, 97, 108
 libido, 167
 libido-theory, 167
 lichen, 97
 lidocaine, 96
 life expectancy, xii, 86, 97, 130, 143, 144, 147
 lifetime, 64, 75
 ligament, xiv, 20, 243, 248, 252, 253
 ligand, 174, 185, 245, 250
 ligands, 244, 248
 light, x, 18, 56, 82, 115, 118, 122, 125, 126, 177, 178, 179, 182, 184, 197
 light-cured composite, 82
 limbic system, 206
 lingual tooth reduction, 52
 lining epithelium, 210
 lithium, 47, 53, 57
 litigation, 211
 liver, x, 86, 173
 liver disease, x, 86
 living environment, 66
 load, xiv, 9, 47, 48, 53, 56, 59, 88, 90, 105, 117, 132, 133, 146, 166, 243, 248
 local anaesthesia, 221
 local anesthesia, 221
 local anesthetic, 99
 localization, 77, 201
 longevity, 55, 71, 75, 77, 78, 121
 longitudinal study, 25, 78, 84, 108, 135, 138, 140
 low risk, 130

lubrication, 74
 Luo, 157, 161
 luting agent, 48, 58
 lymph, 168, 209
 lymph node, 209
 lymph nodes, 209
 lymphocytes, 156, 209, 210, 247, 250, 252
 lysozyme, 178

M

machinable ceramics, 46
 magnesium, 174
 magnetic field, 196, 244, 250, 251, 252, 254
 magnetic resonance, 29, 43
 magnetic resonance imaging, 29, 43
 magnetoencephalography, 29
 magnets, 147, 186, 250
 magnitude, 134
 major depression, 174
 majority, 3, 101, 154, 156, 173, 218, 224, 225
 malignancy, 168, 169, 249
 malnutrition, 99, 154, 156
 malocclusion, 4, 23, 24, 26
 malodor, 212, 238
 malpositions, viii, 15, 16
 man, 163, 188, 195, 233, 234
 management, xii, 23, 25, 87, 99, 100, 101, 106, 108, 109, 110, 111, 143, 164, 183, 184, 185, 187, 189, 194, 200, 224, 225, 227, 228, 229, 230, 233, 238, 239
 mandible, 13, 92, 132, 136, 139, 140, 168, 175
 mandibular movement, 146, 148
 manipulation, 55, 119, 122, 202
 manufacturing, x, 57, 86
 margin, 54, 80
 marginal, ix, 4, 18, 45, 46, 54, 55, 56, 57, 58, 59, 67, 133
 marrow, 245
 martial art, 181
 mass, 73, 80
 massage, 168, 191
 masseter, 31, 169, 185, 186
 masseter muscle, 31, 185, 186
 mast, 175
 mast cells, 175
 mastication, xiv, 3, 40, 41, 42, 43, 47, 53, 70, 134, 148, 219, 243, 248
 masticatory, viii, xi, 7, 8, 9, 28, 29, 30, 31, 34, 35, 39, 40, 41, 42, 47, 67, 75, 98, 109, 130, 134, 135, 139, 140, 147, 168, 176, 181, 188, 201, 204, 205, 206, 207, 233, 234

- materials, ix, x, xii, 45, 46, 48, 53, 54, 58, 61, 66, 67, 68, 73, 74, 80, 81, 82, 85, 92, 95, 115, 116, 117, 118, 119, 120, 121, 122, 123, 125, 126, 143, 144, 145, 146, 147, 148, 153, 154, 158, 162, 163, 209, 210, 213, 239, 240, 252
- matrix, 47, 62, 73, 74, 116, 175, 197, 245, 246, 247, 248, 254
- matrix metalloproteinase, 175, 197
- matter, 3, 153, 183, 217, 219, 228
- maxilla, 12, 20, 23, 24, 175
- maxillary prognatism, 21
- maxillofacial prostheses, 144
- maxillomandibular relationship, 4, 208
- maximal opening position, 168
- maximum bite force, 40, 53, 135
- MB, 10, 12, 161, 193
- measurement, 73, 74, 80, 84
- measurements, 12, 54, 56, 73, 74
- mechanical properties, x, 66, 82, 115, 116, 117, 118, 119, 122, 124, 126
- median, 64, 74, 192
- medical, xii, 2, 7, 90, 102, 109, 147, 151, 153, 168, 171, 173, 180, 183, 187, 190, 193, 196, 212, 214, 215, 216, 217, 222, 224, 226, 230, 249
- medical care, 168, 190
- medical history, 102
- medical science, 183, 193
- medication, x, xiii, 86, 92, 99, 165, 166, 171, 172, 211, 217, 223, 225, 228
- medicinal herb, xiii, 165, 166, 173, 197, 222, 223, 225, 227, 228
- medicine, xiii, 10, 28, 153, 165, 166, 189, 192, 229
- meditation, xiii, 165, 166, 177, 178, 180, 181, 182, 184, 186, 188, 189, 192, 193, 196, 197, 225, 231
- MEG, 29
- mellitus, 106, 164
- memory, 29, 40, 41, 43, 99, 117, 174, 176
- memory function, 174
- menopausal, 187, 188, 191, 201, 237
- menstrual cycle, 174
- mental activity, 43
- mental disorder, 200
- mental energy, 177
- mental foramen, 132, 135
- mental health, 22, 177, 180, 181, 182, 201, 217, 222
- mental impairment, 41
- mercury, 214
- mesenchimal, 245, 254
- messengers, 205
- meta-analysis, 133, 139, 191
- metabolic syndrome, 93
- metabolism, 87, 171, 182
- metal framework, ix, 7, 8, 9, 10, 61, 62, 75, 76, 78
- metal ion, 209, 229, 236
- metal ions, 209, 229, 236
- metals, 84
- methacrylic acid, 122
- methodology, 4
- methyl cellulose, 144, 145
- methyl methacrylate, 119, 123, 125, 126
- methylene chloride, 118
- mice, 105
- microbiota, 103, 212
- micro-fillers, 82
- microflora, 87, 100, 111, 127, 154, 163, 164
- microleakage, 54
- micromassage, 169
- microorganism, xii, 121, 152, 154
- microorganisms, xii, 87, 92, 100, 120, 121, 151, 152, 156, 158, 159, 163
- microscopy, xii, 152, 155
- microstructure, 48, 58
- microwave, x, 115, 119, 120, 121, 122, 123, 124, 125, 126, 158, 159, 161, 162, 163
- migration, 175
- mind-body, vii, xiii, 22, 165, 166, 179, 180, 181, 182, 183, 222, 223, 224, 225, 227, 228
- miniscrew, 20
- Ministry of Education, 41
- minor connector, 29, 77
- minor salivary glands, 96, 211, 237
- misuse, xii, 144, 146, 147
- mitogen, 250
- mixing, 123
- mobility, 64, 75, 78, 212
- mode of fracture, 48
- models, 50, 92, 163, 184, 188
- modulus, 48, 66, 67, 69, 81, 82, 117, 121, 123, 124
- modulus of elasticity, 67, 81, 82, 117, 123
- moisture, 46
- molecules, 68, 157, 210, 211
- molybdenum, 20
- monomers, 116, 117, 119, 120
- mood, 166, 173, 174, 175, 180, 181, 182, 184, 188, 204, 207
- Moon, 124
- morbidity, 100
- morphology, 213
- mortality, xii, 28, 41, 100, 103, 151, 155, 157
- mortality rate, xii, 28, 151, 155, 157
- motivation, 166, 178, 201, 204, 222, 223, 225, 240
- motor abnormalities, 131
- motor activity, 204
- motor behavior, 42
- motor control, 29, 134
- motor skills, 99, 121, 140

mouth rinses, 157
mouthwashes, 111, 157, 158, 164
movement planning, 40
MR, 58, 110, 125, 126, 128, 139, 160, 162, 185
MRI, 29, 214
muco-dental support, 8
mucormycosis, x, 86
mucosa, x, 9, 12, 64, 75, 86, 87, 90, 91, 93, 94, 95, 96, 100, 101, 119, 121, 131, 132, 144, 153, 155, 170, 208, 209, 210, 212, 238, 247
mucosa ridging, 12, 208, 238
mucosal support, 8, 9
mucosal tissues, 7, 8
mucositis, 90, 92, 99, 100, 101, 106, 110, 111, 112
mucous membrane, 94, 106, 107, 145, 161
multiple myeloma, 92, 101, 113
multiple regression, 13
multiple regression analysis, 13
muscle, viii, 27, 28, 29, 30, 31, 34, 35, 37, 40, 41, 98, 134, 141, 168, 169, 171, 176, 179, 182, 204, 205, 207, 209
muscle atrophy, 134
muscle relaxant, 171
muscle relaxation, 169, 176, 181
muscles, 31, 32, 35, 37, 72, 77, 86, 168, 169, 185, 204, 205, 206, 207
muscular, 176, 204, 205, 207
musculoskeletal, 183, 194, 196, 228, 238
music, 185, 239
music therapy, 239
mutation, 109
mutations, 108
mutual agreement, 201
mutual trust, 220
mutuality, 220
myasthenia gravis, 144
myelodysplastic syndromes, 92
myeloma, 92, 101, 102, 113
myofascial pain, 169, 171, 172, 173, 176, 177, 187, 188, 193, 205, 207, 233, 238
myorelaxant, 172

N

narcissistic patients, 179
narcolepsy, 179
narratives, 139
nasopharyngeal carcinoma, 112
National Survey, 130
nausea, 213
Nd, 196
neck cancer, 215
necrosis, 20, 92, 101, 102, 245

necrotizing, 171
negative consequences, 68, 130
negative effects, ix, 61
negative thoughts, 167
neighboring teeth, 16, 18, 19
neon, 112
nerve, 10, 168, 169, 170, 174, 184, 185, 189, 193, 194, 197, 252, 253, 254
nervous structures, 203
neural, 29, 43, 182, 184
neuralgia, 189
neurobiology, 229
neurodegenerative, 98, 99
neurological, x, 28, 41, 85, 98, 213, 214, 217
neurological disease, x, 85, 98
neurologist, 217
neuromuscular coordination, 201
neuronal plasticity, 205
neurons, 42, 98, 176, 203, 205, 234
neuropathic pain, 196
neuropsychology, 185
neuroticism, 178, 182
neurotransmitter, 43
neurotransmitters, 205
neutral, 105
neutropenia, 90, 91, 93, 101
neutrophils, 156, 161
neutrophils, 96
nevus, 168, 249
New England, 103, 107, 110
NHANES, 153
nickel, 18, 19, 20, 212
nicotine, 156
nitrous oxide, 221
noble alloys, 66, 67
non-steroidal anti-inflammatory drugs, 171
norepinephrine, 169, 172, 206
normative evaluation, 183, 227
normatively determined need, 2, 218
North America, 123
NSAIDs, 171
nucleus, 169, 182, 187, 213, 233
nutrient, 163
nutrients, 184
nutritional, 40, 134, 156, 162, 163, 212
nutritional deficiencies, 156, 162
nutritional status, 40, 134, 163

O

obesity, 103
object loss, 201, 216
objectification, 178

obstacles, 167
 obstruction, 94
 obturators, 143, 144
 occlusal, viii, ix, 4, 8, 10, 12, 19, 24, 27, 28, 29, 33, 34, 38, 39, 40, 41, 45, 46, 49, 50, 52, 53, 57, 61, 64, 65, 67, 68, 69, 71, 72, 77, 79, 95, 117, 118, 139, 146, 166, 168, 169, 172, 185, 192, 194, 206, 207, 208, 228, 238
 occlusion, 2, 17, 22, 31, 77, 79, 146, 189, 207, 208, 218
 occupational stress, 223
 occupational therapy, 159
 ocular, 94, 96
 odor, 97
 oil, 145, 157, 191
 old age, 229, 236
 olfaction, 212
 open bite, 21
 operant conditioning, 201
 operations, 193
 ophthalmologist, 217
 opportunities, 103, 140
 optimal retention, 65, 66, 68, 71, 72
 optimism, 182
 optimization, 76
 oral acupuncture, 175
 oral candidiasis, 88, 91, 92, 93, 97, 105, 154, 155, 156, 162, 163
 oral cavity, x, 71, 74, 86, 87, 93, 94, 96, 97, 98, 99, 100, 101, 103, 110, 111, 119, 146, 147, 156, 158, 160, 164, 178, 210, 232, 251
 oral comfort, 13, 211
 oral health, 4, 10, 13, 22, 25, 98, 109, 130, 132, 134, 136, 139, 140, 141, 152, 158, 161, 163, 201, 203, 215, 226, 228, 230, 232, 235, 236, 240, 251
 oral health behavior, 201, 235
 oral lesion, xii, 93, 94, 96, 101, 102, 152
 oral stage, 203
 oral-sadistic stage, 203
 organ, 107, 108, 202, 204
 organic disease, 208
 organize, 226
 organs, 96, 204
 oropyrosis, 177
 orthodontic, viii, 4, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 170, 171, 195, 196, 203, 234, 248, 250, 254
 orthodontic treatment, 17, 21, 24, 25, 26
 orthodontics, vii, viii, 15, 16, 21, 23, 24
 orthognatic surgical procedure, 21
 osseointegrated, 17, 19, 20, 21, 24, 132, 138, 140, 236, 247
 osseointegration, 132

osseus, 25
 osteoarthritis, 170
 osteoblasts, 77, 101, 245, 246, 248, 250
 osteoclast, 175, 244, 245, 246, 248, 249, 250
 osteoclastogenesis, 244
 osteocyte, 252
 osteocytes, 248
 osteogenesis, 247, 251, 254
 osteoid, 248
 osteonecrosis of the jaw, 101, 102, 112, 113, 196
 osteoporosis, 101
 osteotomy, 189, 247, 251, 252
 otolaryngologist, 235
 outpatient, 194, 239
 outpatients, 239
 overcritical patients, 211, 236
 overcriticism, 211
 overdentures, x, xii, 86, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 143, 147, 154
 overextension, 208
 overlap, 82, 208
 overlay, 18
 overload, 7, 95, 205
 ownership, 222
 oxygen, 221

P

pacemaker, 168, 169, 180, 181, 182, 249
 pain perception, 177
 pain tolerance, 178, 204
 palate, xii, 21, 25, 26, 151, 153, 155, 156, 213
 palladium, ix, 62, 82
 palliative, xiii, 165, 166, 167, 168, 183, 223, 225, 228
 palpation, 175, 205
 pancreas, 96
 paradigm shift, 12
 parafunction, xiv, 169, 207, 208, 243, 248
 parafunctions, 207, 213
 parallel, 21, 64, 65, 68, 71, 78
 parallelism, 64
 paranoid psychosis, 175
 parasympathetic, 180
 parathyroid, 246, 252, 253
 parathyroid hormone, 246, 252, 253
 paravertebral, 168, 169, 249
 parents, 202
 parkinsonism, 144
 parotid, 100
 parotid gland, 100
 pathogenesis, 101, 160, 202, 235
 pathogens, 87, 104, 120, 152, 153, 163, 164

- pathology, 86, 94, 101
pathophysiology, 102, 183, 228, 237
pathways, 170, 182, 183, 206, 209, 212, 227, 244
patient management, 25, 200
patient satisfaction, xi, 13, 22, 130, 132, 134, 135, 136, 140, 141, 142, 207, 230, 240
PCR, 87, 91, 104, 155, 156, 164
pellicle, 128, 154, 160, 211
PEMF, vi, xiii, 22, 168, 170, 175, 188, 189, 195, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253
pemphigoid, 94, 95, 96, 106, 107
pemphigus, x, 86, 94, 95, 96, 106
pemphigus vulgaris, x, 86, 94, 95, 106
perceived control, 180
perception, 2, 29, 177, 218, 221
perceptions, 21, 176, 240
periapical, 210
periimplant, 246, 247
periodontal, ix, xiv, 7, 10, 18, 19, 20, 45, 46, 63, 64, 65, 67, 70, 72, 75, 77, 101, 106, 107, 154, 171, 192, 193, 194, 210, 211, 212, 231, 237, 243, 248, 249, 251, 252, 253
periodontal disease, 18, 64, 106, 107, 192, 212, 237
periodontitis, 154, 162, 211
peripheral neuropathy, 196
peripheral vascular disease, 174
peritoneal dialysis, 153, 160
peritonitis, 153, 160
permit, 70
peroxide, xi, 116
personality, 167, 188, 189, 200, 202, 204, 210, 222, 238, 239
PET, 41
phagocytosis, 156
phantom bite, 196, 207, 240
phantom tooth pain, 173, 206
pharmaceutical, 173
phenotype, 252
Philadelphia, 12, 13, 62, 183, 190, 228, 233, 239, 240
phobia, 176, 188
phonation, 211, 213
phosphate, 12, 253
photo-acoustic stimulation, xiii, 22, 165, 166, 178, 179, 184, 189, 221, 225, 231
physical and mechanical properties, 66, 78, 119, 122
physical fitness, 28, 41
physical properties, 81, 127
physical therapy, 240
physicians, 171
Physiological, 72
physiological factors, 72
physiology, 185, 229
physiotherapies, 22, 168, 223, 225, 228
physiotherapy, xiii, 165, 166, 168, 249
pilot drilling, 20
pilot study, 42, 57, 110, 112, 157, 162, 186, 189, 193, 251, 254
pin, 65, 84
pineal gland, 183
pituitary gland, 210
placebo, 107, 169, 170, 194, 195, 197, 223, 225, 228
placebo effect, 169, 170
plaque, viii, xii, 16, 19, 23, 67, 71, 78, 80, 107, 127, 151, 152, 160, 162, 163, 164, 211, 240
plastic deformation, 69, 123
plasticity, 80, 205
plasticizer, 116
platinum, 66, 82
playing, 248
pleasure, 202
PM, 57, 163, 164, 186, 190, 195, 235, 250, 254
PMS, 174
pneumonia, 109, 120
pocket, 19, 154, 211
pocket formation, 19, 211
polar, 122
policymakers, 136
polish, 74
polishing, xi, 73, 81, 82, 115, 122, 159
polymer, xi, 47, 48, 116, 117, 120, 122, 123
polymer matrix, 116
polymer networks, xi, 116
polymeric materials, 120
polymerization, x, 79, 80, 115, 119, 120, 122, 124, 125, 126
polymerization process, 119
polymerization temperature, 122
polymers, 116, 117, 119, 120, 122, 123, 124, 126
polymethylmethacrylate, 119
polymorphism, 113, 155, 184
polyphenols, 197
pontic, 7, 17, 18, 20, 47, 48, 52, 54, 55, 56
population, xi, 12, 28, 34, 41, 84, 88, 97, 98, 102, 104, 129, 130, 139, 143, 147, 153, 154, 156, 159, 161, 197, 209, 234, 238
population group, 130
porosity, 120, 122, 247
positive emotions, 221
positive mood, 180
post and core, 18, 19
postpolymerization, 124, 126
potential benefits, 109
prayer, xiii, 165, 166, 180, 185, 188
precancerosis, 168, 249

- precipitating life event, 215
precision attachment, 8, 9
precursor cells, 244
predictability, 42, 117
prefrontal cortex, viii, 27, 28, 29, 30, 33, 34, 40, 41, 42, 43
pregnancy, 173, 174, 180, 181, 182
prejudice, 220
premenstrual syndrome, 174
premolars, 16, 64, 72
preparation, viii, ix, x, 3, 4, 15, 16, 19, 23, 45, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 61, 62, 82, 85, 210, 219, 252
preservation, 57, 75, 76, 121
preservative, 145
pressing, 47, 122, 208
pressing technique, 47
pressure, 20, 57, 62, 65, 67, 68, 70, 71, 72, 81, 146, 169, 174, 195, 212, 238
presumptive, 214, 227
preventing, 3, 105, 117, 219, 245
prevention, xi, xiii, 11, 24, 28, 41, 86, 97, 100, 101, 103, 106, 111, 112, 129, 130, 158, 164, 165, 166, 183, 186, 197, 199, 212, 214, 217, 220, 222, 224, 227, 231, 251, 253
primary teeth, 58
primate, 42, 43
principles, viii, xiii, 7, 16, 23, 75, 77, 167, 196, 199
private practice, 12
probability, 78
probe, 30
problem behavior, 167
problem solving, 220
product performance, 145
professionals, 92, 101, 116, 122, 171, 173, 214, 217, 224, 226, 227
progesterone, 174
prognosis, 53, 117, 215, 216, 225
progressive muscle relaxation, 176, 181
progressive neurodegenerative disorder, 99
pro-inflammatory, 245
proliferation, 153, 155, 189, 209, 245, 246, 250, 252
promoter, 184, 252
pronunciation, 213
propagation, 48, 53
prophylactic, ix, 4, 61, 62, 67, 68, 71, 75, 76, 77, 79, 82
prophylaxis, 75, 76, 79, 97, 106
prostheses, xii, 4, 11, 29, 55, 59, 92, 93, 94, 95, 99, 100, 116, 121, 124, 125, 127, 132, 133, 134, 135, 136, 137, 138, 139, 141, 143, 144, 153, 160, 161, 190, 233, 234, 239
prosthesis, viii, xi, 9, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 64, 80, 86, 89, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 116, 117, 120, 122, 129, 131, 132, 133, 135, 136, 137, 138, 145, 183, 213, 227, 233
prosthetic device, 87, 90, 93, 97, 120
protease inhibitors, 105
protection, 87, 117, 122, 211
protein synthesis, 244, 251
proteins, 145, 160, 170, 178, 209, 210, 211, 232, 244, 247
proteomics, 232
prototype, 177, 250
protruded, 169
protrusion, 42, 209
provisional crown, 4, 17, 20
proximal boxes, 52
pruritus, 177, 179
pseudohalitus, 212
pseudoneurological, 177, 179, 213
psychiatric, 5, 166, 167, 175, 196, 211, 216, 219, 226
psychiatric disorders, 5, 196, 216, 219
psychiatric patients, 226
psychiatrist, xiii, 165, 166, 172, 173, 217, 223, 224, 226, 228
psychoanalysis, 167
psychoanalytic, 167
psychogenic, xiii, 3, 5, 22, 165, 166, 168, 173, 177, 178, 183, 187, 192, 194, 199, 200, 201, 202, 204, 206, 208, 211, 212, 213, 214, 215, 216, 219, 223, 225, 226, 227, 228, 231, 236, 238
psychological, viii, xiii, 2, 5, 7, 16, 22, 23, 24, 28, 86, 130, 165, 166, 167, 168, 174, 175, 176, 178, 187, 196, 199, 200, 202, 203, 205, 209, 212, 214, 217, 219, 221, 222, 223, 224, 228, 234, 240
psychological development, 203
psychological distress, 130, 205
psychological functions, 176, 196
psychological processes, 168, 200
psychological stress, 5, 175, 212, 217, 219, 221
psychology, 167, 187, 231, 235
psychometric properties, 228
psychopathology, 226, 234, 237
psychoses, 226
psychosis, 109, 175, 230
psychosocial, 2, 5, 16, 22, 131, 166, 174, 200, 203, 206, 208, 211, 213, 214, 215, 216, 217, 218, 219, 228, 235, 239
psychosocial factors, 22, 208
psychosocial functioning, 216
psychosocial stress, 174, 200, 206, 208, 213, 239
psychosomatic, vii, xiii, 10, 21, 22, 165, 166, 167, 168, 170, 171, 172, 173, 174, 176, 178, 179, 180,

181, 184, 186, 187, 192, 194, 199, 200, 201, 202, 204, 212, 213, 216, 223, 224, 225, 226, 227, 228, 229, 231, 237, 238, 239, 240
 psychosomatics, 24, 201
 psychotherapeutic, xiii, 22, 165, 166, 167, 176, 221, 223, 224, 225, 228
 psychotherapy, 167, 168, 184, 186, 192, 224, 226, 231
 psychotic, 179
 psychovegetative disorders, 173, 174
 ptyalismus, 211
 ptyaliorhea, 211
 public figures, xii, 143, 144
 public health, 130, 137
 public service, 131, 135
 pulp, 4, 16, 47, 203, 206, 210, 234, 237
 pulsed electromagnetic field, 22, 24, 168, 186, 188, 189, 192, 194, 195, 249, 250, 251, 252, 253, 254
 PVM, 145

Q

Qigong, xiii, 165, 167, 181, 189, 191
 quality of life, viii, 10, 22, 25, 27, 28, 41, 100, 103, 109, 112, 130, 131, 132, 134, 136, 138, 139, 141, 159, 161, 197, 201, 203, 228, 230, 236
 quantification, 87, 105
 quartz, 68
 quaternary ammonium, 128
 questionnaire, 12, 31, 33, 42, 110, 112, 218, 236, 240
 quetiapine, 99

R

radiation, 100, 110, 111, 112, 144, 163, 204
 Radiation, 111, 112, 204
 radiation therapy, 110
 radicals, 122
 radiography, 102
 radiotherapy, x, 86, 88, 92, 99, 100, 101, 110, 111, 112
 radius, 68
 RE, 10, 42, 111, 125, 195, 253
 reactions, 116, 119, 120, 125, 138, 166, 177, 209, 229, 235, 239
 reactive oxygen, 156, 246
 reactive oxygen species, 156, 246
 reactivity, 178, 193, 210, 234
 reading, 180
 reality, 3, 219, 221, 228
 recall, 147, 152
 receptor sites, 212
 receptors, 173, 244, 246, 254

recognition, 87, 214, 233
 recommendations, 122
 reconstruction, viii, 2, 5, 10, 19, 25, 28, 40, 41
 recovery, 75, 166, 171, 189, 193, 223, 225, 247
 referral, xiii, 30, 165, 166, 173, 194, 205, 217, 223, 224, 226, 228, 239
 referred pain, 204
 reflex splints, 170
 regenerate, 247
 regeneration, 19, 170, 189, 247, 252, 253
 regression, 179, 221
 regrowth, 154, 158
 regulatory influences, 233
 rehabilitation, viii, x, xi, 16, 21, 23, 62, 64, 75, 78, 83, 85, 86, 87, 93, 94, 95, 98, 99, 102, 103, 116, 117, 129, 131, 132, 136, 137, 144, 159, 230, 236, 238, 240
 reinforced composites, 46, 47
 reinforcement, 19, 53, 57, 116, 117, 118, 123, 124, 222
 relapses, 157, 224, 226
 relative analgesia, 221
 relaxation, xiii, 22, 119, 165, 166, 168, 169, 176, 177, 178, 179, 180, 184, 185, 188, 190, 191, 195, 196, 221
 relevance, xiv, 185, 202, 229, 244
 reliability, 73, 140
 relief, 10, 107, 211
 religiosity, 180, 195
 religious, 178, 180, 183, 202
 reline, 90, 95, 99, 120, 123, 126, 162
 relining, 77, 93
 remission, 96, 106, 193, 238
 renal failure, x, 86, 97, 108
 renal osteodystrophy, 108
 renal replacement therapy, 108
 repair, 77, 100, 116, 118, 123, 124, 147, 194, 251, 253
 replication, 121
 requirements, 79, 118, 134, 147
 researchers, 48, 116, 121
 residual alveolar ridge, 64, 80
 residual monomer, xi, 115, 116, 120, 121, 122, 123, 125, 126
 residual ridge, xii, xiii, 133, 134, 137, 143, 146, 170, 193, 243, 246, 253
 resilience, 64
 resilient, viii, 2, 6, 9, 64, 65, 95
 resin, xi, 8, 10, 46, 47, 48, 54, 55, 56, 57, 58, 59, 74, 76, 78, 80, 84, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 131, 132
 resins, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 161, 209

resistance, ix, 45, 47, 48, 52, 53, 55, 56, 57, 58, 66,
 67, 68, 71, 74, 81, 82, 87, 88, 103, 104, 105, 116,
 117, 121, 123, 156, 157, 159, 167, 169
 resolution, 89, 93, 101
 respiration, 121
 respiratory feedback, 179
 response, 4, 17, 19, 23, 25, 54, 86, 149, 183, 197,
 210, 229, 236, 246, 247
 responsiveness, 125
 rest position, 8, 77, 169
 restoration, ix, 7, 8, 9, 10, 16, 17, 18, 19, 20, 21, 23,
 24, 25, 45, 46, 47, 48, 53, 54, 55, 56, 59, 61, 62,
 63, 65, 75, 77, 79, 82, 83
 restorative dentistry, 23, 25, 26
 restorative material, 55
 restorative materials, 55
 restriction fragment length polymorphis, 155
 restrictions, 96, 97, 98
 retainer, 46, 48, 52, 54, 83
 retainers, ix, 29, 45, 46, 48, 55, 56
 retention, ix, xi, 8, 9, 18, 26, 45, 46, 47, 52, 55, 56,
 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73,
 74, 75, 76, 79, 80, 81, 82, 84, 97, 119, 131, 136,
 137, 143, 144, 145, 147, 148, 149, 158, 181, 226
 retention power, 62, 65
 retirement, 203
 retromolar areas, 175
 retronasal olfaction, 212
 RH, 185, 193, 234
 rheumatologist, 217
 riders, 76
 rigidity, 98
 rings, 70
 risk, xi, xii, 4, 5, 10, 22, 28, 40, 54, 87, 88, 91, 92,
 93, 101, 103, 107, 108, 110, 117, 130, 146, 151,
 152, 154, 173, 174, 175, 180, 184, 200, 215, 217,
 219, 222, 223, 227
 risk factors, 103, 108
 risks, 86
 RNA, 121
 root, 16, 18, 19, 20, 24, 26, 65, 127
 roots, 18, 19, 132, 167
 rotate, 52
 rotation axis, vii, 1, 2, 5, 6, 8, 9
 rotator cuff, 184
 rotator cuff tendinitis, 184
 roughening, 54
 roughness, xii, 55, 59, 68, 69, 72, 120, 126, 128, 152,
 154, 156, 158, 161, 162, 163
 rubber, 17, 105

S

sacral, 180
 saddle, 7, 8, 9, 78
 saddles, 78
 safety, 98, 191
 saliva, 72, 74, 87, 93, 96, 97, 123, 145, 147, 157,
 186, 189, 201, 204, 208, 211, 213, 232, 238, 251
 salivary, x, 86, 108, 128, 131, 134, 139, 142, 154,
 156, 157, 160, 161, 178, 179, 180, 211, 212, 231,
 232, 240, 251
 salivary gland, x, 86, 96, 142, 179, 211, 237, 240
 salivary glands, x, 86, 96, 142, 211, 237, 240
 salts, 145
 samarium, 186, 250
 savings, 71
 schizophrenia, 103, 109
 school, 182, 194, 216, 239
 science, 188, 196, 233, 234
 scientific papers, x, 85
 scope, xiii, 138, 165, 166, 171, 172, 223, 224, 226,
 228
 screws, 20
 secondary prophylaxis, 97
 secrete, 245
 secretion, 189, 201, 204, 208, 211, 213, 245, 248
 security, 119, 147
 sedative, 175
 seizure, 169, 180
 selective bonding, 54
 self confidence, 22
 self drilling, 20
 self image, 131, 176, 203
 self-confidence, 134
 self-control, 176
 self-esteem, 22, 23, 130, 220
 self-image, 22, 201, 203
 sensation, 10, 13, 119, 154, 204, 205, 211, 212, 237
 sense, 28, 77, 94, 98, 130, 167, 176, 202, 204
 sensitivity, 10, 12, 78, 183, 202
 sensitization, 119, 125, 176, 209
 sensory, 40, 131, 134, 147, 176, 201, 204, 213, 229
 Serbia, 61
 serology, 88
 serotonin, 169, 172, 173, 184, 185
 serum, 96, 147, 156, 182, 244, 248
 services, 141, 218, 222
 settle, vii, 1, 2, 5, 8, 9
 settlement, 8, 9, 71
 settling, vii, 1, 5, 8
 sexuality, 202, 207
 SFT, 139
 shamanism, 188

- shame, 130, 203
shape, 55, 63, 64, 65, 68, 69, 74, 76, 247
shear, 118, 123, 124, 125
shear strength, 123
shock, 82, 232
showing, xii, 33, 35, 56, 78, 91, 102, 134, 151, 155, 159
side effects, 87, 157, 166, 171
signal transduction, 244, 248
signals, 31, 174, 178, 179, 210
significance level, 33
signs, 88, 90, 119, 179
silane, 48, 54, 117, 123
silicones, 92, 95
silver, ix, xi, 62, 116, 121, 126, 127
Sinai, 192, 238
sinus arrhythmia, 190
skin, 31, 94, 169, 170, 171, 179, 195, 204, 247
sleep deprivation, xiii, 165, 167, 182, 184, 187
Slovakia, 161
smell, 212
social, viii, 2, 7, 12, 16, 22, 23, 99, 130, 131, 134, 178, 181, 184, 202, 203, 204, 216, 218, 220, 222, 227, 238
social interactions, 203
social learning, 221
social network, 216
social participation, 203
social relations, 130, 134
social relationships, 130
socialization, 208
sodium, 144, 145, 188
soft laser, 168, 170
software, 32
solid tumors, 104, 106
solubility, 116, 119, 126, 238
solution, xi, 4, 12, 64, 74, 76, 78, 89, 105, 130, 148, 207, 238
somatization, 214, 216
somatoform, 174, 177, 179, 185, 202, 229
sorption, 116, 119, 120
South Asia, 175
SP, 25, 58, 85, 115, 162, 163, 189, 196
Spain, 162
span, 4, 17, 20, 47, 48, 53, 55, 64, 173
spasm, 169, 171, 204, 205
species, xii, 87, 88, 100, 104, 111, 112, 126, 127, 151, 155, 156, 159, 160, 161, 163, 164, 246
speech, 3, 202, 207, 213, 219
spending, 130
spinal cord, 168, 169, 203, 205, 206, 249
spine, 182
spiritual, 180, 181, 182, 187, 196, 202, 231
spirituality, 195, 196
splint, 168, 169, 185, 239
splinted, 8, 135
splints, 124, 169, 185, 196
squamous cell, 112
Sri Lanka, 10
SS, 123, 127, 162, 192, 230
St John's wort, 173, 174, 188, 191, 192
stability, x, xi, 62, 66, 77, 79, 80, 82, 116, 118, 119, 131, 133, 134, 136, 143, 144, 145, 146, 147, 226
stabilization, ix, 18, 61, 62, 65, 67, 70, 75, 77, 159, 185
state, 54, 65, 71, 78, 105, 130, 141, 176, 177, 178, 179, 200, 222, 233
states, 177, 178, 186, 188, 192, 197, 210, 231
static friction coefficient, 66
steel, 18, 117
stem cells, 203, 234, 244, 245, 254
sterile, 157, 158
steroid inhaler, 154
stimulation, xiii, 22, 24, 32, 75, 165, 166, 168, 169, 174, 175, 177, 178, 179, 180, 184, 185, 186, 187, 189, 193, 196, 197, 209, 221, 225, 231, 232, 238, 243, 245, 246, 247, 248, 249, 250, 251, 253, 254
stimulus, 75, 221, 247
stomatitis, xii, 99, 108, 152, 159, 160, 161, 162, 163, 164
strategic extraction, 4, 219
streptococci, 154
Streptococcus, 121, 160
stress reactions, 180
stress response, 180, 195, 238
stressful life events, 218
stretch, 17, 168
stretching, 182
structural changes, 167
structural characteristics, 71
structure, ix, 29, 45, 46, 56, 65, 68, 73, 79, 86, 98, 99, 200, 207
stump, 4
style, 240
subgroups, 177
subjective evaluation, 183, 227
subjective experience, 176, 178
substitution, 232
substitutions, 215
substrates, 43, 253
success rate, 4, 133, 219, 227
sulfonylurea, 157
sulfur, 212
Sun, 236, 250
supereruption, 170
supernatural, 180

supervision, 145
 support, xii, 4, 7, 8, 9, 47, 49, 94, 131, 134, 135, 139, 143, 144, 146, 147, 151, 153, 154, 167, 222
 suppression, 147, 183, 185
 surface area, 19, 53, 55
 surface layer, 69
 surface protection, 117, 211
 surface roughness, xii, 59, 69, 126, 128, 152, 154, 161, 163
 surface tension, 72, 145, 146
 surface treatment, 55, 56, 69, 117, 119, 122, 124
 surgical intervention, ix, 3, 45, 46, 219, 224
 surplus, 180
 surveillance, 104, 177, 178
 survival, 55, 56, 78, 84, 87, 132, 133, 137, 138
 survival rate, 55, 56, 78, 138
 survivors, 112
 susceptibility, 67, 104, 156, 163, 196
 sweat, 96
 Switzerland, 151
 symbolic meaning, 183, 203
 symmetry, 247
 sympathetic, 179, 182, 210
 sympathetic nervous system, 210
 symptom chronification, 201, 214, 224
 symptom substitutions, 215
 synaptic transmission, 213
 syndrome, x, 10, 85, 96, 97, 107, 172, 173, 174, 184, 185, 188, 190, 191, 192, 193, 194, 195, 196, 206, 228, 229, 233, 234, 235, 236, 237, 238, 239, 240
 synergistic effect, 173, 174
 synthesis, 170, 205, 246
 systemic change, 87

T

tactile stimuli, 43
 Tai-Chi, xiii, 165, 167, 181
 target, 173, 222
 task performance, 31, 32, 34, 43
 taste, 97, 154, 204, 211, 212, 213, 229, 234, 237, 238, 239
 taste alterations, 212
 taste aversion, 229
 taste buds, 212
 taste perception, 211
 taxonomy, 237
 tea, 174, 175, 191, 192, 195, 196, 197
 team members, 217, 223
 technical failures, 77, 78, 84
 technician, ix, 62, 65, 81, 82, 117

techniques, x, 22, 29, 85, 86, 99, 115, 116, 118, 122, 124, 126, 145, 155, 159, 167, 168, 177, 178, 180, 181, 182, 193, 196, 220, 229
 technologies, 86, 104
 technology, 221
 telescope, ix, 8, 61, 62, 64, 65, 66, 68, 71, 72, 73, 74, 78, 83, 84
 telescopic, 68, 69, 74, 78, 82, 83, 84
 temperature, 120, 179, 204, 213
 temporal, 31, 42, 168, 169, 249
 temporary, x, 17, 19, 25, 79, 86
 temporomandibular, 170, 171, 187, 188, 191, 196, 205, 230, 233, 234, 235, 237, 238, 240
 temporomandibular disorders, 191, 230, 235
 tenderness, 174
 tendon, 170
 TENS, 168, 169, 188, 197
 tensile forces, 8
 tensile strength, 125
 tension, 68, 71, 72, 76, 80, 145, 146, 169, 172, 179, 207, 213, 248
 testing, 57, 126, 144, 163, 202, 234
 thalamus, 206
 therapeutic effects, 170
 therapeutic process, xiv, 243, 249
 therapeutic relationship, 194, 225
 therapeutics, xiii, 87, 165, 166, 224, 225, 228
 therapist, 167
 thermal expansion, 82
 thermal resistance, 121
 thickening agents, 144
 thickness, 48, 55, 80, 82, 146
 thoughts, 167
 thrombocytopenia, 90, 92
 thrombosis, 96
 thrush, 154
 time periods, xi, 116
 time resolution, 29
 tinnitus, 174, 213
 tipping, 4, 7, 9, 17
 tissue, ix, x, 18, 19, 20, 25, 45, 46, 54, 57, 62, 63, 64, 65, 66, 67, 70, 75, 77, 86, 92, 94, 95, 100, 121, 122, 127, 137, 144, 145, 149, 153, 155, 169, 171, 204, 208, 209, 211, 247, 250
 titanium, ix, 18, 19, 20, 62, 84, 121, 128, 154, 189, 250
 TNF, 245, 248
 TNF- α , 245, 248
 tongue, 12, 42, 72, 91, 119, 131, 134, 154, 202, 204, 207, 208, 212, 213, 238
 tonic, 174
 toothbrushing, 126
 toothpaste, xii, 152, 158

topical, xii, 93, 95, 96, 97, 151, 157
 topical anesthetic, 96
 torsion, 253
 total bonding technique, 54
 toxic substances, 116
 toxicity, 121
 toxicology, 121
 trabecular, 20
 traditional practices, 182
 traditions, 182
 trafficking, 209
 training, 78, 178, 181, 228
 traits, 195, 234
 transcendent, 180
 transcription, 205, 244, 254
 transcription factors, 205
 transcriptomics, 232
 transducer, 146
 transduction, 244
 transection, 197, 254
 transference, 168, 200, 221
 transformation, 48, 73
 transfusion, 92
 transient hypermobility, 17
 translation, 17, 244
 translucency, 46
 transmission, ix, 61, 65, 67, 174, 195, 250
 transparency, 80
 transplant, 97, 107, 108
 transplant recipients, 108
 transplantation, 88, 97, 106, 111
 trauma, 90, 92, 94, 95, 96, 100, 101, 154, 160, 179, 215
 treatment methods, 186
 trial, 31, 32, 33, 105, 110, 111, 136, 140, 141, 176, 181, 189, 194, 195, 196, 197, 230, 236, 240
 tribology, 83
 trigeminal, 169, 205, 233, 238
 trigeminal nerve, 169, 205
 trigger point, 175, 183, 205, 228
 triggers, 131
 tuberal region, 20
 tumor, 101, 245
 tumor necrosis factor, 245
 tumors, 86, 101

U

UK, 125, 236, 239
 ulcerations, 97, 170, 171
 ulcerative colitis, 191
 ultrasound, 168, 169
 uncomfortable feelings, 207

unconscious, 3, 22, 168, 189, 200, 219
 undercut, 52
 undercuts, 64, 80, 147
 underextension, 208
 underlying tissues, 147
 undermined enamel, 50
 undernutrition, 156
 uniform, 77
 United, 163, 196
 United States, 163, 196
 updating, 122
 urethane, 118
 urinary tract, 127
 urinary tract infection, 127
 urine, 191
 USA, 43, 158, 192, 252, 253

V

vacuum, 57, 80, 158
 vacuum investment, 80
 validation, 24, 112
 valve, 153
 variables, 55, 120, 121, 122, 125, 146, 194, 235, 253
 variations, 72, 118, 123, 132
 vascular, 169, 174, 194, 204, 238, 246
 vasculitis, 96
 vaseline, 96
 vasoactive, 119
 velocity, 32, 37, 187
 veneer, 48, 58, 81, 84
 veneering, ix, 47, 48, 53, 54, 57, 58, 61, 66, 79, 81, 82
 veneers, 76, 81, 82
 vertical, 4, 18, 26, 32, 38, 54, 56, 64, 70, 77, 146, 148, 207, 208
 vertigo, 174
 vestibular, 21
 vestibule, 99
 vestibulum, 175
 vibration, 169, 250
 vibrations, 169
 video games, 221
 vigil, xiii, 165, 167, 182
 virtual reality technology, 221
 virulence, 106, 155
 virus, 97, 104, 105, 106, 156, 162
 virus infection, 106
 viscoelastic properties, 121, 127
 viscosity, 72, 144, 145, 146, 147
 vision, 94
 visual acuity, 152
 visual imagination, 178

visually induced seizure, 180
vital energy, 175, 181
vitality, 166, 181
vocals, 213
voice, 181, 213
volatile sulfur compounds, 212
vulnerability, 207

working memory, 40, 43
working stress, 223
workplace, 22
World Health Organization, 142
worldwide, 86
wound healing, 112, 170, 171, 193, 194, 211, 247, 253

W

warmth, 167, 220
Washington, 105
water, xi, 116, 123, 124, 126, 127, 144, 145, 146, 158, 183
water sorption, 123, 126
waveform, 247, 248
wavelength(s), 31, 170
waviness, 69
wax, 73, 80, 119
weakness, 65, 213
wear, 28, 47, 58, 67, 68, 69, 71, 72, 74, 81, 82, 126, 154, 226, 233
web, 178
weight loss, 187
welding, 67, 69
well fitting, 4, 145, 146
well-being, 174, 203
wetting, 117, 145
WHO, 130, 142
width, 18, 64, 79
wire, 17, 18, 19
wires, xi, 115, 123, 124
withdrawal, 67
workers, 156, 159

X

xerophthalmia, 96
xerostomia, 96, 98, 100, 101, 107, 108, 112, 154, 175, 211
X-ray, 92, 214

Y

yeast, xii, 90, 92, 104, 151, 152, 153, 155, 159, 160
Yoga, xiii, 165, 167, 182, 189, 195, 196
yogic tradition, 182
young adults, 96
yttrium, 47, 53, 57
yttrium-oxide, 47, 53

Z

zeolites, 121
zinc, 12, 121, 126, 145, 147
zircon, 47
zirconia, 46, 47, 48, 53, 56, 57, 154
zirconium, 58